

Infinite 3D medium, Isotropic Point Source, Linearly-Anisotropic Scattering

Gamma-2 Random Flight

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2020 Eugene d'Eon


www.eugenedeon.com/hitchhikers

Path Setup

Put a file at `~/hitchhikerpath` with the path to your hitchhiker repo so that these worksheets can find the MC data from the C++ simulations for verification

```
SetDirectory[Import["~/hitchhikerpath"]]
```

Notation



c - single-scattering albedo

Σ_t - extinction coefficient

r - radial position coordinate in medium (distance from point source at origin)

$u = \cos \theta$ - direction cosine

b - anisotropy parameter

Namespace

```
In[9393]:= Begin["inf3DisopointlinanisoscatterGamma2`"]
```

```
Out[9393]= inf3DisopointlinanisoscatterGamma2`
```

Analytical results

Collision rate density

collision rate density C_c due to correlated emission:

derivation

```
In[9202]:= f00 = Fpc[0, 0, Exp[-#] # &];
f01 = Fpc[0, 1, Exp[-#] # &];
f11 = Fpc[1, 1, Exp[-#] # &];
o = 2;
Clear[A, b, c, r, h];
A[0] := 1; A[1] := b;
hsystem = Table[
  h[k] ==  $\frac{2}{\pi i} c u F[k, 0] + c \text{Sum}[A[m] \times h[m] \times F[k, m], \{m, 0, o-1\}], \{k, 0, o-1\}];
hsystemsolve = Simplify[Solve[hsystem, Table[h[i], {i, 0, o-1}]] /. A[1] → b /.
  F[0, 0] → f00 /. F[0, 1] → f01 /. F[1, 1] → f11 /. F[1, 0] → -f01]$ 
```

```
Out[9209]= { {h[0] →
  (2 c u (b c u^2 + u^4 - b c (1 + u^2) ArcTan[u]^2)) / (π (u^4 (1 - c + u^2) + b c u^2 (2 - c + u^2) -
  2 b c u (1 + u^2) ArcTan[u] + b c^2 (1 + u^2) ArcTan[u]^2)),
  h[1] → (2 c u^3 (-u + (1 + u^2) ArcTan[u])) / (π (u^4 (1 - c + u^2) + b c u^2 (2 - c + u^2) -
  2 b c u (1 + u^2) ArcTan[u] + b c^2 (1 + u^2) ArcTan[u]^2)) } }
```

```
In[9193]:= Clear[r];
(2 k + 1)  $\frac{1}{4 \pi i r c}$  (h[k]) j2[k, r u] /. k → 0 /. hsystemsolve // FullSimplify
```

```
Out[9193]= { (u (b c u^2 + u^4 - b c (1 + u^2) ArcTan[u]^2) Sin[r u]) /
  (2 π^2 r (-b (-2 + c) c u^2 + (1 + (-1 + b) c) u^4 +
  u^6 + b c (1 + u^2) ArcTan[u] (-2 u + c ArcTan[u]))) }
```

result

```
In[9437]:= Ccexact[r_, t_, c_, b_] :=
  NIntegrate[(u (b c u^2 + u^4 - b c (1 + u^2) ArcTan[u]^2) Sin[r u]) /
    (2 π^2 r (-b (-2 + c) c u^2 + (1 + (-1 + b) c) u^4 + u^6 + b c (1 + u^2) ArcTan[u]
      (-2 u + c ArcTan[u]))), {u, 0, Infinity}, Method -> "LevinRule"]

In[9159]:= TraditionalForm[
  HoldForm[Cc[r] = Integrate[(u (b c u^2 + u^4 - b c (1 + u^2) ArcTan[u]^2) Sin[r u]) /
    (2 π^2 r (-b (-2 + c) c u^2 + (1 + (-1 + b) c) u^4 + u^6 +
      b c (1 + u^2) ArcTan[u] (-2 u + c ArcTan[u]))), {u, 0, Infinity}]]]

Out[9159]/TraditionalForm=
```

$$C_c(r) = \int_0^{\infty} \frac{u (b c u^2 + u^4 - b c (1 + u^2) \tan^{-1}(u)^2) \sin(r u)}{2 \pi^2 r (-b (-2 + c) c u^2 + (1 + (-1 + b) c) u^4 + u^6 + b c (1 + u^2) \tan^{-1}(u) (-2 u + c \tan^{-1}(u)))} du$$

load MC data

```
In[9438]:= ppoints[xs_, dr_, maxx_] :=
  Table[{dr (i) - 0.5 dr, xs[[i]]}, {i, 1, Length[xs]}][[1 ;; -2]]

In[9439]:= ppointsu[xs_, du_, Σt_] :=
  Table[{-1.0 + du (i) - 0.5 du, xs[[i]] / (2 Σt)}, {i, 1, Length[xs]}][[1 ;; -1]]

In[9440]:= fs = FileNames["code/3D_medium/infinite3Dmedium/Isotropicpointsource/MCdata/
  inf3D_isotropicpoint_linanisoscatter_gamma2C*"];

In[9441]:= index[x_] := Module[{data, c, mfp, b},
  data = Import[x, "Table"];
  mfp = data[[1, 13]];
  c = data[[2, 3]];
  b = data[[1, 16]];
  {c, mfp, b, data}];
simulations = index /@ fs;
cs = Union[#[[1]] & /@ simulations]

Out[9443]:= {0.01, 0.1, 0.3, 0.5, 0.7, 0.8, 0.9, 0.95, 0.99, 0.999}

In[9444]:= mfps = Union[#[[2]] & /@ simulations]

Out[9444]:= {0.3, 1}

In[9445]:= bs = Union[#[[3]] & /@ simulations]

Out[9445]:= {-0.9, 0.7}

In[9446]:= numcollorders = inf3Disopointlinanisoscatter`simulations[[1]][[-1]][[2, 13]];
```

Compare analytic and MC

Collision-rate density - Exact solution (1) comparison to MC

```
In[9447]:= {{ActionMenu["Set c", "c = "<>ToString[#]>=> (c = #;) & /@cs], Dynamic[c]},  
            {ActionMenu["Set mfp", "mfp = "<>ToString[#]>=> (mfp = #;) & /@mfps],  
              Dynamic[mfp]},  
            {ActionMenu["Set b", "b = "<>ToString[#]>=> (b = #;) & /@bs], Dynamic[b]}}
```


```
Out[9447]= {{Set c, 0.95}, {Set mfp, 1}, {Set b, -0.9}}
```

```

In[9457]:= data = SelectFirst[simulations, #[[1]] == c && #[[2]] == mfp && #[[3]] == b &][[4]];
maxr = data[[2, 5]];
dr = data[[2, 7]];
MCCollisionRate = ppoints[data[[4]], dr, maxr];
exact1CRShallow = Quiet[{#[[1]], 4 Pi #[[1]]^2 Ccexact[#[[1]], 1/mfp, c, b]}] & /@
  MCCollisionRate[[1 ;; 60]];
exact1CR = Quiet[{#[[1]], 4 Pi #[[1]]^2 Ccexact[#[[1]], 1/mfp, c, b]}] & /@
  MCCollisionRate[[61 ;; -1 ;; 10]];
plotϕshallow = Quiet[Show[
  ListPlot[MCCollisionRate[[1 ;; 60]],
    PlotRange → All, PlotStyle → PointSize[.01]],
  ListPlot[exact1CRShallow, PlotRange → All, Joined → True],
  Frame → True,
  FrameLabel -> {{4 π r^2 Cc"[r]}, {r,}}
]];
logplotϕ = Quiet[Show[
  ListLogPlot[MCCollisionRate, PlotRange → All, PlotStyle → PointSize[.01]],
  ListLogPlot[exact1CR, PlotRange → All, Joined → True],
  ListLogPlot[exact1CRShallow, PlotRange → All, Joined → True],
  Frame → True,
  FrameLabel -> {{4 π r^2 Cc"[r]}, {r,}}
]];
Show[GraphicsGrid[{{plotϕshallow, logplotϕ}}, ImageSize → 800],
  PlotLabel -> "Exact solution (1)\nInfinite 3D, isotropic point source,
    linearly-anisotropic scattering, Gamma-2 random flight -
    correlated emission\nCollision-rate density Cc[r], c = "<>
ToString[c]<> ", b = "<> ToString[b]]

```

Out[9465]=



Namespace

In[9392]:= **End[]**

Out[9392]= inf3DisopointlinanisoscatterGamma2`