

Double GGX NDF

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2020 Eugene d'Eon

www.eugenedeon.com/hitchhikers

notation

$u = \mathbf{m} \cdot \mathbf{n} = \cos[\theta_m]$

α = roughness

definitions and derivations

The NDF that is to GGX as GGX is to Beckmann is:

`DoubleGGX`D[u_, α_] :=`

$$\frac{\alpha^2 \left(1 - u^2 + e^{-\frac{u^2 \alpha^2}{-1+u^2}} (1 + u^2 (-1 + \alpha^2)) \operatorname{ExpIntegralEi}\left[\frac{u^2 \alpha^2}{-1+u^2}\right] \right)}{\pi (-1 + u^2)^3} \operatorname{HeavisideTheta}[u]$$

In[683]:= `DoubleGGX`σ[u_, α_] :=` $\frac{1}{2} \left(u + \sqrt{\pi} \operatorname{Abs}[u] \operatorname{HypergeometricU}\left[-\frac{1}{2}, 0, \left(-1 + \frac{1}{u^2}\right) \alpha^2\right] \right)$

In[2702]:= `DoubleGGX`Λ[u_, α_] :=` $\frac{1}{2} \left(-1 + \frac{\sqrt{\pi} \operatorname{Abs}[u] \operatorname{HypergeometricU}\left[-\frac{1}{2}, 0, \left(-1 + \frac{1}{u^2}\right) \alpha^2\right]}{u} \right)$

In[686]:= `(1 + DoubleGGX`Λ[u, α]) u == DoubleGGX`σ[u, α] // FullSimplify`

Out[686]= `True`

In[687]:= `FullSimplify[(DoubleGGX`Λ[u, α]) u == DoubleGGX`σ[-u, α],
Assumptions → 0 < α < 1 && -1 < u < 1]`

Out[687]= `True`

In[690]:= `FunctionExpand[HypergeometricU[-1/2, 0, (-1 + 1/u^2) α^2]]`

Out[690]=
$$-\frac{e^{\frac{1}{2} \left(-1 + \frac{1}{u^2}\right) \alpha^2} (-1 + u^2) \alpha^2 \operatorname{BesselK}\left[0, \frac{1}{2} \left(-1 + \frac{1}{u^2}\right) \alpha^2\right]}{2 \sqrt{\pi} u^2} -$$
$$\frac{e^{\frac{1}{2} \left(-1 + \frac{1}{u^2}\right) \alpha^2} (-1 + u^2) \alpha^2 \operatorname{BesselK}\left[1, \frac{1}{2} \left(-1 + \frac{1}{u^2}\right) \alpha^2\right]}{2 \sqrt{\pi} u^2}$$

In[2703]:= **FullSimplify**[DoubleGGX` Δ [u, $\frac{u}{\sqrt{1-u^2} x}$], Assumptions $\rightarrow 0 < u < 1 \&\& x > 0$]

Out[2703]= $\frac{1}{2} \left(-1 + \sqrt{\pi} \text{HypergeometricU}\left[-\frac{1}{2}, 0, \frac{1}{x^2}\right] \right)$

shape invariant f(x)

In[1233]:= **FullSimplify**[DoubleGGX` \mathcal{D} [u, α] $u^4 \alpha^2 / . u \rightarrow \frac{1}{\sqrt{1+x^2} \alpha^2}$,

Assumptions $\rightarrow 1 - \frac{1}{\sqrt{1+x^2} \alpha^2} > 0$]

Out[1233]= $-\frac{x^2 + e^{\frac{1}{x^2}} (1 + x^2) \text{ExpIntegralEi}\left[-\frac{1}{x^2}\right]}{\pi x^6}$

derivation

In[681]:= **Integrate**[Exp[- αB] GGX` \mathcal{D} [u, $\alpha / \sqrt{\alpha B}$],
{ αB , 0, Infinity}, Assumptions $\rightarrow 0 < u < 1 \&\& 0 < \alpha < 1$]

Out[681]= \$Aborted

In[682]:= **Integrate**[Exp[- αB] GGX` σ [u, $\alpha / \sqrt{\alpha B}$],
{ αB , 0, Infinity}, Assumptions $\rightarrow -1 < u < 1 \&\& 0 < \alpha < 1$]

Out[682]= $\frac{1}{2} \left(u + \sqrt{\pi} \text{Abs}[u] \text{HypergeometricU}\left[-\frac{1}{2}, 0, \left(-1 + \frac{1}{u^2}\right) \alpha^2\right] \right)$

In[684]:= **Integrate**[Exp[- αB] GGX` Δ [u, $\alpha / \sqrt{\alpha B}$],
{ αB , 0, Infinity}, Assumptions $\rightarrow -1 < u < 1 \&\& 0 < \alpha < 1$]

Out[684]= $\frac{1}{2} \left(-1 + \frac{\sqrt{\pi} \text{Abs}[u] \text{HypergeometricU}\left[-\frac{1}{2}, 0, \left(-1 + \frac{1}{u^2}\right) \alpha^2\right]}{u} \right)$

distribution of slopes

In[692]:= **FullSimplify**[DoubleGGX` \mathcal{D} [$\frac{1}{\sqrt{p^2 + q^2 + 1}}$, α] $\left(\frac{1}{\sqrt{p^2 + q^2 + 1}} \right)^4$,

Assumptions $\rightarrow 0 < \alpha < 1 \&\& p > 0 \&\& q > 0$]

Out[692]= $-\frac{\alpha^2 \left(p^2 + q^2 + e^{\frac{\alpha^2}{p^2 + q^2}} (p^2 + q^2 + \alpha^2) \text{ExpIntegralEi}\left[-\frac{\alpha^2}{p^2 + q^2}\right] \right)}{\pi (p^2 + q^2)^3}$

In[695]:= DoubleGGX`P22[p_, q_, α] := $-\frac{\alpha^2 \left(p^2 + q^2 + e^{\frac{\alpha^2}{p^2 + q^2}} (p^2 + q^2 + \alpha^2) \text{ExpIntegralEi}\left[-\frac{\alpha^2}{p^2 + q^2}\right] \right)}{\pi (p^2 + q^2)^3}$

```
In[696]:= Integrate[DoubleGGX`P22[p, q, α], {p, -Infinity, Infinity},
               {q, -Infinity, Infinity}, Assumptions → 0 < α < 1]
```

```
Out[696]= 1
```


```
In[698]:= Integrate[DoubleGGX`P22[p, q, 1],
               {q, -Infinity, Infinity}, Assumptions → α > 0 && Im[p] == 0]
```

```
Out[698]= $Aborted
```

compare σ to delta integral:

```
In[699]:= Delta`σ[u_, ui_] := Re[2 ( √(1 - u² - ui²) + u ui ArcCos[ - (u ui) / (√(1 - u²) √(1 - ui²)) ] )]
```

```
In[705]:= With[{α = .7},
  Plot[{
    Quiet[NIntegrate[DoubleGGX`D[ui, α] × Delta`σ[u, ui], {ui, 0, 1}]],
    DoubleGGX`σ[u, α]
  }, {u, -1, 1}]
```



importance sampling

```

In[4489]:= With[{α = .9},
  Show[
    Histogram[Table[ $\sqrt{\frac{1}{1 - \left(\alpha / \sqrt{\#}\right)^2 + \frac{(\alpha / \sqrt{\#})^2}{\text{RandomReal}[]}}}$  & [-Log[RandomReal[]]]],
    {i, Range[10 000]}], 200, "PDF"],
    Plot[DoubleGGX`D[u, α] 2 Pi u, {u, 0, .999}, PlotRange → All]
  ]
]

```

