

Ei NDF

This is code to accompany the book:

A Hitchhiker's Guide to Multiple Scattering

© 2020 Eugene d'Eon

www.eugenedeon.com/hitchhikers

notation

$u = \mathbf{m} \cdot \mathbf{n} = \cos[\theta_m]$

α = roughness

Definitions and derivations

$$\text{In}[4444]:= \text{Ei}^{\text{D}}[u_, \alpha_] := \frac{\text{Gamma}\left[0, \frac{-1 + \frac{1}{u^2}}{\alpha^2}\right]}{\pi u^4 \alpha^2} \text{HeavisideTheta}[u]$$

$$\text{In}[2623]:= \text{Ei}^{\text{L}}[u_, \alpha_] := \frac{2 e^{\frac{u^2}{(-1+u^2)\alpha^2}} \sqrt{1-u^2} \alpha (-\alpha^2 + u^2 (-1 + \alpha^2)) + \sqrt{\pi} u (3 \alpha^2 + u^2 (2 - 3 \alpha^2)) \text{Erfc}\left[\frac{u}{\sqrt{1-u^2} \alpha}\right]}{6 \sqrt{\pi} u (-1 + u^2) \alpha^2}$$

$$\text{In}[1331]:= \text{Ei}^{\text{S}}[u_, \alpha_] := \frac{1}{6 \sqrt{\pi} (-1 + u^2) \alpha^2} \left(\alpha \left(3 \sqrt{\pi} u (-1 + u^2) \alpha + 2 e^{\frac{u^2}{(-1+u^2)\alpha^2}} \sqrt{1-u^2} (-\alpha^2 + u^2 (-1 + \alpha^2)) \right) + 3 \sqrt{\pi} u (-\alpha^2 + u^2 (-2 + \alpha^2)) \text{Erf}\left[\frac{u}{\sqrt{1-u^2} \alpha}\right] + 2 \sqrt{\pi} u^2 \text{Abs}[u] \left(1 + 2 \text{Erf}\left[\frac{u^2 \sqrt{1-u^2}}{\alpha \text{Abs}[u] - u^2 \alpha \text{Abs}[u]}\right] \right) \right)$$

$$\text{In}[2624]:= \text{Ei}^{\text{G1}}[u_, a_] := \frac{1}{1 + \text{Ei}^{\text{L}}[u, a]}$$

$$\text{In}[2689]:= \text{FullSimplify}\left[\text{Ei}^{\text{L}}\left[u, \frac{u}{\sqrt{1-u^2} x}\right], \text{Assumptions} \rightarrow 0 < u < 1 \ \&\& \ x > 0\right]$$

$$\text{Out}[2689]= \frac{e^{-x^2} (1 + x^2)}{3 \sqrt{\pi} x} - \frac{1}{6} (3 + 2 x^2) \text{Erfc}[x]$$

derivation

$$\text{Beckmann`D}[u_, \alpha_] := \frac{e^{-1 + \frac{1}{\alpha^2} u^2}}{\alpha^2 \pi u^4} \text{HeavisideTheta}[u]$$

In[1304]:= `Integrate[Beckmann`D[u, $\alpha \sqrt{m}$], {m, 0, 1}, Assumptions $\rightarrow 0 < u < 1 \&\& \alpha > 0$]`

$$\text{Out[1304]} = \frac{\text{Gamma}\left[0, \frac{-1 + \frac{1}{\alpha^2} u^2}\right]}{\pi u^4 \alpha^2}$$

In[1326]:= `Integrate[Beckmann` σ [u, $\alpha \sqrt{m}$], {m, 0, 1}, Assumptions $\rightarrow -1 < u < 1 \&\& \alpha > 0$]`

$$\begin{aligned} \text{Out[1326]} = & \frac{1}{6 \sqrt{\pi} (-1 + u^2) \alpha^2} \left(\alpha \left(3 \sqrt{\pi} u (-1 + u^2) \alpha + 2 e^{\frac{u^2}{(-1+u^2) \alpha^2}} \sqrt{1 - u^2} (-\alpha^2 + u^2 (-1 + \alpha^2)) \right) \right) + \\ & 3 \sqrt{\pi} u (-\alpha^2 + u^2 (-2 + \alpha^2)) \text{Erf}\left[\frac{u}{\sqrt{1 - u^2} \alpha}\right] + \\ & 2 \sqrt{\pi} u^2 \text{Abs}[u] \left(1 + 2 \text{Erf}\left[\frac{u^2 \sqrt{1 - u^2}}{\alpha \text{Abs}[u] - u^2 \alpha \text{Abs}[u]}\right] \right) \end{aligned}$$

In[1308]:= `Integrate[Beckmann` Δ [u, $\alpha \sqrt{m}$], {m, 0, 1}, Assumptions $\rightarrow 0 < u < 1 \&\& \alpha > 0$]`

$$\begin{aligned} \text{Out[1308]} = & \frac{2 e^{\frac{u^2}{(-1+u^2) \alpha^2}} \sqrt{1 - u^2} \alpha (-\alpha^2 + u^2 (-1 + \alpha^2)) + \sqrt{\pi} u (3 \alpha^2 + u^2 (2 - 3 \alpha^2)) \text{Erfc}\left[\frac{u}{\sqrt{1 - u^2} \alpha}\right]}{6 \sqrt{\pi} u (-1 + u^2) \alpha^2} \end{aligned}$$

shape invariant f(x)

In[1314]:= `FullSimplify[Ei`D[u, α] u4 α^2 /. u -> $\frac{1}{\sqrt{1 + x^2} \alpha^2}$,`

$$\text{Assumptions} \rightarrow 1 - \frac{1}{\sqrt{1 + x^2} \alpha^2} > 0 \&\& x > 0 \&\& \alpha > 0]$$

$$\text{Out[1314]} = \frac{\text{Gamma}[0, x^2]}{\pi}$$

height field normalization

In[1315]:= `Integrate[2 Pi u Ei`D[u, α], {u, 0, 1}, Assumptions $\rightarrow 0 < \alpha < 1$]`

$$\text{Out[1315]} = 1$$


distribution of slopes

```
In[1316]:= FullSimplify[Ei`D[ $\frac{1}{\sqrt{p^2 + q^2 + 1}}$ ,  $\alpha$ ]]  $\left(\frac{1}{\sqrt{p^2 + q^2 + 1}}\right)^4$ ,
Assumptions  $\rightarrow 0 < \alpha < 1 \ \&\& p > 0 \ \&\& q > 0$ ]
Out[1316]:=  $\frac{\text{Gamma}\left[0, \frac{p^2 + q^2}{\alpha^2}\right]}{\pi \alpha^2}$ 
```

```
In[1318]:= Ei`P22[p_, q_,  $\alpha$ _] :=  $\frac{\text{Gamma}\left[0, \frac{p^2 + q^2}{\alpha^2}\right]}{\pi \alpha^2}$ 
In[1319]:= Integrate[Ei`P22[p, q,  $\alpha$ ], {p, -Infinity, Infinity},
{q, -Infinity, Infinity}, Assumptions  $\rightarrow 0 < \alpha < 1$ ]
Out[1319]:= 1
```

compare σ to delta integral:

```
In[1322]:= Delta` $\sigma$ [u_, ui_] := Re[ $2 \left( \sqrt{1 - u^2 - ui^2} + u ui \text{ArcCos}\left[-\frac{u ui}{\sqrt{1 - u^2} \sqrt{1 - ui^2}}\right] \right)$ ]
In[1332]:= With[{ $\alpha$  = .7},
Plot[{
Quiet[NIntegrate[Ei`D[ui,  $\alpha$ ]  $\times$  Delta` $\sigma$ [u, ui], {ui, 0, 1}]],
Quiet[Ei` $\sigma$ [u,  $\alpha$ ]]
}, {u, -1, 1}]
]
```



importance sampling

```
In[4500]:= With[{α = 1.7},
  Show[
    Histogram[Table[1 / (Sqrt[1 - (α Sqrt[RandomReal[]])^2] (Log[RandomReal[]]))],
      {i, Range[10 000]}], 200, "PDF"],
    Plot[Ei`D[u, α] 2 Pi u, {u, 0, 1}, PlotRange → All]
  ]
]
```

... **General:** $\text{Exp}[-8.29137 \times 10^8]$ is too small to represent as a normalized machine number; precision may be lost.

