

---

## Table of contents

### Velocity - Movement of Matter in Space and Time

#### Matter and Vacuum - Refraction, max. phase velocity, max. group velocity:

Lightspeed, group velocity max.:

$$\begin{aligned} c &= \frac{1}{\sqrt{\epsilon_0 \mu_0}} = \frac{|\mathcal{E}|}{|\mathcal{B}|} = \sqrt{v_{phase} \cdot v_{matter}} = \eta \sqrt{\epsilon \mu} = \eta \sqrt{\epsilon_0 \epsilon_r \mu_0 \mu_r} = \\ \eta \cdot c_{ph} &= \frac{\omega}{k} = \frac{\lambda}{\mathcal{T}} = \nu \lambda = \\ -\frac{d\lambda}{d\nu} \nu^2 &= \lambda^2 \frac{\mathcal{J}_\lambda}{\mathcal{J}_\nu} = \frac{\langle |\mathcal{S}| \rangle}{\langle w_{EM} \rangle} = v_{gr}^{max} \\ &\leq v_{ph} - \lambda \frac{d}{d\lambda} v_{ph} = \frac{d}{dk} \omega = v_{gr} \end{aligned} \tag{1}$$

where:

$\mathcal{J} \equiv$  Flux Density flow <sup>1</sup>

$\mathcal{S} \equiv$  Poynting-Vector intensity

$w_{EM} \equiv$  Energy Density electromagnetic

Refraction Index:

Refraction for identifying matter (property of matter):

$$\begin{aligned} \eta &\stackrel{opt.}{=} \frac{\sin(\alpha)}{\sin(\beta)} = \frac{\eta_\beta}{\eta_\alpha} \stackrel{prism}{=} \frac{\sin[\frac{1}{2}(\varpi + \varsigma_{min})]}{\sin[\frac{1}{2}\varpi]} \\ &\stackrel{phys.}{=} \sqrt{\epsilon_r \mu_r} = \frac{\lambda_\alpha}{\lambda_\beta} = \frac{v_{ph,\alpha}}{v_{ph,\beta}} = \frac{1}{2} \frac{\lambda}{d_l} \frac{n}{\cos(\beta)} = \sqrt{\left(\frac{n}{2} \frac{\lambda}{d_l}\right)^2 + \sin(\alpha)^2} \\ &\stackrel{dispersion}{=} \eta(\lambda) = \eta(\omega) \stackrel{anisotropy}{=} \eta(r) \end{aligned} \tag{2}$$

Vacuum  $\Leftrightarrow \eta = 1$ : [1]

$$f = FocalLength \quad D = RefractivePower(dioptry) \tag{3}$$

---

<sup>1</sup>Flux Density

---

$n \equiv$  order (“quantum”):

$$n = \frac{\nu}{c} \Delta s = \frac{\Delta \varphi}{2\pi} = 2\eta \frac{d_l}{\lambda} \sqrt{\eta^2 - \sin(\alpha)^2} = 2\eta \frac{d_l}{\lambda} \cos(\beta) = \frac{q}{e_0} \quad \in \mathbb{N} \quad (4)$$

$$v^2 = \left( \frac{\nu \lambda}{\eta} \right)^2 = \left( \frac{\omega}{k\eta} \right)^2 \quad (5)$$

$$\left( \frac{\eta}{c} \right)^2 = \epsilon \mu \quad \epsilon = \epsilon_0 \epsilon_r \quad \mu = \mu_0 \mu_r \quad (6)$$

where:

$d_l :=$  distance length

$\alpha :=$  irradiation angle, incoming radiation direction, from source/sender (Einstrahlungswinkel zur Normalen auf Grenzfläche des Mediums)

$\beta :=$  refraction angle for bented transmitted radiation  $A_t$  orientation in matter (Durchstrahlungswinkel zur Normalen aus Grenzfläche ins Medium)

[1] Brechkraft

## Index

Flux Density, [1](#)

group velocity, [1](#)

Lightspeed, [1](#)

phase velocity, [1](#)