

## Exercise and Homework Round 4

These exercises (except for the last) will be gone through on **Tuesday October 6th 12:15–14:00** in the exercise session. The last exercise is a homework which you should return via mycourses by **Tuesday October 13th at 12:00**.

### Exercise 1. (Gradient descent for scalar linear model)

Let us consider a scalar linear model

$$y = g x + r \tag{1}$$

and the corresponding least squares cost function

$$J(x) = (y - g x)^2. \tag{2}$$

- (a) Write down the gradient descent algorithm for minimizing  $J(x)$ .
- (b) Compute the optimal step size  $\gamma$  by minimizing  $J$  with respect to it.

### Exercise 2. (Gauss–Newton for scalar linear model)

Consider the Gauss–Newton method for the model in the previous exercise.

- (a) Show that the Gauss–Newton method converges for this model in a single step.
- (b) What is the relationship of the Gauss–Newton method with the optimal step size that we computed in the previous exercise?

### Exercise 3. (Implementation of Gauss–Newton)

Consider the model

$$y_n = g(\mathbf{x}) + r_n, \quad (3)$$

where  $n = 1, \dots, N$ ,  $r_n \sim \mathcal{N}(0, R)$ , and

$$g(\mathbf{x}) = \begin{bmatrix} \alpha \sqrt{x_1} \\ \beta \sqrt{x_2} \end{bmatrix}. \quad (4)$$

- (a) Write down the corresponding weighted least squares problem for this model.
- (b) Derive the Jacobian of  $g$  and write down the pseudo-code for the Gauss–Newton algorithm.
- (c) Simulate data from this model with suitable parameters and implement Gauss–Newton algorithm for minimizing the cost function.

### Homework 4 (DL Tuesday October 13th at 12:00)

Implement gradient descend algorithm to minimize  $J(x) = (1.1 - \sin(x))^2$ . Also empirically test the effect of the step size to the convergence speed.