

We will provide a brute-force proof of equation (8.125), which was shown in the text via symmetry arguments. Start with

$$\boldsymbol{\sigma} \cdot \hat{\mathbf{r}} = \begin{bmatrix} \cos \theta & e^{-i\phi} \sin \theta \\ e^{i\phi} \sin \theta & -\cos \theta \end{bmatrix} \quad (1)$$

and the definition of spin-angular functions (see eq (3.384))

$$\mathcal{Y}_l^{j=l\pm 1/2, m} = \frac{1}{\sqrt{2l+1}} \begin{bmatrix} \pm \sqrt{l \pm m + \frac{1}{2}} Y_l^{m-1/2}(\theta, \phi) \\ \sqrt{l \mp m + \frac{1}{2}} Y_l^{m+1/2}(\theta, \phi) \end{bmatrix} \quad (2)$$

When j , instead of l , is fixed, we have

$$\mathcal{Y}_{l=j+1/2}^{j, m} = \frac{1}{\sqrt{2j+2}} \begin{bmatrix} -\sqrt{j-m+1} Y_{j+1/2}^{m-1/2}(\theta, \phi) \\ \sqrt{j+m+1} Y_{j+1/2}^{m+1/2}(\theta, \phi) \end{bmatrix} \quad \mathcal{Y}_{l=j-1/2}^{j, m} = \frac{1}{\sqrt{2j}} \begin{bmatrix} \sqrt{j+m} Y_{j-1/2}^{m-1/2}(\theta, \phi) \\ \sqrt{j-m} Y_{j-1/2}^{m+1/2}(\theta, \phi) \end{bmatrix} \quad (3)$$

We wish to prove equation (8.125)

$$(\boldsymbol{\sigma} \cdot \hat{\mathbf{r}}) \mathcal{Y}_{l=j\mp 1/2}^{j, m}(\theta, \phi) = -\mathcal{Y}_{l=j\pm 1/2}^{j, m}(\theta, \phi) \quad (4)$$

But only one sign needs to be proved, since $(\boldsymbol{\sigma} \cdot \hat{\mathbf{r}})^2 = 1$ implies the other. Now we pick to prove

$$\begin{bmatrix} \cos \theta & e^{-i\phi} \sin \theta \\ e^{i\phi} \sin \theta & -\cos \theta \end{bmatrix} \begin{bmatrix} \sqrt{\frac{j+m}{2j}} Y_{j-1/2}^{m-1/2}(\theta, \phi) \\ \sqrt{\frac{j-m}{2j}} Y_{j-1/2}^{m+1/2}(\theta, \phi) \end{bmatrix} = - \begin{bmatrix} -\sqrt{\frac{j-m+1}{2j+2}} Y_{j+1/2}^{m-1/2}(\theta, \phi) \\ \sqrt{\frac{j+m+1}{2j+2}} Y_{j+1/2}^{m+1/2}(\theta, \phi) \end{bmatrix} \quad (5)$$

Our strategy is to keep writing (5) as equivalent identities, until the end where they are recognized as well known facts.

Expand (5) into component identities:

$$\cos \theta \sqrt{\frac{j+m}{2j}} Y_{j-1/2}^{m-1/2}(\theta, \phi) + e^{-i\phi} \sin \theta \sqrt{\frac{j-m}{2j}} Y_{j-1/2}^{m+1/2}(\theta, \phi) = \sqrt{\frac{j-m+1}{2j+2}} Y_{j+1/2}^{m-1/2}(\theta, \phi) \quad (6)$$

$$e^{i\phi} \sin \theta \sqrt{\frac{j+m}{2j}} Y_{j-1/2}^{m-1/2}(\theta, \phi) - \cos \theta \sqrt{\frac{j-m}{2j}} Y_{j-1/2}^{m+1/2}(\theta, \phi) = -\sqrt{\frac{j+m+1}{2j+2}} Y_{j+1/2}^{m+1/2}(\theta, \phi) \quad (7)$$

With the general spherical harmonics formula

$$Y_l^m(\theta, \phi) = C_l^m e^{im\phi} P_l^m(\cos \theta) \quad \text{where} \quad C_l^m = (-1)^m \sqrt{\frac{(l-m)!}{(l+m)!}} \sqrt{\frac{2l+1}{4}} \quad (8)$$

we can see in (6) and (7) the ϕ dependency is clearly equal on both sides. So now we need to prove

$$\cos \theta \sqrt{\frac{j+m}{2j}} C_{j-1/2}^{m-1/2} P_{j-1/2}^{m-1/2} + \sin \theta \sqrt{\frac{j-m}{2j}} C_{j-1/2}^{m+1/2} P_{j-1/2}^{m+1/2} = \sqrt{\frac{j-m+1}{2j+2}} C_{j+1/2}^{m-1/2} P_{j+1/2}^{m-1/2} \quad (9)$$

$$\sin \theta \sqrt{\frac{j+m}{2j}} C_{j-1/2}^{m-1/2} P_{j-1/2}^{m-1/2} - \cos \theta \sqrt{\frac{j-m}{2j}} C_{j-1/2}^{m+1/2} P_{j-1/2}^{m+1/2} = -\sqrt{\frac{j+m+1}{2j+2}} C_{j+1/2}^{m+1/2} P_{j+1/2}^{m+1/2} \quad (10)$$

Divide (9) and (10) by $C_{j-1/2}^{m-1/2}$, while noticing (8) gives

$$\frac{C_{j+1/2}^{m+1/2}}{C_{j-1/2}^{m-1/2}} = -\sqrt{\frac{(j-m-1)!}{(j+m)!}} \sqrt{\frac{(j+m-1)!}{(j-m)!}} = -\frac{1}{\sqrt{(j+m)(j-m)}} \quad (11)$$

$$\frac{C_{j+1/2}^{m-1/2}}{C_{j-1/2}^{m-1/2}} = \sqrt{\frac{(j-m+1)!}{(j+m)!}} \sqrt{2j+2} \sqrt{\frac{(j+m-1)!}{(j-m)!}} \frac{1}{\sqrt{2j}} = \sqrt{\frac{2j+2}{2j}} \sqrt{\frac{j-m+1}{j+m}} \quad (12)$$

$$\frac{C_{j+1/2}^{m+1/2}}{C_{j-1/2}^{m-1/2}} = -\sqrt{\frac{(j-m)!}{(j+m+1)!}} \sqrt{2j+2} \sqrt{\frac{(j+m-1)!}{(j-m)!}} \frac{1}{\sqrt{2j}} = -\sqrt{\frac{2j+2}{2j}} \frac{1}{\sqrt{(j+m+1)(j+m)}} \quad (13)$$

we end up with the equivalent claims

$$\cos \theta \sqrt{\frac{j+m}{2j}} P_{j-1/2}^{m-1/2} - \sin \theta \sqrt{\frac{j-m}{2j}} \frac{1}{\sqrt{(j+m)(j-m)}} P_{j-1/2}^{m+1/2} = \sqrt{\frac{j-m+1}{2j+2}} \sqrt{\frac{2j+2}{2j}} \sqrt{\frac{j-m+1}{j+m}} P_{j+1/2}^{m-1/2} \quad (14)$$

$$\sin \theta \sqrt{\frac{j+m}{2j}} P_{j-1/2}^{m-1/2} + \cos \theta \sqrt{\frac{j-m}{2j}} \frac{1}{\sqrt{(j+m)(j-m)}} P_{j-1/2}^{m+1/2} = \sqrt{\frac{j+m+1}{2j+2}} \sqrt{\frac{2j+2}{2j}} \frac{1}{\sqrt{(j+m+1)(j+m)}} P_{j+1/2}^{m+1/2} \quad (15)$$

which are simplified into

$$\cos \theta \sqrt{\frac{j+m}{2j}} P_{j-1/2}^{m-1/2} - \frac{\sin \theta}{\sqrt{2j(j+m)}} P_{j-1/2}^{m+1/2} = \frac{j-m+1}{\sqrt{2j(j+m)}} P_{j+1/2}^{m-1/2} \quad (16)$$

$$\sin \theta \sqrt{\frac{j+m}{2j}} P_{j-1/2}^{m-1/2} + \frac{\cos \theta}{\sqrt{2j(j+m)}} P_{j-1/2}^{m+1/2} = \frac{1}{\sqrt{2j(j+m)}} P_{j+1/2}^{m+1/2} \quad (17)$$

or,

$$\cos \theta (j+m) P_{j-1/2}^{m-1/2} - \sin \theta P_{j-1/2}^{m+1/2} = (j-m+1) P_{j+1/2}^{m-1/2} \quad (18)$$

$$\sin \theta (j+m) P_{j-1/2}^{m-1/2} + \cos \theta P_{j-1/2}^{m+1/2} = P_{j+1/2}^{m+1/2} \quad (19)$$

If we denote $l' = j - 1/2$, $m' = m - 1/2$, $x = \cos \theta$, these are equivalent to

$$x(l' + m' + 1) P_{l'}^{m'} - \sqrt{1 - x^2} P_{l'}^{m'+1} = (l' - m' + 1) P_{l'+1}^{m'} \quad (20)$$

$$\sqrt{1 - x^2} (l' + m' + 1) P_{l'}^{m'} + x P_{l'}^{m'+1} = P_{l'+1}^{m'+1} \quad (21)$$

But these are well known recurrence relations of the associated Legendre functions (for reference, see equation (2.5.23) and (2.5.22) in A. R. Edmonds, *Angular Momentum in Quantum Mechanics*, Princeton University Press, 2nd edition (1960)).