

Entrainment of low Mach number thermals in stratified atmospheres

EVAN H. ANDERS,^{1,2} DANIEL LECOANET,³ AND BENJAMIN P. BROWN^{1,2}

¹*Dept. Astrophysical & Planetary Sciences, University of Colorado – Boulder, Boulder, CO 80309, USA*

²*Laboratory for Atmospheric and Space Physics, Boulder, CO 80303, USA*

³*Stuff*

(Received October 19, 2018; Revised February 18, 2019; Accepted ??)

Submitted to ApJ

ABSTRACT

Keywords: hydrodynamics — turbulence — entrainment

1. INTRODUCTION

2. EXPERIMENT

We perform direct numerical simulations of the evolution of dry thermals by solving the fully compressible Navier-Stokes equations,

$$\frac{D \ln \rho}{Dt} = -\nabla \cdot (\mathbf{u}) \quad (1)$$

$$\frac{D \mathbf{u}}{Dt} = -\frac{1}{\rho} \nabla P + \mathbf{g} + \nabla \cdot (\bar{\bar{\boldsymbol{\Pi}}}), \quad (2)$$

$$\frac{DT}{Dt} + (\gamma - 1)T \nabla \cdot (\mathbf{u}) + \frac{1}{\rho c_V} \nabla \cdot (-\rho \chi \nabla T) = \frac{1}{\rho c_V} (\bar{\bar{\boldsymbol{\Pi}}} \cdot \nabla) \cdot \mathbf{u}, \quad (3)$$

3. RESULTS

4. DISCUSSION

This work was supported by NASA Headquarters under the NASA Earth and Space Science Fellowship Program – Grant 80NSSC18K1199. This work was additionally supported by NASA LWS grant number NNX16AC92G. Computations were conducted with support by the NASA High End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center on Pleiades with allocation GID s1647.

APPENDIX

A. TABLE OF SIMULATIONS