Schwarzschild or Ledoux: composition gradients are fragile

EVAN H. ANDERS,[1,2] ADAM S. JERMYN,[3,2] DANIEL LECOANET,[1,4,2] ADRIAN E. FRASER,[5,2] IMOGEN G. CRESSWELL,[6,2] AND
J. R. FUENTES[7]

[1]CIERA, Northwestern University, Evanston IL 60201, USA
[2]Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA
[3]Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA
[4]Department of Engineering Sciences and Applied Mathematics, Northwestern University, Evanston IL 60208, USA
[5]University of California, Santa Cruz, Santa Cruz, California 95064, U.S.A
[6]Department Astrophysical and Planetary Sciences & LASP, University of Colorado, Boulder, CO 80309, USA
[7]Department of Physics and McGill Space Institute, McGill University, 3600 rue University, Montreal, QC H3A 2T8, Canada

ABSTRACT

This will be an abstract.

Keywords: UAT keywords

1. INTRODUCTION

2. CENTRAL RESULT:

APPENDIX

Corresponding author: Evan H. Anders
evan.anders@northwestern.edu

A. TABLE OF SIMULATION PARAMETERS

Input parameters and summary statistics of the simulations presented in this work are shown in Table **??**.

REFERENCES