



Modeling the Modeler: A Normative Theory of Experimental Design

Evan Piermont

Royal Holloway, University of London



Fernando Payró

Universitat Autònoma de Barcelona

University of Surrey

The Twentieth Day of November in the Year of Our Lord Twenty Twenty Five



An experimenter seeks to learn about a subject:

- ◇ has a theory about how subjects makes choices, conditional on their type
 - ◇ $\text{type} =_{def}$ parameters of preference / beliefs / whatever drives choices
- ◇ provides experiments to the subject, observes outcome
 - ◇ $\text{experiment} =_{def}$ a choice problem, from which the subject's behavior is observed

Our paper: how should the experimenter's goals (and her theory of behavior) determine her valuation of experiments

There are different goals for different experimenters:

Information Maximizing: wishes to learn as much as possible about the subject, independently of what she learns

There are different goals for different experimenters:

Information Maximizing: wishes to learn as much as possible about the subject, independently of what she learns

Theory Testing: Gains more value from validating a hypothesis than from anything else

There are different goals for different experimenters:

Information Maximizing: wishes to learn as much as possible about the subject, independently of what she learns

Theory Testing: Gains more value from validating a hypothesis than from anything else

Profit Seeking: Will use her information about the subject to sell him something

There are different goals for different experimenters:

Information Maximizing: wishes to learn as much as possible about the subject, independently of what she learns

Theory Testing: Gains more value from validating a hypothesis than from anything else

Profit Seeking: Will use her information about the subject to sell him something

Deceitful: Uninterested in truth, only wishes to validate hypothesis

- ◇ We offer a framework unifying the many goals of experimentation
- ◇ We propose three normative principles for how to rank experiments
 - ◇ minimal rationality properties, independent of specific motivations
- ◇ We show that they imply a particular representation
 - ◇ Relates a experiment to the expected value of identification
 - ◇ Provides a recipe for choosing experiments
 - ◇ Distinguish between various ‘types’ of experimenters
 - ◇ Test to ensure experimenter does not have an ‘agenda’

- ◇ A space of choice parameters Θ
- ◇ A (costly) experiment A has
 - ◇ possible observable outcomes $\mathcal{P} = \{P_1, \dots, P_n\}$
 - ◇ cost on implementation $c \geq 0$
- ◇ Observing $P \in \mathcal{P}$ *partially identifies* a set of parameters:
 - ◇ $W_{A,P} \subseteq \Theta$ consistent with observation

Normative Principles



Structural Invariance: Prefer the least costly implementation of a given identification of parameters

Information Monotonicity: (Weakly) prefer experiments that induce sharper identification

Identification Separability: Consider only the elements of experiments that can be controlled

Expected Identification Value



These principles characterize *expected identification value* maximization

- ◇ Exists some τ : for $W \subseteq \Theta$, $\tau(W)$ is the value of identifying W
- ◇ Experiment (A, \mathcal{P}, c) is valued according to:

$$\sum_{P \in \mathcal{P}} \tau(W_{A,P}) \mu(W_{A,P}) - c$$

- ◇ where μ is the (exogenous) prior probability

Special Case: Entropy



$$\tau(W) = -\log(\mu(W))$$

- ◇ Value of experiment is expected reduction in entropy
- ◇ Exp is valued proportional to information gain
- ◇ ‘Information Maximizing’ experimenter

Special Case: Hypothesis Testing



$$\tau(W) = \begin{cases} 1 & \text{if } W \subseteq W^* \text{ or } W^* \subseteq W^c \\ 0 & \text{otherwise .} \end{cases}$$

- ◇ Hypothesis: the parameter lies in W^*
- ◇ Value of exp is the probability the hypothesis can be accepted or rejected
- ◇ ‘Theory Testing’ experimenter

Special Case: Actions



$$\tau(W) = \max_{\alpha \in \mathbb{A}} \int_{\theta \in W} \xi(\alpha, \theta) d\mu.$$

- ◇ The analyst will take action $\alpha \in \mathbb{A}$
- ◇ Utility of outcome depends on the parameter: $\xi(a, \theta)$
- ◇ Value of exp is expected value of conditionally optimal action
- ◇ ‘Profit Seeking’ experimenter



Experimental Paradigms




An **Experimental Paradigm** is (Z, E, T)

- ◇ Z is a set of **choice patterns**
- ◇ E is a set of **experiments**.
- ◇ T is the experimenters **theory of behavior**.

An **experiment** $e = (A, \mathcal{P})$ is:

- ◇ $A \subseteq Z$ is finite choice problem
- ◇ \mathcal{P} is a partition of A



- ◇ Represents observability constraints
- ◇ Allows for dynamic experiments, non-lab settings, etc

A **theory of behavior** $T = \langle \Theta, \Omega, \mu \rangle$:

- ◇ Θ is a set of types (choice parameters), each $\theta \in \Theta$ associated with a choice function

$$c_\theta : 2^Z \rightarrow Z \quad \text{such that} \quad c_\theta(A) \in A.$$

- ◇ Ω algebra of measurable sets of Θ
- ◇ μ prior over (Θ, Ω)

Given (A, \mathcal{P}) , define the *identified set*:

$$W_{A,P} = \{\theta \in \Theta \mid c_\theta(A) \in P\}$$

Observing $P \in \mathcal{P}$ identifies that the subject's type is in $W_{A,P}$

Concordant Exp. Paradigms



We call an experimental paradigm (Z, E, T) **concordant** if the following hold:

- (1) Any observable outcome can be measured
- (2) Any measurable hypothesis can be tested

Concordant Exp. Paradigms



We call an experimental paradigm (Z, E, T) **concordant** if the following hold:

(1) Any observable outcome can be measured

◇ For all $(A, \mathcal{P}) \in E$ and $P \in \mathcal{P}$, $W_{A,P} \in \Omega$

(2) Any measurable hypothesis can be tested

Concordant Exp. Paradigms



We call an experimental paradigm (Z, E, T) **concordant** if the following hold:

(1) Any observable outcome can be measured

◇ For all $(A, \mathcal{P}) \in E$ and $P \in \mathcal{P}$, $W_{A,P} \in \Omega$

(2) Any measurable hypothesis can be tested

◇ For all Ω -meas. partitions $\{W_1, \dots, W_n\}$ of Θ , there exists $(A, \mathcal{P}) \in E$:

$$\{W_{A,P} | P \in \mathcal{P}\} =_{\mu} \{W_1, \dots, W_n\}$$

($=_{\mu}$ means up to μ -measure 0 differences)

Example: Choice Functions



Let X be a finite set of alternatives, subject chooses element out of $A \subseteq X$

- ◇ $Z = X \times \dots \times X$
- ◇ $E = \{(A_1 \times \dots \times A_n, \mathcal{P}) \mid A_i \subseteq X, \mathcal{P} \text{ partitions } A_1 \times \dots \times A_n\}$
- ◇ Θ is all (strict) preference orders over X :

$$c_\theta(A_1 \times \dots \times A_n) = \prod_{i \leq n} \{x \in A_i \mid x \succ_\theta y, \text{ for all } y \in A_i\}$$

- ◇ Then for discrete Ω , any μ forms a concordant paradigm

Example: Choice Functions



Let X be a finite set of alternatives, subject chooses element out of $A \subseteq X$

- ◇ $Z = X \times \dots \times X$
- ◇ $E = \{(A_1 \times \dots \times A_n, \mathcal{P}) \mid A_i \subseteq X, \mathcal{P} \text{ partitions } A_1 \times \dots \times A_n\}$
- ◇ Θ is all (strict) preference orders over X :

$$c_\theta(A_1 \times \dots \times A_n) = \prod_{i \leq n} \{x \in A_i \mid x \succ_\theta y, \text{ for all } y \in A_i\}$$

- ◇ Then for discrete Ω , any μ forms a concordant paradigm
 - ◇ this is **not** true for weak preferences; no way to test for tie-breaking

Example: Choice Correspondences



Let X be a finite set of alternatives, subject chooses subset out of $A \subseteq X$

- ◇ $Z = 2^X \times \dots \times 2^X$
- ◇ $E = \{(2^{Y_1} \times \dots \times 2^{Y_n}, \mathcal{P}) \mid Y_i \subseteq X, \mathcal{P} \text{ partitions } 2^{Y_1} \times \dots \times 2^{Y_n}\}$
- ◇ Θ is all preference orders over Z :

$$c_\theta(2^{Y_1} \times \dots \times 2^{Y_n}) = \prod_{i \leq n} \{x \in Y_i \mid x \succ_\theta y, \text{ for all } y \in Y_i\}$$

- ◇ Then for discrete Ω , any μ forms a concordant paradigm

Example: Adaptive Experiments



Let $X = \{a, b, c_a, c_b, d_a, d_b\}$

◇ $Z = \{(a, c_a), (a, d_a), (b, c_b), (b, d_b)\}$


◇ $E = \{(Z, \mathcal{P}) \mid \mathcal{P} \text{ partitions } Z\}$

◇ Θ collect utility functions over X : could represent naive subjects

$$c_\theta(Z) = (x, y) \text{ such that } x \in \arg \max_{z \in \{a, b\}} \theta(z) \text{ and } y \in \arg \max_{w \in \{c_x, d_x\}} \theta(w)$$

or sophisticated subjects

$$c_\theta(Z) = (x, y) \text{ such that } x \in \arg \max_{(z, w) \in Z} \theta(z) + \theta(w)$$



Example: Adaptive Experiments




Let $X = \{a, b, c_a, c_b, d_a, d_b\}$

◇ $Z = \{(a, c_a), (a, d_a), (b, c_b), (b, d_b)\}$

◇ $E = \{(Z, \mathcal{P}) \mid \mathcal{P} \text{ partitions } Z\}$

◇ In either case: Ω must reflect the inability to simultaneously measure preferences over $\{c_a, d_a\}$ and also $\{c_b, d_b\}$

◇ Ω must be generated by the four choice patterns in Z



Example: Expected Utility





- ◇ Z is lotteries over $\{a, b, c\}$
- ◇ E is all finite subsets of Z (and all partitions thereof)
- ◇ Θ is all affine functions over Z :


$$\theta(A) \in \arg \max_{x \in A} \theta(x)$$


(so each parameter also specifies how ties are broken)


$$\left\{\frac{2}{3}a + \frac{1}{3}b, \frac{2}{3}a + \frac{1}{3}c, \frac{1}{3}a + \frac{1}{3}b + \frac{1}{3}c\right\}$$
























Example: Expected Utility



- ◇ Z is lotteries over $\{a, b, c\}$
- ◇ E is all finite subsets of Z (and all partitions thereof)
- ◇ Θ is all affine functions over Z :

$$\theta(A) \in \arg \max_{x \in A} \theta(x)$$

Theorem: Random Expected Utility

$(Z, (\Theta, \Omega, \mu), E)$ is concordant iff μ is regular à la Gul & Pesendorfer (2006):

$$\text{for all } p, q \in Z, \quad \mu(\{\theta \mid \theta(p) = \theta(q)\}) = 0$$



Normative Principles



Primitive



- ◇ Fix some concordant paradigm: (Z, E, T) :
- ◇ Our primitive is a ranking \succsim over the set of all *costly* experiments:

$$(A, \mathcal{P}, c) \quad \text{where } (A, \mathcal{P}) \in E \quad \text{and} \quad c \geq 0$$

- ◇ We entertain 4 axioms:
 - ◇ 3 for our normative principles
 - ◇ 1 for quasi-linearity

“ Prefer the least costly implementation of a given identification of parameters ”



(P1) - Structural Invariance

Let $\{ W_{A,P} | P \in \mathcal{P} \} =_{\mu} \{ W_{B,Q} | Q \in \mathcal{Q} \}$:



$$(A, \mathcal{P}, c) \succsim (B, \mathcal{Q}, c') \quad \text{if and only if} \quad c' \geq c.$$

- ◇ Structural properties of experiments are irrelevant
- ◇ Also, 0-probability events are irrelevant ($=_{\mu}$)



Consider our EU maximizing subject choosing lotteries over $\{a, b, c\}$.





Consider our EU maximizing subject choosing lotteries over $\{a, b, c\}$.





Consider our EU maximizing subject choosing lotteries over $\{a, b, c\}$.



Consider our EU maximizing subject choosing lotteries over $\{a, b, c\}$.



Consider our EU maximizing subject choosing lotteries over $\{a, b, c\}$.



- ◇ Structural invariance reflects the symmetries of the given domain
- ◇ With linear utility, the symmetry is *translation invariance*:

Structural Invariance for Expected Utility

$$(A, \{P_1, \dots, P_n\}, c) \sim (A + B, \{P_1 + B, \dots, P_n + B\}, c)$$

“ (Weakly) prefer experiments that induce sharper identification ”

(P2) - Information Monotonicity

If \mathcal{P} refines \mathcal{Q} then $(A, \mathcal{P}, c) \succcurlyeq (A, \mathcal{Q}, c)$.

- ◇ Preference respects Blackwell order


“ Consider only the elements of experiments that can be controlled ”

(P3) - Identification Separability


$$(A, \mathcal{P}, c) \sim (A, \mathcal{Q}_B \mathcal{P}, 0) \quad \text{if and only if} \quad (A, \mathcal{P}_B \mathcal{Q}, c) \sim (A, \mathcal{Q}, 0).$$

- ◇ If \mathcal{P} and \mathcal{Q} partition of A and $B \subseteq A$, then $\mathcal{P}_B \mathcal{Q}$ denotes the partition that coincides with \mathcal{P} over B and with \mathcal{Q} over $A \setminus B$


(A, \mathcal{P})



(A, \mathcal{Q})




(A, \mathcal{P})




$(B, \mathcal{Q}_B \mathcal{P})$



(A, \mathcal{Q})




(A, \mathcal{P})



$(B, \mathcal{Q}_B \mathcal{P})$



$(A, \mathcal{P}_B \mathcal{Q})$




(A, \mathcal{Q})



$$(A, \mathcal{P}, c) \sim (A, \mathcal{Q}_B \mathcal{P}, 0) \quad \text{if and only if} \quad (A, \mathcal{P}_B \mathcal{Q}, c) \sim (A, \mathcal{Q}, 0).$$


(A, \mathcal{P})



$(B, \mathcal{Q}_B \mathcal{P})$



$(A, \mathcal{P}_B \mathcal{Q})$



(A, \mathcal{Q})



Let \succsim be an quasi-linear preference, represented by index $U : E \times \mathbb{R}_+ \rightarrow \mathbb{R}$.

Theorem: Expected Identification Value Maximization

Then \succsim satisfies P1-3 if and only if there exists a $\tau : \Omega \rightarrow \mathbb{R}$ such that:

$$U(A, \mathcal{P}, c) = \sum_{P \in \mathcal{P}} \tau(W_{A,P}) \mu(W_{A,P}) - c$$

with $W \subseteq V$ implies

- ◇ $\tau(V) \leq \tau(W) \mu(W|V) + \tau(V \setminus W) (1 - \mu(W|V))$
- ◇ if also $\mu(V \setminus W) = 0$ then $\tau(W) = \tau(V)$

Representation reflects our normative principles:

$$\sum_{P \in \mathcal{P}} \tau(W_{A,P}) \mu(W_{A,P})$$

- ◇ Only depends on $W_{A,P} \implies$ Structural Invariance
- ◇ Additive \implies Identification Separability
- ◇ $\tau(V) \leq \tau(W) \mu(W|V) + \tau(V \setminus W) (1 - \mu(W|V)) \implies$ Monotonicity



Information Based Models



Information Based Models



- ◇ Often, we seek only to reduce uncertainty
 - ◇ Experimenter does not have a preference for *what* is learned
- ◇ Value of identification $\tau(W)$, depends only of $\mu(W)$.
 - ◇ We can specialize structural invariance to capture this.
- ◇ E.g., value for an experiment is the (expected) reduction in entropy

$$U(A, \mathcal{P}, c) = - \sum_{P \in \mathcal{P}} \log(\mu(W_{A,P})) \mu(W_{A,P}) - c$$

Probability Vectors



For each $e = (A, \{P_1, \dots, P_n\}) \in E$ let

$$\mathbf{p}_e = \{\mu(W_{A,P_1}), \dots, \mu(W_{A,P_n})\} \in \Delta^n$$

denote the probability vector induced by the experiment.

(P2*) - Information Based Structural Invariance

Let $\mathbf{p}_e = \mathbf{p}_{e'}$ and $c' \geq c$: then

$$(e, c) \succcurlyeq (e', c'),$$

with a strict preference whenever $c' > c$.

(P2**) - Schur Concavity

Let $\mathbf{p}_e = \mathbf{D} \mathbf{p}_{e'}$ for some doubly stochastic matrix \mathbf{D} , and $c' \geq c$: then

$$(e, c) \succcurlyeq (e', c'),$$

with a strict preference whenever $c' > c$.

Let \succsim be an Expected Identification Value Maximization preference:

Theorem: Belief Based Models

Let μ be non-atomic. \succsim satisfies P2* if and only

$$U(A, \mathcal{P}, c) = \sum_{P \in \mathcal{P}} h(\mu(W_{A,P})) \mu(W_{A,P}) - c.$$

for some $h : [0, 1] \rightarrow \mathbb{R}$ with $h(0) = 0$ is such that

◇ For all $p, \alpha \in [0, 1]$,

$$h(p) \leq \alpha h(\alpha p) + (1 - \alpha) h((1 - \alpha)p)$$

◇ $p \mapsto h(p)p$ is concave iff P2* is strengthened to P2**.

Entropic Partitions

- ◇ Fix $(A, \mathcal{P} = \{P_1, \dots, P_n\})$ and let $\mathcal{P}^1 = \{P_1^1, \dots, P_k^1\}$ partition P_1 .
- ◇ Then $\mathcal{P}^\dagger = \{P_1^1, \dots, P_k^1, P_2, \dots, P_n\}$ is also partition of A .
 - ◇ As if observing \mathcal{P} and then if P_1 is realized, further observing \mathcal{P}^1
 - ◇ \mathcal{P} observed with prob 1, \mathcal{P}^1 observed with probability $\mu(W_{A, P_1})$
- ◇ Let $(B, \mathcal{Q} = \{Q_1, \dots, Q_k\})$ with $\mu(W_{B, Q_i}) = \mu(W_{A, P_i^1} \mid W_{A, P_1})$
 - ◇ Observing \mathcal{Q} has same ‘informational content’ as observing \mathcal{P}^1 conditional on realization of P_1

(A, \mathcal{P})

P_1	P_2	P_3	P_4
-------	-------	-------	-------

(A, \mathcal{P})

P_1	P_2	P_3	P_4
-------	-------	-------	-------

(A, \mathcal{P}^\dagger)

P_1^1	P_2^1	P_3^1	P_2	P_3	P_4
---------	---------	---------	-------	-------	-------

(A, \mathcal{P})

P_1	P_2	P_3	P_4
-------	-------	-------	-------

(A, \mathcal{P}^\dagger)

P_1^1	P_2^1	P_3^1	P_2	P_3	P_4
---------	---------	---------	-------	-------	-------

(B, \mathcal{Q})

Q_1	Q_2	Q_3
-------	-------	-------

(A, \mathcal{P})

P_1	P_2	P_3	P_4
-------	-------	-------	-------

(A, \mathcal{P}^\dagger)

P_1^1	P_2^1	P_3^1	P_2	P_3	P_4
---------	---------	---------	-------	-------	-------

(B, \mathcal{Q})

Q_1	Q_2	Q_3
-------	-------	-------

(A, \mathcal{P})

P_1	P_2	P_3	P_4
-------	-------	-------	-------


(A, \mathcal{P}^\dagger)

P_1^1	P_2^1	P_3^1	P_2	P_3	P_4
---------	---------	---------	-------	-------	-------


(B, \mathcal{Q})

Q_1	Q_2	Q_3
-------	-------	-------


(A, \mathcal{P})



(A, \mathcal{P}^\dagger)



(B, \mathcal{Q})



(P3*) - Identification Separability for Entropy

Fix $(A, \mathcal{P} = \{P_1, \dots, P_n\})$ and $\mathcal{P}^\dagger = \{P_1^1, \dots, P_k^1, P_2, \dots, P_n\}$.

Then if $(B, \{Q_1, \dots, Q_k\})$ is such that $\mu(W_{B, Q_i}) = \mu(W_{A, P_i^1} \mid W_{A, P_1})$, it follows that

$$(B, \mathcal{Q}, c) \sim (A, \{A\}, 0) \quad \text{if and only if} \quad (A, \mathcal{P}^\dagger, \mu(W_{A, P_1}) \cdot c) \sim (A, P, 0)$$

- ◇ Conditional on P_1 being realized, observing \mathcal{P}^1 has the same information content as (unconditionally) observing \mathcal{Q}
- ◇ P_1 is realized with probability $\mu(W_{A, P_1})$.

Let \succsim be an Expected Identification Value Maximization preference:

Theorem: Entropy Minimization

Let μ be non-atomic. \succsim satisfies P2** and P3* if and only

$$U(A, \mathcal{P}, c) = - \sum_{P \in \mathcal{P}} \log(\mu(W_{A,P})) \mu(W_{A,P}) - c.$$



Thank You!

