KAPASA MAKASA UNIVERSITY

GROUP 4

NAMES	SIN
CHISHIMBA SYDNEY JEREMIAH	22167595
NGWELE MICHEAL	20230382
MULONGOTI COSMAS	20231586
CHIBANGA BENJAMIN	20231800
ZULU FRANCISCO	20231750
WINA PRISCILLA	20232049
MWALE SARAH	20231693
NAKAPITE TRIGHT	20231925
SIAKAPOGO RACHEAL	20231605
MWINGA MARTHA	


INTRODUCTION

The Misamfu regional research station was established in 1957 by the government initiatives. It's part of Zambia Agriculture Research Institute (ZARI). Misamfu regional research station is located in Northern Province kasama Zambia. It is a prominent agriculture research institute located in Zambia, dedicated to advancing sustainable agriculture practices and enhancing food security in the region. The centers research initiatives encompasses a variety of topics including soil and water management, rice research system, plant protection and acquired resistance farming system. This report presents field-based descriptions of a soil pit at Misamfu Research Center, conducted without laboratory analysis or inclusion of organic horizons.

A soil profile refers to a vertical cross-section of soil, showcasing its layers or horizons. It provides valuable information about soil composition, structure, texture, color and organic matter content. Understanding soil profiles is key for sustainable agricultural development and land planning (FAO, 2006). In Kasama, Northern Zambia, the Misamfu Red Soil Series is a predominant soil type, recognized for its deep profile and characteristic red coloration (ZARI, 2023).

The Misamfu Red Soil Series is a notable soil formation. Due to the buildup of iron oxides, these soils are distinguished by their characteristic red hue. The parent minerals, mainly quartzite and schist, which have seen significant weathering over time, have a significant impact on the formation and properties of these soils. Because Misamfu soils are vital to the region's agricultural operations, it is imperative to comprehend their characteristics, constraints, and management techniques.

Comprehending the Misamfu Red Soil Series offers valuable perspectives on horizon differentiation, soil formation processes, and the physical and chemical elements influencing their potential for agriculture. The morphology, texture, structure, surface characteristics, and suitability for cultivation of the Misamfu Red Soil Series are the main topics of this report's thorough investigation of the soil profile.



OBJECTIVES

- To describe the morphological and physical characteristics of the Misamfu Red Soil Series.

- To reflect on the implications of these soil properties for agricultural practices.

- To examine soil features such as color, texture, structure, and crusting.

- To assess the limitations and agricultural implications of the Misamfu soil profile

- To propose sustainable management practices for enhancing soil productivity.

METHODOLOGY

- Soil pits were dug at regular intervals along the transect, exposing the vertical soil profile for detailed examination.


- Morphological characteristics were recorded, including horizon color (using the Munsell Soil Color Chart), texture (feel method), structure, and consistence.

SITE DESCRIPTION

The study site lies at Misamfu Research Center in Kasama District, a region characterized by high rainfall and deeply weathered soils derived from granitic and gneissic rock formations (Sanchez and Logan, 1992). The natural vegetation of the area was originally dominated by Miombo woodland species such as Brachystegia and Julbernardia (ZARI, 2023). The soils are classified as Ferralsols according to the World Reference Base (FAO, 2006).

RESULTS AND DISCUSSION

SOIL MORPHOLOGY



A series of discrete horizons define the Misamfu soil profile. The AP horizon has a sandy clay loam texture and is usually dark reddish-brown (5YR 3/4) at the surface. This horizon serves as the main area for plant growth since it is friable when wet and has a modest root density. Below this, the Bt1 horizon is identified by its clay texture, moderate sub angular blocky structure, and reddish-brown color (5YR 4/4). The presence of clay films, a certain sign of clay illuviation, makes this horizon noteworthy.

Deeper down, the Bt2 horizon has a thicker, clay-rich texture and a more intense red tint (2.5YR 4/6). Its firmness and sturdy blocky structure frequently prevent roots from penetrating. Given how narrow this horizon is, it might physically prevent roots from growing in deeper strata. The BC horizon at the profile's base is a transitional zone that gradually transitions from clay-rich

material to the parent material. Because of its weak blocky structure and sandy clay loam texture, this area is less limiting for root growth.

SURFACE FEATURES AND CRUSTING

The Misamfu Red Soil Series frequently exhibits surface crusting, especially in regions where the soil is exposed. There were two primary kinds of crusts found: depositional and structural. Raindrops hitting bare soil compacts the surface and decreases water uptake, forming structural crusts. In farmed fields when the soil surface is disturbed, these crusts frequently occur. Conversely, when fine particles are carried by runoff and land on the soil surface, depositional crusts form. Because it is more compact, this kind of crust can significantly restrict the emergence of seedlings. A number of variables, including soil texture, slope, vegetation cover, and rainfall intensity, affect the likelihood of crusting.

SOIL CONSTRAINTS AND AGRICULTURAL IMPLICATIONS

For agricultural development, the Misamfu soils pose a number of difficulties. The Misamfu soil profile supports agriculture due to its well-drained surface layer, but productivity can be hindered by poor drainage in deeper horizons. To optimize crop production, soil conservation practices like mulching and reduced tillage are recommended (FAO, 2006). Fertility management is also essential due to nutrient depletion through leaching in high rainfall areas (Sanchez and Logan, 1992).

The Bt2 horizon's compactness, which limits root growth and decreases water circulation, is the main limitation. This issue is made worse by surface crusting, which inhibits water infiltration and lowers seedling emergence. With pH values normally falling between 5.5 and 6.0, the soils are also somewhat acidic, which could have an impact on crops' access to nutrients. The clay concentration of the Misamfu soils, however, maintains their good nutrient-holding capacity in spite of these drawbacks. The Bt horizons' clay films are a definite sign of illuviation, which contributes fine clay particles and related nutrients to these layers. These soils may sustain a variety of crops, such as cassava, beans, and maize, with the right care.

MANAGEMENT RECOMMENDATIONS

The Misamfu Red Soil Series should be managed in a number of ways to increase its yield. Among these is the application of organic materials, such manure or compost, to enhance soil structure and moisture retention. To reduce soil disturbance and stop surface crusting, conservation tillage techniques ought to be used. Correcting the acidity of the soil may need liming, especially in regions where the pH is less than 5.5. Cover crops can also improve the amount of organic matter in the soil, stop erosion, and shield the soil's surface.

CONCLUSION

If properly maintained, the Misamfu Red Soil Series is a well-established, clay-rich soil type with substantial agricultural potential. Long-term weathering and clay illuviation are responsible for its characteristic red color and clay-rich Bt layers. But there are drawbacks to the soils as well, such as surface crusting and restricted root penetration in the Bt2 horizon. For soil fertility and productivity to be maintained, sustainable soil management techniques including liming, conservation tillage, and organic matter addition are crucial.

APPENDIX



- Soil Color References from Munsell Chart

- Field Texture Analysis Table

- Depth and Horizon Measurement Sheet

REFERENCES

1. FAO, 2006. Guidelines for Soil Description. 4th edition. Rome: Food and Agriculture Organization of the United Nations.
2. Sanchez, P.A. and Logan, T.J., 1992. Myths and science about the chemistry and fertility of soils in the tropics. In: Myths and Science of Soils of the Tropics, SSSA Special Publication No. 29. Madison: Soil Science Society of America, pp. 35–46.
3. ZARI (Zambia Agriculture Research Institute), 2023. Misamfu Research Station Soil Survey Manual. Lusaka: Ministry of Agriculture.
4. Fritsch, E. and Valentin, C., 1987. Characterization of a soil transect in Misamfu Research Station (Zambia). Africaland. Land development and management of acid soils in Africa. II. IBSRAM Proceedings, (7), pp.287-310.
5. MALAMA, C. and KONDOWE, K., 2003. Performance of green manures and grain legumes on severely acidic soils in northern Zambia, and their effect on soil fertility improvement. In Grain Legumes and Green Manures for Soil Fertility in Southern Africa: Taking Stock of Progress: Proceedings of a Conference Held 8-11 October 2002 at the Leopard Rock Hotel, Vumba, Zimbabwe (p. 185). CIMMYT.
6. Maynard, T.G., 1983. A partial budget analysis of the first six years of the long term fertilizer trial (MN. 1074) at Misamfu Regional Research Station [minimum returns analysis].
7. Sleen, L.V., 1976. Detailed soil survey of the Misamfu Regional Research Station, Northern Province.