

# Equilibrium States are (at least somewhat) Gibbs

Some new results on equilibrium states for amenable group subshifts.

C. Evans Hedges - University of Denver

Great Lakes Mathematical Physics Meeting: 07 June 2024

# Presentation Outline

- ▶ Introduction to  $\mathbb{Z}^d$  Subshifts and Symbolic Dynamics
- ▶ Introduction to Thermodynamic Formalism in the  $\mathbb{Z}$  Setting
  - ▶ Entropy
  - ▶ Pressure
  - ▶ Equilibrium States
- ▶ A jump to the amenable  $G$  subshift setting
- ▶ Equilibrium States are Always Gibbs-like.

# Subshifts Represent Lattice Models

- ▶ Lattice gas model,
- ▶ Ising model (and generalizations like the Potts Model).
- ▶ a variety of discretizations of state space,
- ▶ etc.

# $\mathbb{Z}^d$ -Subshifts

- ▶ Finite set  $\mathcal{A}$  called the **alphabet** - Think  $\{0, 1\}$  or  $\{-1, +1\}$ .
- ▶ The **left shift map**  $\sigma : \mathcal{A}^{\mathbb{Z}^d} \rightarrow \mathcal{A}^{\mathbb{Z}^d}$  is defined by

$$\sigma_k(x)_n = x_{k+n}.$$

- ▶ We call a closed,  $\sigma$ -invariant  $X \subset \mathcal{A}^{\mathbb{Z}^d}$  a **subshift**.
- ▶ Let  $L_{[-n,n]^d}(X)$  denote the  $[-n, n]^d$ -**language** of  $X$ . I.e. the **configurations** or **words** of shape  $[-n, n]^d$  that appear in  $X$ .
- ▶ For a given word  $w \in L_{[-n,n]^d}(X)$ , we define the **cylinder set**

$$[w] = \{x \in X : x_{[-n,n]^d} = w\}.$$

# Golden Mean Subshift

Example:

- ▶  $X = \{x \in \{0, 1\}^{\mathbb{Z}} : \forall i \in \mathbb{Z}, x_{i,i+1} \neq 11\}.$
- ▶  $\dots 100.010\dots = x \in X.$
- ▶  $\sigma(x) = \dots 1000.10\dots$
- ▶  $L_3(X) = \{000, 001, 010, 100, 101\}.$

(This is the  $1 - D$  lattice gas model with radius 1. )

# Invariant Measures

For a subshift  $X$ , let  $M_\sigma(X)$  be the collection of all  $\sigma$ -invariant Borel probability measures on  $X$ .

►  $\mu(A) = \mu(\sigma^{-1}A)$  for all measurable  $A \subset X$ .

In the subshift setting, cylinder sets generate the Borel  $\sigma$ -algebra so we can restrict ourselves computing  $\mu([w])$  for  $w \in L(X)$ .

# Measure Theoretic Entropy

- For each  $\mu \in M_\sigma(X)$  we can compute the Kolmogorov-Sinai entropy of  $\mu$  by

$$h(\mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{w \in L_n(X)} -\mu([w]) \log \mu([w]).$$

# Entropy for Subshifts

Example on  $X = \{0, 1\}^{\mathbb{Z}}$ :

- ▶ Let  $\nu$  be Bernoulli assigning  $(1/3, 2/3)$  to  $[0]$  and  $[1]$  respectively. Then

$$h(\nu) = -\frac{1}{3} \log \frac{1}{3} - \frac{2}{3} \log \frac{2}{3} < \log 2.$$

- ▶ Let  $\mu$  be Bernoulli assigning  $(1/2, 1/2)$  to  $[0]$  and  $[1]$  respectively. Then

$$h(\mu) = -\log \frac{1}{2} = \log 2.$$



# Potentials

- ▶ Given a subshift  $X$ , a continuous, real valued  $\phi \in C(X)$  is called a **potential**.
- ▶ Potentials assign *energy* to points in  $x$ .
  - ▶ (Really  $\phi \approx -\frac{E}{kT}$ )
- ▶ By additionally considering the energy of points, we can construct a generalization of topological entropy: **topological pressure**.

# Pressure and Equilibrium States

We define **topological pressure** of a potential  $\phi$  to be:

$$P_{top}(\phi) = \sup\{h(\mu) + \int \phi d\mu : \mu \in M_\sigma(X)\}.$$

► With  $\phi \approx -\frac{E}{kT}$ ,  $\max h + \int \phi$  is equivalent to  $\min E - TS$ .

We say  $\mu$  is an **equilibrium state** for  $\phi$  if

$$h(\mu) + \int \phi d\mu = P_{top}(\phi).$$

# Equilibrium State Example

Example:

- ▶ Let  $X = \{0, 1\}^{\mathbb{Z}}$ ,
- ▶ Define  $\phi([0]) = 0$ ,  $\phi([1]) = 1$ .
- ▶ Find  $\mu$  that maximizes  $h(\mu) + \int \phi d\mu$ .
- ▶ Balance the entropy and average energy.

# Equilibrium State Example

- ▶ Let's look at  $\mu_p$  Bernoulli  $(p, 1 - p)$
- ▶  $h(\mu_p) + \int \phi d\mu_p$ .

$$h(\mu_p) = -p \log p - (1 - p) \log(1 - p),$$

$$\int \phi d\mu_p = 0 \cdot p + 1 \cdot (1 - p) = 1 - p.$$

- ▶ ... some calculus later...
- ▶ Maximized at  $p = \frac{1}{1+e}$ .

# Equilibrium State Example

- ▶ For  $X = \{0, 1\}^{\mathbb{Z}}$ ,
- ▶  $\phi([0]) = 0$ ,  $\phi([1]) = 1$ ,
- ▶  $\mu$  Bernoulli  $(\frac{1}{1+e}, \frac{e}{1+e})$ ,

$\mu$  satisfies the following equation for any  $w \in L_n(X)$ ,  $x \in [w]$ :

$$\mu([w]) = \frac{\exp(\phi_n(x))}{\sum_{v \in L_n(X)} \exp(\phi_n(y))} \quad (1)$$

where  $\phi_n(x) = \sum_{i=0}^{n-1} \phi(\sigma^i(x))$ .

# Uniqueness of Equilibrium States in $\mathbb{Z}$

**Fact:** Given a mixing SFT  $X \subset \{0, 1\}^{\mathbb{Z}}$  and a Hölder continuous potential  $\phi \in C(X)$ , there exists a unique equilibrium state  $\mu_\phi \in M_\sigma(X)$  and  $\mu_\phi$  is Gibbs for  $\phi$ .

$\mu_\phi$  being Gibbs for  $\phi$  means (1) holds after conditioning on background configurations.

# Extended DLR Theorems

For nice subshifts  $X \subset \mathcal{A}^G$ , and sufficiently regular potentials  $\phi \in C(X)$ , the following are equivalent:

- ▶  $\mu$  is an equilibrium state for  $\phi$
- ▶  $\mu$  is a  $\sigma$ -invariant Gibbs measure for  $\phi$ .

Outside of  $\mathbb{Z}$ , uniqueness is out the window (see Ising model in  $\mathbb{Z}^2$ ).

What about when  $X$  isn't a nice subshift?

## $\mathbb{Z}^d$ Measures of Maximal Entropy

For words  $v, w \in L_F(X)$ , we denote  $v \rightarrow w$  to mean  $v$  can be replaced by  $w$ .

- **Garcia-Ramos and Pavlov, 2019:** Let  $X \subset \mathcal{A}^{\mathbb{Z}^d}$  be any subshift,  $w, v \in L_F(X)$ , and  $\mu$  be any m.m.e.. If  $v \rightarrow w$ , then

$$\mu([v]) \leq \mu([w]).$$



# G Equilibrium States

Let  $G$  is a countable, amenable group,  $X \subset \mathcal{A}^G$  be a subshift, and  $\phi \in C(X)$  have summable variation, and  $\mu_\phi$  an equilibrium state for  $\phi$ .

► **H., 2024:** If  $v, w \in L_F(X)$  such that  $v \rightarrow w$ , then

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \sup_{x \in [v]} \exp \left( \sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x))) \right).$$

# A Strange Equivalent Gibbs Definition

Let  $X \subset \mathcal{A}^G$  be a subshift,  $\phi \in C(X)$  be a potential, and  $\mu \in M_\sigma(X)$ .

►  $\mu$  is Gibbs for  $\phi$  if for all  $v, w \in L_F(X)$ ,

$$\frac{d(\mu \circ \xi_{v,w})}{d\mu}(x) = \exp \left( \sum_{n \in \mathbb{Z}^d} \phi(\sigma_n(\xi_{v,w}(x))) - \phi(\sigma_n(x)) \right).$$

## $\mathbb{Z}^d$ Bounds when $v \leftrightarrow w$

Let  $X \subset \mathcal{A}^{\mathbb{Z}^d}$  be a subshift,  $\phi \in C(X)$  be a potential with  $d$ -summable variation, and  $\mu \in M_\sigma(X)$  an equilibrium state for  $\phi$ .

► **Meyerovitch, 2013:** For any  $v, w \in L_F(X)$  such that  $v \leftrightarrow w$ ,

$$\frac{d(\mu \circ \xi_{v,w})}{d\mu}(x) = \exp \left( \sum_{g \in G} \phi(\sigma_g(\xi_{v,w}(x))) - \phi(\sigma_g(x)) \right).$$

## $G$ Bounds when $\nu \rightarrow w$

Let  $G$  is a countable, amenable group,  $X \subset \mathcal{A}^G$  be a subshift, and  $\phi \in C(X)$  have summable variation, and  $\mu_\phi$  an equilibrium state for  $\phi$ .

- **H., 2024:** If  $\nu, w \in L_F(X)$  such that  $\nu \rightarrow w$ , then for  $\mu_\phi$ -a.e.  $x \in [w]$ :

$$\frac{d(\mu \circ \xi_{\nu, w})}{d\mu}(x) \leq \exp \left( \sum_{g \in G} \phi(\sigma_g(\xi_{\nu, w}(x))) - \phi(\sigma_g(x)) \right).$$

## $G$ Bounds when $v \leftrightarrow w$

As a corollary we extend the results of Meyerovitch to the amenable group subshift setting:

- ▶ **H., 2024:** Let  $X \subset \mathcal{A}^G$  be any subshift, let  $\phi \in C(X)$  have summable variation, and let  $\mu_\phi$  be an equilibrium state for  $\phi$ . Then  $\mu_\phi$  is topologically Gibbs.

# Thank you!

Questions?