

under Graduate Homework In Mathematics

FunctionalAnalysis 12

王胤雅

201911010205

201911010205@mail.bnu.edu.cn

2023 年 11 月 25 日



General fire extinguisher

PROBLEM I \mathcal{X} is a linear space on \mathbb{C} . p is a seminorm on \mathcal{X} . $p(x_0) \neq 0, x_0 \in \mathcal{X}$. Prove: $\exists f$ is a linear functional on \mathcal{X} such that

1. $f(x_0) = 1$
2. $|f(x)| \leq \frac{p(x)}{p(x_0)}, \forall x \in \mathcal{X}$.

SOLUTION. Consider $p^* : \mathcal{X} \rightarrow \mathbb{C}, x \mapsto \frac{p(x)}{p(x_0)}$. Obviously p^* is a seminorm on \mathcal{X} . Let $\mathcal{X}_0 := \text{Span}\{x_0\} \subset \mathcal{X}$ is a subspace of \mathcal{X} . $f : \mathcal{X}_0 \rightarrow \mathbb{C}, \alpha x_0 \mapsto \alpha f(x_0)$, where $f(x_0) = 1$. So f is a linear functional on \mathcal{X}_0 . And $\forall x \in \mathcal{X}_0, x = \alpha x_0, |f(x)| = |\alpha| |f(x_0)| \leq |\alpha| = \frac{p(\alpha x_0)}{p(x_0)} = p^*(x)$. Thus, by Hahn-Banach theorem, $\exists \tilde{f} : \mathcal{X} \rightarrow \mathbb{R}$ is a linear functional on \mathcal{X} such that

1. $\tilde{f}(x) = f(x), \forall x \in \mathcal{X}_0$.
2. $|\tilde{f}(x)| \leq p^*(x) = \frac{p(x)}{p(x_0)}, \forall x \in \mathcal{X}$.

So $\tilde{f}(x_0) = f(x_0)$. □

PROBLEM II \mathcal{X} is a B^* space, $\{x_n\}_{n=1}^\infty \subset \mathcal{X}$ such that $\forall f \in \mathcal{X}^*, \{f(x_n)\}_{n=1}^\infty$ is bounded. Prove that $\{x_n\}_{n=1}^\infty$ is bounded.

SOLUTION. Since there is an embedding map from $\mathcal{X} \rightarrow \mathcal{X}^{**}$, which keeps norm. Regard $\{x_n\}_{n=1}^\infty$ as subset of \mathcal{X}^{**} . And $\mathcal{X}^{**} = \mathcal{L}(\mathcal{X}^*, \mathbb{K}), \mathcal{X}^* = \mathcal{L}(X, \mathbb{K})$. \mathbb{K} is complete, so \mathcal{X}^* is a B space. Besides, $\forall f \in \mathcal{X}^*, \sup_{n \in \mathbb{N}_+} |x_n(f)| = \sup_{n \in \mathbb{N}_+} |f(x_n)| < \infty$. By Banach-Steinhaus theorem, $\sup_{n \in \mathbb{N}_+} \|x_n\| < \infty$. □

PROBLEM III \mathcal{X} is a B^* space, \mathcal{X}_0 is a closed subspace of \mathcal{X} . Prove that $\forall x \in \mathcal{X}, \inf_{y \in \mathcal{X}_0} \|x - y\| = \sup\{|f(x)| : f \in \mathcal{X} \|f\| = 1, f|_{\mathcal{X}_0} = 1\}$.

Lemma 1. \mathcal{X} is a B^* space, let $H_f^\lambda := \{x \in \mathcal{X} : f(x) = \lambda\}$ where f is a linear functional on \mathcal{X} . If $\|f\| = 1$, then $|f(x)| = d(x, H_f^0), \forall x \in \mathcal{X}$, where $d(x, H_f^0) := \inf_{z \in H_f^0} \|x - z\|$.

证明. 1. $|f(x)| \leq d(x, H_f^0)$: Since $\forall \varepsilon > 0, \exists y \in H_f^0$ such that $\|x - y\| \geq d(x, H_f^0) + \varepsilon$. And $|f(x)| = |f(x - y)|$ □

SOLUTION. □