


# **The hamburger theorem**

Mikio Kano and Jan Kynčl


EPFL

### Problem:

Given  $n$  red and  $n$  blue points in the plane in general position, draw  $n$  noncrossing red-blue segments.




**Solution:** take the shortest red-blue perfect matching.




**Theorem:** (Akiyama and Alon, 1989)

Given point sets  $X_1, X_2, \dots, X_d \subset \mathbb{R}^d$  in general position, with  $|X_1| = |X_2| = \dots = |X_d| = n$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.



**Theorem:** (Akiyama and Alon, 1989)


Given point sets  $X_1, X_2, \dots, X_d \subset \mathbb{R}^d$  in general position, with  $|X_1| = |X_2| = \dots = |X_d| = n$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.



**Proof:** recursive cutting by hyperplanes, using the discrete ham-sandwich theorem.

**Theorem:** (Akiyama and Alon, 1989)

Given point sets  $X_1, X_2, \dots, X_d \subset \mathbb{R}^d$  in general position, with  $|X_1| = |X_2| = \dots = |X_d| = n$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.




**Proof:** recursive cutting by hyperplanes, using the discrete ham-sandwich theorem.

**The discrete ham-sandwich theorem:** (Stone and Tukey, 1942)

If  $X_1, X_2, \dots, X_d \subset \mathbb{R}^d$  are disjoint finite sets in general position, then there is a hyperplane that bisects each  $X_i$  exactly in half.


**Theorem:** (Kano, Suzuki and Uno, 2014)

Let  $R, G, B \subset \mathbb{R}^2$  be sets of red, green and blue points in general position such that  $|R| + |G| + |B| = 2n$  and  $|R|, |G|, |B| \leq n$ . Then there are  $n$  disjoint rainbow segments.



**Proof:** using a special result for partitioning colored sets on a line

**Alternative solution:** shortest rainbow perfect matching



**Theorem:** (Kano, Suzuki and Uno, 2014)

Let  $R, G, B \subset \mathbb{R}^2$  be sets of red, green and blue points in general position such that  $|R| + |G| + |B| = 2n$  and  $|R|, |G|, |B| \leq n$ . Then there are  $n$  disjoint rainbow segments.

**Corollary:** Given point sets  $X_1, X_2, \dots, X_r \subset \mathbb{R}^2$  in general position such that  $|X_1| + |X_2| + \dots + |X_r| = 2n$  and  $|X_i| \leq n$  for every  $i \in [d+1]$ , then there are  $n$  disjoint rainbow segments.

**Proof:** merging the smallest sets:

$$(4, 4, 3, 2, 1) \rightarrow (4, 4, 3, 3) \rightarrow (4, 4, 6)$$

(also shortest rainbow perfect matching)

**Conjecture:** (Kano and Suzuki) Let  $r \geq d \geq 3$  and  $n \geq 1$ . Given point sets  $X_1, X_2, \dots, X_r \subset \mathbb{R}^d$  in general position such that  $|X_1| + |X_2| + \dots + |X_r| = dn$  and  $|X_i| \leq n$  for every  $i \in [r]$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.



**Conjecture:** (Kano and Suzuki) Let  $r \geq d \geq 3$  and  $n \geq 1$ . Given point sets  $X_1, X_2, \dots, X_r \subset \mathbb{R}^d$  in general position such that  $|X_1| + |X_2| + \dots + |X_r| = dn$  and  $|X_i| \leq n$  for every  $i \in [r]$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.

**Theorem:** The conjecture is true for  $d \geq 2$  and  $r = d + 1$ .

**Conjecture:** (Kano and Suzuki) Let  $r \geq d \geq 3$  and  $n \geq 1$ . Given point sets  $X_1, X_2, \dots, X_r \subset \mathbb{R}^d$  in general position such that  $|X_1| + |X_2| + \dots + |X_r| = dn$  and  $|X_i| \leq n$  for every  $i \in [r]$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.

**Theorem:** The conjecture is true for  $d \geq 2$  and  $r = d + 1$ .

**Plan of proof:** recursive cutting by a hyperplane, into a pair of **balanced** subsets; that is, for each of the two halfspaces  $H$ , we want

$$|H \cap X_i| \leq \frac{1}{d} \sum_{j=1}^r |H \cap X_j|.$$

Moreover, the total number of points in  $H$  must be positive and divisible by  $d$ .

**Conjecture: (Kano and Suzuki)** Let  $r \geq d \geq 3$  and  $n \geq 1$ . Given point sets  $X_1, X_2, \dots, X_r \subset \mathbb{R}^d$  in general position such that  $|X_1| + |X_2| + \dots + |X_r| = dn$  and  $|X_i| \leq n$  for every  $i \in [r]$ , then there are  $n$  disjoint rainbow  $(d - 1)$ -dimensional simplices.

**Theorem:** The conjecture is true for  $d \geq 2$  and  $r = d + 1$ .

**Plan of proof:** recursive cutting by a hyperplane, into a pair of **balanced** subsets; that is, for each of the two halfspaces  $H$ , we want

$$|H \cap X_i| \leq \frac{1}{d} \sum_{j=1}^r |H \cap X_j|.$$

Moreover, the total number of points in  $H$  must be positive and divisible by  $d$ .

→ we need a generalization of the discrete ham-sandwich theorem to  $d + 1$  sets in  $\mathbb{R}^d$ .

First we show a continuous version, then discretize it.

**Definition:**

Let  $r \geq d$  and let  $\mu_1, \mu_2, \dots, \mu_r$  be finite Borel measures on  $\mathbb{R}^d$ . We say that  $\mu_1, \mu_2, \dots, \mu_r$  are **balanced** in a subset  $X \subseteq \mathbb{R}^d$  if for every  $i \in [r]$ , we have

$$\mu_i(X) \leq \frac{1}{d} \cdot \sum_{j=1}^r \mu_j(X).$$

## The hamburger theorem:

Let  $d \geq 2$ , let  $\mu_1, \mu_2, \dots, \mu_{d+1}$  be absolutely continuous finite Borel measures on  $\mathbb{R}^d$ .

## The hamburger theorem:

Let  $d \geq 2$ , let  $\mu_1, \mu_2, \dots, \mu_{d+1}$  be absolutely continuous finite Borel measures on  $\mathbb{R}^d$ . Let

- $\omega_i := \mu_i(\mathbb{R}^d)$  for  $i \in [d+1]$ ,
- $\omega := \min\{\omega_i; i \in [d+1]\}$ ,

## The hamburger theorem:

Let  $d \geq 2$ , let  $\mu_1, \mu_2, \dots, \mu_{d+1}$  be absolutely continuous finite Borel measures on  $\mathbb{R}^d$ . Let

- $\omega_i := \mu_i(\mathbb{R}^d)$  for  $i \in [d+1]$ ,
- $\omega := \min\{\omega_i; i \in [d+1]\}$ ,

and assume that

- $\sum_{j=1}^{d+1} \omega_j = 1$ ,
- $\mu_1, \mu_2, \dots, \mu_{d+1}$  are balanced in  $\mathbb{R}^d$ .

## The hamburger theorem:

Let  $d \geq 2$ , let  $\mu_1, \mu_2, \dots, \mu_{d+1}$  be absolutely continuous finite Borel measures on  $\mathbb{R}^d$ . Let

- $\omega_i := \mu_i(\mathbb{R}^d)$  for  $i \in [d+1]$ ,
- $\omega := \min\{\omega_i; i \in [d+1]\}$ ,

and assume that

- $\sum_{j=1}^{d+1} \omega_j = 1$ ,
- $\mu_1, \mu_2, \dots, \mu_{d+1}$  are balanced in  $\mathbb{R}^d$ .

Then there exists a hyperplane  $h$  such that for each open halfspace  $H$  defined by  $h$ , the measures  $\mu_1, \mu_2, \dots, \mu_{d+1}$  are balanced in  $H$  and

$$\sum_{j=1}^{d+1} \mu_j(H) \geq \min\left(\frac{1}{2}, 1 - d\omega\right) \geq \frac{1}{d+1}.$$



## The hamburger theorem:

Let  $d \geq 2$ , let  $\mu_1, \mu_2, \dots, \mu_{d+1}$  be absolutely continuous finite Borel measures on  $\mathbb{R}^d$ . Let

- $\omega_i := \mu_i(\mathbb{R}^d)$  for  $i \in [d+1]$ ,
- $\omega := \min\{\omega_i; i \in [d+1]\}$ ,

and assume that

- $\sum_{j=1}^{d+1} \omega_j = 1$ ,
- $\mu_1, \mu_2, \dots, \mu_{d+1}$  are balanced in  $\mathbb{R}^d$ .


Then there exists a hyperplane  $h$  such that for each open halfspace  $H$  defined by  $h$ , the measures  $\mu_1, \mu_2, \dots, \mu_{d+1}$  are balanced in  $H$  and

$$\sum_{j=1}^{d+1} \mu_j(H) \geq \min\left(\frac{1}{2}, 1 - d\omega\right) \geq \frac{1}{d+1}.$$


---

For  $\omega = 0$  we get exactly the ham-sandwich theorem.

- The lower bound  $1/(d+1)$  is tight:



- discretization is nontrivial:



## Proof of the hamburger theorem

analogous to the proof of the ham-sandwich theorem:

- parametrize half-spaces in  $\mathbb{R}^d$  by the points of  $S^d = \{\mathbf{u} = (u_0, u_1, \dots, u_d) \in \mathbb{R}^{d+1}; u_0^2 + u_1^2 + \dots + u_d^2 = 1\}$ :  
if  $|u_0| < 1$ , then

$$H^-(\mathbf{u}) := \{(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_d) \in \mathbb{R}^d; u_1 \mathbf{x}_1 + u_2 \mathbf{x}_2 + \dots + u_d \mathbf{x}_d < u_0\},$$

$$H^+(\mathbf{u}) := \{(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_d) \in \mathbb{R}^d; u_1 \mathbf{x}_1 + u_2 \mathbf{x}_2 + \dots + u_d \mathbf{x}_d > u_0\},$$

## Proof of the hamburger theorem

analogous to the proof of the ham-sandwich theorem:

- parametrize half-spaces in  $\mathbb{R}^d$  by the points of  $S^d = \{\mathbf{u} = (u_0, u_1, \dots, u_d) \in \mathbb{R}^{d+1}; u_0^2 + u_1^2 + \dots + u_d^2 = 1\}$ :  
if  $|u_0| < 1$ , then

$$H^-(\mathbf{u}) := \{(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_d) \in \mathbb{R}^d; u_1 x_1 + u_2 x_2 + \dots + u_d x_d < u_0\},$$

$$H^+(\mathbf{u}) := \{(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_d) \in \mathbb{R}^d; u_1 x_1 + u_2 x_2 + \dots + u_d x_d > u_0\},$$

and

$$H^-(1, 0, 0, \dots, 0) := \mathbb{R}^d, \quad H^+(1, 0, 0, \dots, 0) := \emptyset,$$

$$H^-(-1, 0, 0, \dots, 0) := \emptyset, \quad H^+(-1, 0, 0, \dots, 0) := \mathbb{R}^d.$$

## Proof of the hamburger theorem

analogous to the proof of the ham-sandwich theorem:

- parametrize half-spaces in  $\mathbb{R}^d$  by the points of  $S^d = \{\mathbf{u} = (u_0, u_1, \dots, u_d) \in \mathbb{R}^{d+1}; u_0^2 + u_1^2 + \dots + u_d^2 = 1\}$ :  
if  $|u_0| < 1$ , then

$$H^-(\mathbf{u}) := \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d; u_1 x_1 + u_2 x_2 + \dots + u_d x_d < u_0\},$$

$$H^+(\mathbf{u}) := \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d; u_1 x_1 + u_2 x_2 + \dots + u_d x_d > u_0\},$$

and

$$H^-(1, 0, 0, \dots, 0) := \mathbb{R}^d, \quad H^+(1, 0, 0, \dots, 0) := \emptyset,$$

$$H^-(-1, 0, 0, \dots, 0) := \emptyset, \quad H^+(-1, 0, 0, \dots, 0) := \mathbb{R}^d.$$

We have  $H^-(\mathbf{u}) = H^+(-\mathbf{u})$  for every  $\mathbf{u} \in S^d$ .

- define  $f = (f_1, \dots, f_{d+1}) : S^d \rightarrow \mathbb{R}^{d+1}$  by

$$f_i(\mathbf{u}) := \mu_i(H^-(\mathbf{u})).$$

- define  $f = (f_1, \dots, f_{d+1}) : S^d \rightarrow \mathbb{R}^{d+1}$  by

$$f_i(\mathbf{u}) := \mu_i(H^-(\mathbf{u})).$$

- $f$  is continuous since  $\mu_i(h) = 0$  for every hyperplane  $h$

- define  $f = (f_1, \dots, f_{d+1}) : S^d \rightarrow \mathbb{R}^{d+1}$  by

$$f_i(\mathbf{u}) := \mu_i(H^-(\mathbf{u})).$$

- $f$  is continuous since  $\mu_i(h) = 0$  for every hyperplane  $h$
- $f[S^d] \subseteq B := \prod_{i=1}^{d+1} [0, \omega_i]$



- define  $f = (f_1, \dots, f_{d+1}) : S^d \rightarrow \mathbb{R}^{d+1}$  by



$$f_i(\mathbf{u}) := \mu_i(H^-(\mathbf{u})).$$



- $f$  is continuous since  $\mu_i(h) = 0$  for every hyperplane  $h$
- $f[S^d] \subseteq B := \prod_{i=1}^{d+1} [0, \omega_i]$
- $f(\mathbf{u})$  and  $f(-\mathbf{u})$  symmetric about the center  $\mathbf{b}$  of  $B$

- define  $f = (f_1, \dots, f_{d+1}) : S^d \rightarrow \mathbb{R}^{d+1}$  by



$$f_i(\mathbf{u}) := \mu_i(H^-(\mathbf{u})).$$

- $f$  is continuous since  $\mu_i(h) = 0$  for every hyperplane  $h$
- $f[S^d] \subseteq B := \prod_{i=1}^{d+1} [0, \omega_i]$
- $f(\mathbf{u})$  and  $f(-\mathbf{u})$  symmetric about the center  $\mathbf{b}$  of  $B$
- our goal is to show that the image of  $f$  intersects the **target polytope**, determined by the conditions “balanced” and “nontrivial”






- assume  $\omega = \omega_{d+1}$
- $t := \min \left( \frac{1}{2d}, \frac{1}{d} - \omega \right)$



- assume  $\omega = \omega_{d+1}$
- $t := \min\left(\frac{1}{2d}, \frac{1}{d} - \omega\right)$
- $\mathbf{a} := (t, t, \dots, t, 0)$ ,  $\mathbf{b} := (\omega_1/2, \omega_2/2, \dots, \omega_{d+1}/2)$ ,
- $\mathbf{c} := (\omega_1 - t, \omega_2 - t, \dots, \omega_d - t, \omega_{d+1})$



- assume  $\omega = \omega_{d+1}$
- $t := \min\left(\frac{1}{2d}, \frac{1}{d} - \omega\right)$
- $\mathbf{a} := (t, t, \dots, t, 0)$ ,  $\mathbf{b} := (\omega_1/2, \omega_2/2, \dots, \omega_{d+1}/2)$ ,
- $\mathbf{c} := (\omega_1 - t, \omega_2 - t, \dots, \omega_d - t, \omega_{d+1})$
- $\ell := \text{line } \mathbf{ac}$
- $\pi_\ell := \text{projection to a hyperplane orthogonal to } \ell$
- $g(\mathbf{u}) := \pi_\ell(f(\mathbf{u}) - \mathbf{b})$

$g$  is antipodal map from  $S^d$  to  $\mathbb{R}^d$ . By the Borsuk–Ulam theorem, there exists  $\mathbf{u} \in S^d$  such that  $g(\mathbf{u}) = \mathbf{0}$ , which means that  $f(\mathbf{u}) \in \ell$ .