

Studies of Quantum Dots using Machine Learning

Even Marius Nordhagen

University of Oslo

evenmn@fys.uio.no

October 1, 2019

Quantum mechanics

Machine learning

Our approach

Results

$$i\hbar \frac{\partial}{\partial t} |\Psi(\mathbf{r}, t)\rangle = \hat{\mathcal{H}} |\Psi(\mathbf{r}, t)\rangle$$

The time-independent Schrödinger equation

$$E_n = \frac{\int d\mathbf{r} \Psi_n^*(\mathbf{r}) \hat{\mathcal{H}} \Psi_n(\mathbf{r})}{\int d\mathbf{r} \Psi_n^*(\mathbf{r}) \Psi_n(\mathbf{r})}$$

$$\hat{\mathcal{H}} = \sum_{i=1}^N \left(-\frac{1}{2} \nabla_i^2 + \frac{1}{2} \omega^2 r_i^2 \right)$$

Many-body problem

$$\hat{\mathcal{H}} = \sum_{i=1}^N \left(-\frac{1}{2} \nabla_i^2 + \frac{1}{2} \omega^2 r_i^2 \right) + \sum_{i < j} \frac{1}{r_{ij}}$$

$$\begin{aligned} E_0 < E &= \frac{\int d\mathbf{r} \Psi_0(\mathbf{r})^* \hat{\mathcal{H}} \Psi_0(\mathbf{r})}{\int d\mathbf{r} \Psi_0(\mathbf{r})^* \Psi_0(\mathbf{r})} \\ &= \int d\mathbf{r} E_L(\mathbf{r}) P(\mathbf{r}) \\ &\approx \frac{1}{M} \sum_{i=1}^M E_L(\mathbf{r}_i) \end{aligned}$$

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something

something, something something