

Studies of Quantum Dots using Machine Learning



Even Marius Nordhagen

University of Oslo

evenmn@fys.uio.no

December 4, 2019

Outline

- ▶ Motivation
- ▶ Quantum Theory
- ▶ Machine Learning
- ▶ Methods
- ▶ Results
- ▶ Conclusion
- ▶ (Code)



Motivation



Why quantum dots?

- ▶ Technology¹
- ▶ Theoretically
- ▶ Experimentally (2D)



Why is it challenging?

- ▶ Quantum many-body problem
- ▶ Fermi-Dirac statistics



How to overcome the challenges?

- ▶ Efforts..
- ▶ gh
- ▶ Our approach: Machine Learning^{2,3}



Quantum Theory



The Time-independent Schrödinger Equation

A stationary quantum mechanical system is described by

$$E_n = \frac{\int d\mathbf{R} \Psi_n^*(\mathbf{R}) \hat{\mathcal{H}} \Psi_n(\mathbf{R})}{\int d\mathbf{R} \Psi_n^*(\mathbf{R}) \Psi_n(\mathbf{R})}$$

which gives the energy of state n .




Machine Learning Theory




Machine Learning

Machine learning is the science of getting computers to act without being explicitly programmed.

Feed-forward Neural Network



Restricted Boltzmann machines



Methods



13



UiO : **University of Oslo**

Variational Monte Carlo (VMC)

Exploit the variational principle in order to obtain the ground state energy

$$\begin{aligned} E_0 < E_{\text{VMC}} &= \frac{\int d\mathbf{R} \Psi_T(\mathbf{R})^* \hat{\mathcal{H}} \Psi_T(\mathbf{R})}{\int d\mathbf{R} \Psi_T(\mathbf{R})^* \Psi_T(\mathbf{R})} \\ &= \int d\mathbf{R} \underbrace{\frac{\Psi_T(\mathbf{R})^* \Psi_T(\mathbf{R})}{\int d\mathbf{R} \Psi_T(\mathbf{R})^* \Psi_T(\mathbf{R})}}_{P(\mathbf{R})} \cdot \underbrace{\frac{1}{\Psi_T(\mathbf{R})} \hat{\mathcal{H}} \Psi_T(\mathbf{R})}_{E_L(\mathbf{R})} \end{aligned}$$

Monte Carlo Integration

We attempt to solve the integral by sampling from the probability density function $P(\mathbf{r})$

$$\begin{aligned} E_{\text{VMC}} &= \int d\mathbf{R} E_L(\mathbf{R}) P(\mathbf{R}) \\ &\approx \frac{1}{M} \sum_{i=1}^M E_L(\mathbf{R}_i) \end{aligned}$$

Trial Wave Function

$$P(\mathbf{R}) \propto \Psi_T(\mathbf{R})^* \Psi_T(\mathbf{R})$$

Use the Slater-Jastrow function as our trial wave function

$$\Psi_T(\mathbf{R}) = |\hat{D}(\mathbf{R})|J(\mathbf{R})$$

where the Slater matrix, $\hat{D}(\mathbf{R})$, contains all the single-particle functions

$$\hat{D}(\mathbf{R}) = \begin{pmatrix} \phi_1(\mathbf{r}_1) & \phi_2(\mathbf{r}_1) & \dots & \phi_N(\mathbf{r}_1) \\ \phi_1(\mathbf{r}_2) & \phi_2(\mathbf{r}_2) & \dots & \phi_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ \phi_1(\mathbf{r}_N) & \phi_2(\mathbf{r}_N) & \dots & \phi_N(\mathbf{r}_N) \end{pmatrix}$$

Single-particle Functions

The Hermite functions,

$$\phi_n(\mathbf{r}) \propto H_n(\mathbf{r}) \exp\left(-\frac{1}{2}\alpha\omega|\mathbf{r}|^2\right),$$

are used as the single-particle functions for quantum dots in standard VMC. The Gaussian can be factorized out from the Slater determinant.

$$|\hat{D}(\mathbf{R}; \alpha)| \propto \exp\left(-\frac{1}{2}\alpha\omega|\mathbf{R}|^2\right) \begin{vmatrix} H_1(\mathbf{r}_1) & H_2(\mathbf{r}_1) & \dots & H_N(\mathbf{r}_1) \\ H_1(\mathbf{r}_2) & H_2(\mathbf{r}_2) & \dots & H_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ H_1(\mathbf{r}_N) & H_2(\mathbf{r}_N) & \dots & H_N(\mathbf{r}_N) \end{vmatrix}$$

Restricted Boltzmann Machine

We use the marginal distribution of the visible units as the single-particle functions in the Slater determinant, and see if they can model the correlations

$$\phi_n(\mathbf{r}) \propto H_n(\mathbf{r})P(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W})$$

where $P(\mathbf{r})$ is the marginal distribution of the visible units.

$$|\hat{D}(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W})| \propto P(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W}) \begin{vmatrix} H_1(\mathbf{r}_1) & H_2(\mathbf{r}_1) & \dots & H_N(\mathbf{r}_1) \\ H_1(\mathbf{r}_2) & H_2(\mathbf{r}_2) & \dots & H_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ H_1(\mathbf{r}_N) & H_2(\mathbf{r}_N) & \dots & H_N(\mathbf{r}_N) \end{vmatrix}$$

Jastrow Factor

The Jastrow factor is added to account for the correlations
Simple Jastrow factor

$$J(\mathbf{r}; \boldsymbol{\beta}) = \exp \left(\sum_{i=1}^N \sum_{j>i}^N \beta_{ij} r_{ij} \right).$$

Padé-Jastrow factor

$$J(\mathbf{r}; \boldsymbol{\beta}) = \exp \left(\sum_{i=1}^N \sum_{j>i}^N \frac{a_{ij} r_{ij}}{1 + \beta r_{ij}} \right).$$

Results



Ground State Energy

Table: Ground state energy of two-dimensional quantum dot with $N = 2$ electrons and frequency $\omega = 1.0$.

ω	RBM	RBM+SJ	RBM+PJ	HF [†]	Exact ^{††}
1/6	0.7036(1)	0.67684(7)	0.66715(6)	0.768675	2/3
1	3.0803(2)	3.02108(5)	2.999587(5)	3.1690	3



One-body density





Figure: Plots of the one-body density profiles, $\rho(x, y)$, for two-dimensional quantum dots with $N = 20$ electrons and frequency $\omega = 1.0$.

Computational Cost



Conclusion



Conclusions

- ▶ RBM is able to account for most of the correlations
- ▶ RBM+PJ implies to give a lower ground state energy and model the correlations better than a traditional VMC
- ▶ RBM+SJ is both more expensive and less accurate than its fellow methods, and we see no reason to choose it

Future Work






- ▶ Repeat the exercise using spherical coordinates - interactions are easier to model in spherical coordinates
- ▶ Check the ability of modeling the three-body correlations, considering nuclear systems
- ▶ Reduce the computational cost

Thank you!



UiO : **University of Oslo**

References

-  Samsung QLED TV | The Frame | Premium UHD TV.
<http://www.samsung.com/global/tv/blog/why-are-quantum-dot-displays-so-good/>.
-  Carleo, G. & Troyer, M. Solving the Quantum Many-Body Problem with Artificial Neural Networks. *Science* **355**, 602 (2017).
-  Pfau, D., Spencer, J. S., Matthews, A. G. G. & Foulkes, W. M. C. Ab-Initio Solution of the Many-Electron Schrödinger Equation with Deep Neural Networks. [arXiv: 1909.02487](https://arxiv.org/abs/1909.02487) (2019).
-  Mariadason, A. A. *Quantum Many-Body Simulations of Double Dot System*. MA thesis (2018).
-  Taut, M. Two electrons in an external oscillator potential: Particular analytic solutions of a Coulomb correlation problem. *Physica Scripta* **48**, 3561 (1993).