

Studies of Quantum Dots using Machine Learning

Even Marius Nordhagen

University of Oslo

evenmn@fys.uio.no

December 3, 2019



- ▶ Motivation
- ▶ Quantum Theory
- ▶ Machine Learning
- ▶ Methods
- ▶ Results
- ▶ Conclusion
- ▶ (Code)

Motivation

Why quantum dots?



- ▶ Technology
- ▶ Theoretically
- ▶ Experimentally

Why is it challenging?



- ▶ Quantum many-body problem
- ▶ Fermi-Dirac statistics

How to overcome the challenges?



- ▶ Efforts..
- ▶ gh
- ▶ Our approach: Machine Learning

Theory

The Time-independent Schrödinger Equation



A stationary quantum mechanical system is described by


$$E_n = \frac{\int d\mathbf{r} \Psi_n^*(\mathbf{r}) \hat{\mathcal{H}} \Psi_n(\mathbf{r})}{\int d\mathbf{r} \Psi_n^*(\mathbf{r}) \Psi_n(\mathbf{r})}$$

which gives the energy of state n .




“ Machine learning is the science of getting computers to act without being explicitly programmed. ”

Feed-forward Neural Network



Restricted Boltzmann machines



Methods

Exploit the variational principle in order to obtain the ground state energy

$$\begin{aligned} E_0 < E_{\text{VMC}} &= \frac{\int d\mathbf{r} \Psi_T(\mathbf{r})^* \hat{\mathcal{H}} \Psi_T(\mathbf{r})}{\int d\mathbf{r} \Psi_T(\mathbf{r})^* \Psi_T(\mathbf{r})} \\ &= \int d\mathbf{r} \underbrace{\frac{\Psi_T(\mathbf{r})^* \Psi_T(\mathbf{r})}{\int d\mathbf{r} \Psi_T(\mathbf{r})^* \Psi_T(\mathbf{r})}}_{P(\mathbf{r})} \cdot \underbrace{\frac{1}{\Psi_T(\mathbf{r})} \hat{\mathcal{H}} \Psi_T(\mathbf{r})}_{E_L(\mathbf{r})} \end{aligned}$$



We attempt to solve the integral by sampling from the probability density function $P(\mathbf{r})$

$$\begin{aligned} E_{\text{VMC}} &= \int d\mathbf{r} E_L(\mathbf{r}) P(\mathbf{r}) \\ &\approx \frac{1}{M} \sum_{i=1}^M E_L(\mathbf{r}_i) \end{aligned}$$

$$P(\mathbf{r}) \propto \Psi_T(\mathbf{r})^* \Psi_T(\mathbf{r})$$

Use the Slater-Jastrow function as our trial wave function

$$\Psi_T(\mathbf{r}) = |\hat{D}(\mathbf{r})|J(\mathbf{r})$$

where the Slater matrix, $\hat{D}(\mathbf{r})$, contains all the single-particle functions

$$\hat{D}(\mathbf{r}) = \begin{pmatrix} \phi_1(\mathbf{r}_1) & \phi_2(\mathbf{r}_1) & \dots & \phi_N(\mathbf{r}_1) \\ \phi_1(\mathbf{r}_2) & \phi_2(\mathbf{r}_2) & \dots & \phi_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ \phi_1(\mathbf{r}_N) & \phi_2(\mathbf{r}_N) & \dots & \phi_N(\mathbf{r}_N) \end{pmatrix}$$



The Hermite functions,

$$\phi_n(\mathbf{r}) \propto H_n(\mathbf{r}) \exp\left(-\frac{1}{2}\alpha\omega|\mathbf{r}|^2\right),$$

are used as the single-particle functions for quantum dots in standard VMC. The Gaussian can be factorized out from the Slater determinant.

$$|\hat{D}(\mathbf{r}; \alpha)| \propto \exp\left(-\frac{1}{2}\alpha\omega|\mathbf{r}|^2\right) \begin{vmatrix} H_1(\mathbf{r}_1) & H_2(\mathbf{r}_1) & \dots & H_N(\mathbf{r}_1) \\ H_1(\mathbf{r}_2) & H_2(\mathbf{r}_2) & \dots & H_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ H_1(\mathbf{r}_N) & H_2(\mathbf{r}_N) & \dots & H_N(\mathbf{r}_N) \end{vmatrix}$$



We use the marginal distribution of the visible units as the single-particle functions in the Slater determinant, and see if them can model the correlations

$$\phi_n(\mathbf{r}) \propto H_n(\mathbf{r})P(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W})$$

where $P(\mathbf{r})$ is the marginal distribution of the visible units.

$$|\hat{D}(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W})| \propto P(\mathbf{r}; \mathbf{a}, \mathbf{b}, \mathbf{W}) \begin{vmatrix} H_1(\mathbf{r}_1) & H_2(\mathbf{r}_1) & \dots & H_N(\mathbf{r}_1) \\ H_1(\mathbf{r}_2) & H_2(\mathbf{r}_2) & \dots & H_N(\mathbf{r}_2) \\ \vdots & \vdots & \ddots & \vdots \\ H_1(\mathbf{r}_N) & H_2(\mathbf{r}_N) & \dots & H_N(\mathbf{r}_N) \end{vmatrix}$$



The Jastrow factor is added to account for the correlations
Simple Jastrow factor


$$J(\mathbf{r}; \boldsymbol{\beta}) = \exp \left(\sum_{i=1}^N \sum_{j>i}^N \beta_{ij} r_{ij} \right).$$

Padé-Jastrow factor

$$J(\mathbf{r}; \boldsymbol{\beta}) = \exp \left(\sum_{i=1}^N \sum_{j>i}^N \frac{a_{ij} r_{ij}}{1 + \beta r_{ij}} \right).$$

Results

Computational Cost



Conclusion



- ▶ RBM is able to account for most of the correlations
- ▶ RBM+PJ implies to give a lower ground state energy and model the correlations better than a traditional VMC
- ▶ RBM+SJ is both more expensive and less accurate than its fellow methods, and we see no reason to choose it



- ▶ Repeat the exercise using spherical coordinates - interactions are easier to model in spherical coordinates
- ▶ Check the ability of modeling the three-body correlations, considering nuclear systems
- ▶ Reduce the computational cost

Thank you!