# COVID-19 subject 2456

2021-03-01


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Туре	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP0608-1	single experiment	NA	Saliva	2021-01-01	29.94	B.1.243	99.9%	99.8%

#### Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



#### Saliva 2021-01-01


241 intergenic	4198
1578 orf1ab V438A	677
3037 orf1ab silent	2031
6618 orf1ab G2118D	4344
7833 orf1ab N2523T	2980
8782 orf1ab silent	1482
8841 orf1ab T2859I	1485
10741 orf1ab silent	2040
14408 orf1ab P314L	4032
16647 orf1ab silent	2690
18060 orf1ab silent	2065
20268 orf1ab silent	293
21145 orf1ab L2560F	1932
23403 S D614G	23389
23604 S P681H	8894
23905 S silent	2350
24076 S silent	2035
28144 orf8 S84L	9567
28854 N S194L	1683
29266 N silent	8920
29710 intergenic	296
	VSP0608-1




#### Analyses of individual experiments and composite results

### VSP0608-1 | 2021-01-01 | Saliva | 2456 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	2.3.3
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.0.0
tidyverse	1.2.1
ShortRead	1.34.2
$\operatorname{GenomicAlignments}$	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
$\operatorname{GenomeInfoDb}$	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1