

# COVID-19 subject UPHS-0488

*2021-06-23*

The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with  $> 90\%$  sequence coverage.

Table 1. Sample summary.

| Experiment | Type              | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage ( $\geq 5$ reads) |
|------------|-------------------|---------|-------------|-------------|---------------------|---------|-------------------------|---|
| VSP1614-1  | single experiment | NA      | NA          | 2021-03-20  | 22.98               | B.1.575 | 99.9%                   | 99.6%                                     |

## Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



|                        | NA<br>2021-03-20 |  |
|------------------------|------------------|--|
| 241 intergenic         | 4280             |  |
| 1059 ORF1ab T265I      | 1161             |  |
| 1180 ORF1ab silent     | 1580             |  |
| 2305 ORF1ab K680N      | 4208             |  |
| 3037 ORF1ab silent     | 3197             |  |
| 3340 ORF1ab silent     | 7201             |  |
| 4504 ORF1ab E1413D     | 8416             |  |
| 5462 ORF1ab S1733G     | 5959             |  |
| 6195 ORF1ab P1977L     | 7227             |  |
| 6513 ORF1ab del 3      | 710              |  |
| 6730 ORF1ab silent     | 1264             |  |
| 9190 ORF1ab silent     | 11905            |  |
| 9319 ORF1ab silent     | 4411             |  |
| 10029 ORF1ab T3255I    | 437              |  |
| 11514 ORF1ab T3750I    | 11680            |  |
| 14408 ORF1ab P314L     | 11751            |  |
| 16375 ORF1ab P970S     | 8588             |  |
| 19999 ORF1ab V2178F    | 7169             |  |
| 20748 ORF1ab silent    | 15574            |  |
| 23042 S S494P          | 3384             |  |
| 23403 S D614G          | 14191            |  |
| 23604 S P681H          | 11178            |  |
| 23709 S T716I          | 11388            |  |
| 25563 ORF3a Q57H       | 7128             |  |
| 26767 M I82T           | 4352             |  |
| 28271 intergenic del 1 | 6825             |  |
| 28655 N D128Y          | 11994            |  |
| 28851 N S193I          | 728              |  |
| 28887 N T205I          | 701              |  |
| 29711 intergenic del 1 | 6545             |  |
| 29719 intergenic       | 5543             |  |
|                        | VSP1614-1        |  |


Base change

- Expected
- A
- T
- C
- G
- N
- Ins/Del
- No data


# Analyses of individual experiments and composite results

VSP1614-1 | 2021-03-20 | NA | UPHS-0488 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according to variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

| Software/R package   | Version   |
|----------------------|---|
| R                    | 3.4.0   |
| bwa                  | 0.7.17-r1198-dirty                                      |
| samtools             | 1.10 Using htlib 1.10                                   |
| bcftools             | 1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3 |
| pangolin             | 3.1.3   |
| genbankr             | 1.4.0   |
| optparse             | 1.6.0   |
| forcats              | 0.3.0   |
| stringr              | 1.4.0   |
| dplyr                | 0.8.1   |
| purrr                | 0.2.5   |
| readr                | 1.1.1   |
| tidyr                | 0.8.1   |
| tibble               | 2.1.2   |
| ggplot2              | 3.3.3   |
| tidyverse            | 1.2.1   |
| ShortRead            | 1.34.2  |
| GenomicAlignments    | 1.12.2  |
| SummarizedExperiment | 1.6.5   |
| DelayedArray         | 0.2.7   |
| matrixStats          | 0.54.0  |
| Biobase              | 2.36.2  |
| Rsamtools            | 1.28.0  |
| GenomicRanges        | 1.28.6  |
| GenomeInfoDb         | 1.12.3  |
| Biostrings           | 2.44.2  |
| XVector              | 0.16.0  |
| IRanges              | 2.10.5  |
| S4Vectors            | 0.14.7  |
| BiocParallel         | 1.10.1  |
| BiocGenerics         | 0.22.1  |