COVID-19 subject Lennon_canine1

2021-03-08


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Туре	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP0909-1	single experiment	NA	Canine	2021-03-01	26.97	B.1.509	99.8%	99.4%

Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



Canine 2021-03-01


241 intergenic	8684
1059 orf1ab T265I	24787
1385 orf1ab H374Y	22741
1738 orf1ab silent	1260
3037 orf1ab silent	6699
3242 orf1ab G993S	7482
8734 orf1ab silent	38
8782 orf1ab silent	41
9365 orf1ab silent	633
10982 orf1ab K3573E	219
11665 orf1ab silent	47070
11916 orf1ab S3884L	33504
14408 orf1ab P314L	20577
18060 orf1ab silent	6301
23403 S D614G	60220
25449 orf3a E19D	31293
25563 orf3a Q57H	27498
27431 orf7a A13V	8720
28087 orf8 A65V	25065
28144 orf8 S84L	7593
28373 N G34R	69569
28881 N R203K	239
	09–1
	SP0909-1




Analyses of individual experiments and composite results

VSP0909-1 | 2021-03-01 | Canine | Lennon_canine1 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	2.3.3
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.0.0
tidyverse	1.2.1
ShortRead	1.34.2
$\operatorname{GenomicAlignments}$	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
$\operatorname{GenomeInfoDb}$	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1