# COVID-19 subject UPHS-0714

2021-06-23


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Туре	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP1932-1	single experiment	NA	Saliva	2021-03-18	10.56	NA	92.5%	92.2%

#### Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



#### Saliva 2021-03-18


	2021 00 10
1059 ORF1ab T265I	207
3037 ORF1ab silent	125
6101 ORF1ab G1946S	194
8809 ORF1ab silent	54
9867 ORF1ab L3201P	56
11075 ORF1ab F3604L	151
11288 ORF1ab del 9	130
14408 ORF1ab P314L	50
16500 ORF1ab Q1011H	1073
20262 ORF1ab silent	393
21575 S L5F	25
21618 S T19I	25
21846 S T95I	337
22320 S D253G	100
22992 S S477N	67
23403 S D614G	963
24432 S Q957R	147
27739 ORF7a L116F	990
27925 ORF8 T11I	537
28271 intergenic del 1	694
28311 N P13L	745
28879 N S202R	362
	VSP1932-1




### Analyses of individual experiments and composite results

#### VSP1932-1 | 2021-03-18 | Saliva | UPHS-0714 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

Software/R package	Version				
R	3.4.0				
bwa	0.7.17-r1198-dirty				
samtools	1.10 Using htslib 1.10				
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3				
pangolin	3.1.3				
genbankr	1.4.0				
optparse	1.6.0				
forcats	0.3.0				
stringr	1.4.0				
dplyr	0.8.1				
purrr	0.2.5				
readr	1.1.1				
tidyr	0.8.1				
tibble	2.1.2				
ggplot2	3.3.3				
tidyverse	1.2.1				
ShortRead	1.34.2				
GenomicAlignments	1.12.2				
${\bf Summarized Experiment}$	1.6.5				
DelayedArray	0.2.7				
matrixStats	0.54.0				
Biobase	2.36.2				
Rsamtools	1.28.0				
GenomicRanges	1.28.6				
GenomeInfoDb	1.12.3				
Biostrings	2.44.2				
XVector	0.16.0				
IRanges	2.10.5				
S4Vectors	0.14.7				
BiocParallel	1.10.1				
BiocGenerics	0.22.1				