

COVID-19 subject JHV54

2021-06-23


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with $> 90\%$ sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (≥ 5 reads)
VSP9971	composite	NA	Negative control	2021-01-30	NA	NA	8.0%	4.4%
VSP9971-1	single experiment	NA	Negative control	2021-01-30	NA	NA	6.3%	2.9%
VSP9971-2	single experiment	NA	Negative control	2021-01-30	NA	NA	5.8%	3.7%


Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.





Negative control
2021-01-30



7626 ORF1ab N2454S



10665 ORF1ab V3467A



29665 ORF10 silent



Base change




VSP9971-1

VSP9971-2


Analyses of individual experiments and composite results

VSP9971 | 2021-01-30 | Negative control | JHV54 | composite result

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according to variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




No contig data available.

VSP9971-1 | 2021-01-30 | Negative control | JHV54 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




No contig data available.

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



No contig data available.

Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htlib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3
pangolin	3.1.3
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.3.3
tidyverse	1.2.1
ShortRead	1.34.2
GenomicAlignments	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
GenomeInfoDb	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1