COVID-19 subject UPHS-0545

2021-06-03


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage $(>= 5 \text{ reads})$
VSP1671-1	single experiment	NA	NA	2021-03-22	29.82	B.1.526	99.7%	99.7%

Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



NA 2021-03-22


	2021-03-22
241 intergenic	526
1059 ORF1ab T265I	1939
3037 ORF1ab silent	1769
9207 ORF1ab S2981F	1475
9867 ORF1ab L3201P	1587
14408 ORF1ab P314L	1554
16500 ORF1ab Q1011H	2803
20262 ORF1ab silent	526
21575 S L5F	629
21846 S T95I	1237
22320 S D253G	355
23012 S E484K	489
23403 S D614G	5325
23664 S A701V	2115
25489 ORF3a A33S	1033
25517 ORF3a P42L	781
25563 ORF3a Q57H	1548
27019 M K166R	5890
27925 ORF8 T11I	1376
28271 intergenic del 1	1320
28869 N P199L	291
28975 N M234I	364
	71–1
	VSP1671-1
	»




Analyses of individual experiments and composite results

VSP1671-1 | 2021-03-22 | NA | UPHS-0545 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	2.3.8
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.3.3
tidyverse	1.2.1
ShortRead	1.34.2
${\it Genomic Alignments}$	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
$\operatorname{GenomeInfoDb}$	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1