# COVID-19 subject SARS_CoV_306

*2021-06-29*

The table below provides a summary of subject samples for which sequencing data is available.
The experiments column shows the number of sequencing experiments performed for each specimen.
Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

| Experiment | Type | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage (>= 5 reads) |
|---|---|---|---|---|---|---|---|---|
| VSP3097-1 | single experiment | NA | NA | 2021-04-27 | 29.85 | B.1.1.7 | 99.8% | 99.8% |

### Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/NC_04551 are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in $> 50\%$ of read pairs and the variant yields a PHRED score $> 20$. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.




NA
2021−04−27
241 intergenic
820 ORF1ab silent
913 ORF1ab silent
1103 ORF1ab I280L
3037 ORF1ab silent
3267 ORF1ab T1001I
5175 ORF1ab T1637I
5388 ORF1ab A1708D
5986 ORF1ab silent
6954 ORF1ab I2230T
11288 ORF1ab del 9
14408 ORF1ab P314L
14676 ORF1ab silent
15096 ORF1ab silent
15279 ORF1ab silent
16176 ORF1ab silent
21765 S del 6
21991 S del 3
23063 S N501Y
23271 S A570D
23403 S D614G
23604 S P681H
23709 S T716I
24506 S S982A
24742 S silent
24912 S T1117I
24914 S D1118H
25785 ORF3a W131C
26528 M silent
27972 ORF8 Q27stop
28048 ORF8 R52I
28095 ORF8 K68stop
28111 ORF8 Y73C
28271 intergenic del 1
28280 N D3L
28281 N D3L
28282 N D3L
28881 N R203K
28882 N R203K
28883 N G204R
28977 N S235F
29272 N silent
29764 intergenic
3495
4449
4481
1950
2928
2640
2010
2938
1866
828
3223
4034
2784
3183
5500
5400
2360
746
2685
3945
4476
4844
3915
1955
3333
3748
3848
5516
598
10835
9894
8479
8036
7460
5763
5763
5865
653
651
656
468
6701
12130
VSP3097−1
Base change
Expected
A
T
C
G
N
Ins/Del
No data

## Analyses of individual experiments and composite results

### VSP3097-1 | 2021-04-27 | NA | SARS_CoV_306 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

| Software/R package | Version |
|---|---|
| R | 3.4.0 |
| bwa | 0.7.17-r1198-dirty |
| samtools | 1.10 Using htslib 1.10 |
| bcftools | 1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3 |
| pangolin | 3.1.3 |
| genbankr | 1.4.0 |
| optparse | 1.6.0 |
| forcats | 0.3.0 |
| stringr | 1.4.0 |
| dplyr | 0.8.1 |
| purrr | 0.2.5 |
| readr | 1.1.1 |
| tidyr | 0.8.1 |
| tibble | 2.1.2 |
| ggplot2 | 3.3.3 |
| tidyverse | 1.2.1 |
| ShortRead | 1.34.2 |
| GenomicAlignments | 1.12.2 |
| SummarizedExperiment | 1.6.5 |
| DelayedArray | 0.2.7 |
| matrixStats | 0.54.0 |
| Biobase | 2.36.2 |
| Rsamtools | 1.28.0 |
| GenomicRanges | 1.28.6 |
| GenomeInfoDb | 1.12.3 |
| Biostrings | 2.44.2 |
| XVector | 0.16.0 |
| IRanges | 2.10.5 |
| S4Vectors | 0.14.7 |
| BiocParallel | 1.10.1 |
| BiocGenerics | 0.22.1 |