

COVID-19 subject 211-TCE

2021-06-23

The table below provides a summary of subject samples for which sequencing data is available.

The experiments column shows the number of sequencing experiments performed for each specimen.


Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with $> 90\%$ sequence coverage.

Table 1. Sample summary.

| Experiment | Type | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage (≥ 5 reads) |
|------------|-------------------|---------|-------------|-------------|---------------------|---------|-------------------------|---|
| VSP0184-1m | single experiment | NA | NP | 2020-04-08 | 29.86 | B.1 | 100.0% | 99.9% |
| VSP0185-1m | single experiment | NA | OP | 2020-04-13 | 29.88 | B.1 | 99.9% | 99.8% |
| VSP0186-1m | single experiment | NA | ETA | 2020-04-13 | 29.96 | B.1 | 99.9% | 99.8% |

Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



| | ETA 2020-04-13 | NP 2020-04-08 | OP 2020-04-13 | |
|---------------------|-------------------|------------------|------------------|--|
| 241 intergenic | 1938 | 2184 | 2035 | |
| 1059 ORF1ab T265I | 1314 | 4984 | 5385 | |
| 3037 ORF1ab silent | 1199 | 2086 | 3406 | |
| 4113 ORF1ab A1283V | 2514 | 3703 | 3527 | |
| 6285 ORF1ab T2007I | 1869 | 3554 | 263 | |
| 9615 ORF1ab T3117I | 667 | 123 | 1329 | |
| 11916 ORF1ab S3884L | 1165 | 980 | 2590 | |
| 14408 ORF1ab P314L | 1062 | 1416 | 3291 | |
| 18998 ORF1ab A1844V | 1341 | 2846 | 217 | |
| 23403 S D614G | 4146 | 3933 | 3124 | |
| 24197 S A879S | 1106 | 4543 | 4675 | |
| 25563 ORF3a Q57H | 1329 | 4824 | 5037 | |
| 29540 intergenic | 760 | 1223 | 889 | |
| | VSP0186-1m | VSP0184-1m | VSP0185-1m | |


Base change

- Expected
- A
- T
- C
- G
- N
- Ins/Del
- No data


Analyses of individual experiments and composite results

VSP0184-1m | 2020-04-08 | NP | 5 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according to variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

| Software/R package | Version |
|----------------------|---|
| R | 3.4.0 |
| bwa | 0.7.17-r1198-dirty |
| samtools | 1.10 Using htlib 1.10 |
| bcftools | 1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3 |
| pangolin | 3.1.3 |
| genbankr | 1.4.0 |
| optparse | 1.6.0 |
| forcats | 0.3.0 |
| stringr | 1.4.0 |
| dplyr | 0.8.1 |
| purrr | 0.2.5 |
| readr | 1.1.1 |
| tidyr | 0.8.1 |
| tibble | 2.1.2 |
| ggplot2 | 3.3.3 |
| tidyverse | 1.2.1 |
| ShortRead | 1.34.2 |
| GenomicAlignments | 1.12.2 |
| SummarizedExperiment | 1.6.5 |
| DelayedArray | 0.2.7 |
| matrixStats | 0.54.0 |
| Biobase | 2.36.2 |
| Rsamtools | 1.28.0 |
| GenomicRanges | 1.28.6 |
| GenomeInfoDb | 1.12.3 |
| Biostrings | 2.44.2 |
| XVector | 0.16.0 |
| IRanges | 2.10.5 |
| S4Vectors | 0.14.7 |
| BiocParallel | 1.10.1 |
| BiocGenerics | 0.22.1 |