COVID-19 subject SARS_CoV_107

2021-06-29


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Туре	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP3020-1	single experiment	NA	NA	2021-02-10	29.83	B.1	99.8%	99.8%

Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/NC_0455) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



NA 2021-02-10


203 intergenic	1044
222 intergenic	1040
241 intergenic	1043
5974 ORF1ab silent	1283
10029 ORF1ab T3255I	746
10954 ORF1ab silent	1492
11117 ORF1ab I3618V	698
12789 ORF1ab T4175I	3274
14408 ORF1ab P314L	2204
15765 ORF1ab silent	1249
17550 ORF1ab silent	3128
19839 ORF1ab silent	1460
21306 ORF1ab silent	692
22995 S T478K	262
23403 S D614G	3230
23604 S P681H	1809
23756 S T732A	873
27722 ORF7a I110T	2178
27904 ORF8 L4P	4513
28087 ORF8 A65V	3260
29197 N silent	2435
	VSP3020-1
	S>




Analyses of individual experiments and composite results

VSP3020-1 | 2021-02-10 | NA | SARS_CoV_107 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version				
R	3.4.0				
bwa	0.7.17-r1198-dirty				
samtools	1.10 Using htslib 1.10				
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3				
pangolin	3.1.3				
genbankr	1.4.0				
optparse	1.6.0				
forcats	0.3.0				
stringr	1.4.0				
dplyr	0.8.1				
purrr	0.2.5				
readr	1.1.1				
tidyr	0.8.1				
tibble	2.1.2				
ggplot2	3.3.3				
tidyverse	1.2.1				
ShortRead	1.34.2				
GenomicAlignments	1.12.2				
${\bf Summarized Experiment}$	1.6.5				
DelayedArray	0.2.7				
matrixStats	0.54.0				
Biobase	2.36.2				
Rsamtools	1.28.0				
GenomicRanges	1.28.6				
GenomeInfoDb	1.12.3				
Biostrings	2.44.2				
XVector	0.16.0				
IRanges	2.10.5				
S4Vectors	0.14.7				
BiocParallel	1.10.1				
BiocGenerics	0.22.1				