# COVID-19 subject UPHS-0323

*2021-04-17*

The table below provides a summary of subject samples for which sequencing data is available.
The experiments column shows the number of sequencing experiments performed for each specimen.
Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

| Experiment | Type | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage (>= 5 reads) |
|---|---|---|---|---|---|---|---|---|
| VSP1368-1 | single experiment | NA | Saline | 2021-03-22 | 29.85 | B.1.1.7 | 99.9% | 99.8% |

**Variants shared across samples**

The heat map below shows how variants (reference genome /home/everett/projects/SARS-CoV-2-Philadelphia/USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.




Saline
2021−03−22
241 intergenic
913 orf1ab silent
2110 orf1ab silent
3004 orf1ab E913D
3037 orf1ab silent
3267 orf1ab T1001I
5388 orf1ab A1708D
5986 orf1ab silent
6954 orf1ab I2230T
7042 orf1ab M2259I
8025 orf1ab A2587V
8782 orf1ab silent
11288 orf1ab del 9
12499 orf1ab silent
14120 orf1ab P218L
14408 orf1ab P314L
14676 orf1ab silent
15279 orf1ab silent
16176 orf1ab silent
17766 orf1ab silent
18060 orf1ab silent
21765 S del 6
21991 S del 3
23063 S N501Y
23271 S A570D
23403 S D614G
23604 S P681H
23709 S T716I
24506 S S982A
24914 S D1118H
27401 orf7a I3T
27972 orf8 Q27stop
28048 orf8 R52I
28095 orf8 K68stop
28111 orf8 Y73C
28144 orf8 S84L
28271 intergenic del 1
28280 N D3L
28281 N D3L
28282 N D3L
28881 N R203K
28882 N R203K
28883 N G204R
28977 N S235F
1455
4805
1955
2241
1950
3081
3739
1783
724
2343
7926
3268
2880
3968
3433
2160
3354
6013
7094
1923
2837
1691
1207
191
4311
4658
3285
2708
3605
5183
2948
2701
2663
4276
4640
5153
3257
2157
2157
2335
559
555
557
1234
VSP1368−1
Base change
Expected
A
T
C
G
N
Ins/Del
No data

## Analyses of individual experiments and composite results

### VSP1368-1 | 2021-03-22 | Saline | UPHS-0323 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

| Software/R package | Version |
|---|---|
| R | 3.4.0 |
| bwa | 0.7.17-r1198-dirty |
| samtools | 1.10 Using htslib 1.10 |
| bcftools | 1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3 |
| pangolin | 2.3.8 |
| genbankr | 1.4.0 |
| optparse | 1.6.0 |
| forcats | 0.3.0 |
| stringr | 1.4.0 |
| dplyr | 0.8.1 |
| purrr | 0.2.5 |
| readr | 1.1.1 |
| tidyr | 0.8.1 |
| tibble | 2.1.2 |
| ggplot2 | 3.0.0 |
| tidyverse | 1.2.1 |
| ShortRead | 1.34.2 |
| GenomicAlignments | 1.12.2 |
| SummarizedExperiment | 1.6.5 |
| DelayedArray | 0.2.7 |
| matrixStats | 0.54.0 |
| Biobase | 2.36.2 |
| Rsamtools | 1.28.0 |
| GenomicRanges | 1.28.6 |
| GenomeInfoDb | 1.12.3 |
| Biostrings | 2.44.2 |
| XVector | 0.16.0 |
| IRanges | 2.10.5 |
| S4Vectors | 0.14.7 |
| BiocParallel | 1.10.1 |
| BiocGenerics | 0.22.1 |