

COVID-19 subject 219

2020-08-13


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report.

Table 1. Sample summary.

Experiment	Type	Input genomes	Sample type	Sample date	Largest contig (KD)	Reference read coverage	Reference read coverage (>= 5 reads)
VSP0017	composite	NA	ETA	4/17/2020	7.66	93.1%	92.0%
VSP0017-1a	single experiment	1870	ETA	4/17/2020	0.47	19.4%	11.1%
VSP0017-1b	single experiment	1870	ETA	4/17/2020	0.40	22.9%	20.4%
VSP0017-2a	single experiment	1870	ETA	4/17/2020	1.23	45.9%	42.8%
VSP0017-2b	single experiment	1870	ETA	4/17/2020	NA	NA	NA
VSP0017-3	single experiment	9350	ETA	4/17/2020	1.53	48.6%	40.2%
VSP0017-4	single experiment	1870	ETA	4/17/2020	1.05	49.1%	46.4%
VSP0017-5	single experiment	1870	ETA	4/17/2020	1.69	43.2%	38.4%
VSP0017-6	single experiment	1870	ETA	4/17/2020	3.40	64.1%	61.6%

Variants shared across samples


The heat map below shows how variants are shared across subject samples. The quality scores are PHRED scaled values [$Q = -10\log_{10}(\text{error rate})$] where a score of 30 represents a probability of 99.9% that a variant is called correctly and a score of 50 represents a probability of 99.999%. Gray tiles denote that 10 or more reads covered the variant position and the reference base was observed. Tiles are omitted if there are less than 10 reads covering a variant position.




Analyses of individual experiments and composite results.

VSP0017 | 4/17/2020 | ETA | 219-tri | composite result

The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




VSP0017-2b | 4/17/2020 | ETA | 219-tri | 1870 genomes | single experiment


No pileup data available.

No contig data available.


The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



The plot below shows the number of reads covering each nucleotide position in the reference genome (USA-WA1-2020). Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.

