# COVID-19 subject UK strain SARS-CoV-2

*2021-05-05*

The table below provides a summary of subject samples for which sequencing data is available.
The experiments column shows the number of sequencing experiments performed for each specimen.
Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

| Experiment | Type | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage (>= 5 reads) |
|---|---|---|---|---|---|---|---|---|
| VSP1975-1 | single experiment | NA | RNA | 2021-04-07 | 29.86 | B.1.1.7 | 99.9% | 99.7% |

**Variants shared across samples**

The heat map below shows how variants (reference genome /home/everett/projects/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.




RNA
2021−04−07
241 intergenic
913 ORF1ab silent
3037 ORF1ab silent
3267 ORF1ab T1001I
4999 ORF1ab silent
5388 ORF1ab A1708D
5986 ORF1ab silent
6633 ORF1ab A2123V
6954 ORF1ab I2230T
11288 ORF1ab del 9
13339 ORF1ab N4358K
14408 ORF1ab P314L
14676 ORF1ab silent
15279 ORF1ab silent
16176 ORF1ab silent
17615 ORF1ab K1383R
19079 ORF1ab E1871G
19854 ORF1ab silent
21765 S del 6
21991 S del 3
22015 S S151R
23014 S E484D
23063 S N501Y
23271 S A570D
23403 S D614G
23604 S P681H
23709 S T716I
24506 S S982A
24914 S D1118H
27972 ORF8 Q27stop
28048 ORF8 R52I
28111 ORF8 Y73C
28271 intergenic del 1
28280 N D3L
28281 N D3L
28282 N D3L
28881 N R203K
28882 N R203K
28883 N G204R
28977 N S235F
647
2465
1631
699
1360
3227
352
1367
529
1960
8749
1342
1170
2165
2974
2480
2064
1477
722
307
307
1744
1679
2485
2591
3075
1310
944
3221
1503
1563
1401
1750
1522
1522
1554
1919
1908
1919
1601
VSP1975−1
Base change
Expected
A
T
C
G
N
Ins/Del
No data

## Analyses of individual experiments and composite results

### VSP1975-1 | 2021-04-07 | RNA | UK strain SARS-CoV-2 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

| Software/R package | Version |
|---|---|
| R | 3.4.0 |
| bwa | 0.7.17-r1198-dirty |
| samtools | 1.10 Using htslib 1.10 |
| bcftools | 1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3 |
| pangolin | 2.3.8 |
| genbankr | 1.4.0 |
| optparse | 1.6.0 |
| forcats | 0.3.0 |
| stringr | 1.4.0 |
| dplyr | 0.8.1 |
| purrr | 0.2.5 |
| readr | 1.1.1 |
| tidyr | 0.8.1 |
| tibble | 2.1.2 |
| ggplot2 | 3.0.0 |
| tidyverse | 1.2.1 |
| ShortRead | 1.34.2 |
| GenomicAlignments | 1.12.2 |
| SummarizedExperiment | 1.6.5 |
| DelayedArray | 0.2.7 |
| matrixStats | 0.54.0 |
| Biobase | 2.36.2 |
| Rsamtools | 1.28.0 |
| GenomicRanges | 1.28.6 |
| GenomeInfoDb | 1.12.3 |
| Biostrings | 2.44.2 |
| XVector | 0.16.0 |
| IRanges | 2.10.5 |
| S4Vectors | 0.14.7 |
| BiocParallel | 1.10.1 |
| BiocGenerics | 0.22.1 |