COVID-19 subject UPHS-0612

2021-06-01

The table below provides a summary of subject samples for which sequencing data is available.
The experiments column shows the number of sequencing experiments performed for each specimen.
Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP1797-1	single experiment	NA	Saline	2021-03-24	29.80	B.1.2	99.7%	99.5%

Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in $> 50\%$ of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.


Saline
2021−03−24
241 intergenic
65
835 ORF1ab silent
230
1059 ORF1ab T265I
81
2539 ORF1ab silent
135
3037 ORF1ab silent
158
3497 ORF1ab A1078T
189
4261 ORF1ab silent
200
6063 ORF1ab D1933G
110
6618 ORF1ab G2118D
344
9207 ORF1ab S2981F
150
10319 ORF1ab L3352F
149
11083 ORF1ab L3606F
86
14408 ORF1ab P314L
249
18060 ORF1ab silent
111
18424 ORF1ab N1653D
206
21304 ORF1ab R2613C
284
23403 S D614G
331
25563 ORF3a Q57H
137
25907 ORF3a G172V
147
27964 ORF8 S24L
384
28472 N P67S
236
28869 N P199L
51
VSP1797−1
Base change
Expected
A
T
C
G
N
Ins/Del
No data

Analyses of individual experiments and composite results

VSP1797-1 | 2021-03-24 | Saline | UPHS-0612 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	2.3.8
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.3.3
tidyverse	1.2.1
ShortRead	1.34.2
GenomicAlignments	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
GenomeInfoDb	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1