

COVID-19 subject UPHS-0132

2021-03-29

The table below provides a summary of subject samples for which sequencing data is available.

The experiments column shows the number of sequencing experiments performed for each specimen.


Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with $> 90\%$ sequence coverage.

Table 1. Sample summary.

| Experiment | Type | Genomes | Sample type | Sample date | Largest contig (KD) | Lineage | Reference read coverage | Reference read coverage (≥ 5 reads) |
|------------|-------------------|---------|-------------|-------------|---------------------|---------|-------------------------|---|
| VSP1117-1 | single experiment | NA | VTM | 2021-03-15 | 11.56 | B.1.166 | 98.5% | 90.1% |

Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



VTM

5388 orf1ab A1708D

25

11288 orf1ab del 9

21

14408 orf1ab P314L

21

14676 orf1ab silent

14

15096 orf1ab silent

20

16176 orf1ab silent

23

23271 S A570D

25

23403 S D614G

15

23604 S P681H

26

23709 S T716I

18

24506 S S982A

15

24914 S D1118H

70

25785 orf3a W131C

14

27972 orf8 Q27stop

17

28048 orf8 R52I

24

28095 orf8 K68stop

24

28111 orf8 Y73C

20

29764 intergenic

23

Base change

Expected

A

T

C

G

N

Ins/Del


No data

VSP1117-1


Analyses of individual experiments and composite results

VSP1117-1 | 2021-03-15 | VTM | UPHS-0132 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

| Software/R package | Version |
|----------------------|---|
| R | 3.4.0 |
| bwa | 0.7.17-r1198-dirty |
| samtools | 1.10 Using htlib 1.10 |
| bcftools | 1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3 |
| pangolin | 2.3.3 |
| genbankr | 1.4.0 |
| optparse | 1.6.0 |
| forcats | 0.3.0 |
| stringr | 1.4.0 |
| dplyr | 0.8.1 |
| purrr | 0.2.5 |
| readr | 1.1.1 |
| tidyr | 0.8.1 |
| tibble | 2.1.2 |
| ggplot2 | 3.0.0 |
| tidyverse | 1.2.1 |
| ShortRead | 1.34.2 |
| GenomicAlignments | 1.12.2 |
| SummarizedExperiment | 1.6.5 |
| DelayedArray | 0.2.7 |
| matrixStats | 0.54.0 |
| Biobase | 2.36.2 |
| Rsamtools | 1.28.0 |
| GenomicRanges | 1.28.6 |
| GenomeInfoDb | 1.12.3 |
| Biostrings | 2.44.2 |
| XVector | 0.16.0 |
| IRanges | 2.10.5 |
| S4Vectors | 0.14.7 |
| BiocParallel | 1.10.1 |
| BiocGenerics | 0.22.1 |