COVID-19 subject AHT2YGAFX2-1

2021-03-25


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Туре	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP9961-1	single experiment	NA	Negative control	2021-03-22	29.84	B.1.234	99.8%	99.8%

Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



Negative control 2021–03–22


241 intergenic	1835
1738 orf1ab silent	4143
2258 orf1ab V665I	2194
3037 orf1ab silent	4453
4596 orf1ab T1444N	4531
5812 orf1ab silent	8643
6441 orf1ab K2059R	8360
8140 orf1ab silent	5114
8782 orf1ab silent	4370
9204 orf1ab D2980G	6750
14408 orf1ab P314L	5686
18060 orf1ab silent	4226
20268 orf1ab silent	1227
21575 S L5F	1724
23202 S T547K	1613
23403 S D614G	6034
28144 orf8 S84L	2119
28854 N S194L	765
29384 N D371Y	1593
29445 N T391I	1907
	VSP9961-1




Analyses of individual experiments and composite results

$VSP9961-1 \mid 2021-03-22 \mid Negative\ control \mid AHT2YGAFX2-1 \mid genomes \mid single\ experiment$


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	2.3.3
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.0.0
tidyverse	1.2.1
ShortRead	1.34.2
$\operatorname{GenomicAlignments}$	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
$\operatorname{GenomeInfoDb}$	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1