

COVID-19 subject 209

2021-01-19


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. The code base for this analysis can be found ([here](#)).

Table 1. Sample summary.

Experiment	Type	Input genomes	Sample type	Sample date	Largest contig (KD)	Reference read coverage	Reference read coverage (≥ 5 reads)
VSP0011	composite	NA	ETA	2020-04-08	29.90	99.8%	99.6%
VSP0011-1m	single experiment	NA	ETA	2020-04-08	0.72	52.0%	30.9%
VSP0011-2	single experiment	2150000	ETA	2020-04-08	29.90	99.8%	99.6%

Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



ETA
2020-04-08

241 intergenic	0	4084
1059 orf1ab T265I	97	1033
3037 orf1ab silent	16	390
3312 orf1ab L1016P	0	19102
8782 orf1ab silent	0	1093
14408 orf1ab P314L	2	364
18060 orf1ab silent	6	133
23403 S D614G	2	44329
25563 orf3a Q57H	86	887
28144 orf8 S84L	8	11411

Base change

- Expected
- A
- T
- C
- G
- N
- Ins/Del
- No data


VSP0011-1m

VSP0011-2


Analyses of individual experiments and composite results.

VSP0011 | 2020-04-08 | ETA | 209-tri | composite result


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according to variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.




The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.




Excerpt from plot above focusing on reads coverage from 0 to 50 NT.




The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.

