

COVID-19 subject HUP-PH-0006

2021-05-05

The table below provides a summary of subject samples for which sequencing data is available.

The experiments column shows the number of sequencing experiments performed for each specimen.


Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with $> 90\%$ sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (≥ 5 reads)
VSP0821-1	single experiment	NA	Saline	2021-02-12	29.87	B.1.243	99.9%	99.8%

Variants shared across samples

The heat map below shows how variants (reference genome `/home/everett/projects/SARS-CoV-2-Philadelphia/Wuhan-Hu-1`) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



Saline
2021-02-12

241 intergenic	4910
527 ORF1ab silent	4978
1578 ORF1ab V438A	5173
3037 ORF1ab silent	9926
6618 ORF1ab G2118D	33869
7833 ORF1ab N2523T	12855
10741 ORF1ab silent	20284
14408 ORF1ab P314L	11062
20268 ORF1ab silent	8451
23403 S D614G	17080
23604 S P681H	12922
24076 S silent	10968
27197 intergenic	17517
28077 ORF8 V62L	11693
28854 N S194L	2083
29266 N silent	5163
29710 intergenic	2768
29764 intergenic	2935

Base change




VSP0821-1


Analyses of individual experiments and composite results

VSP0821-1 | 2021-02-12 | Saline | HUP-PH-0006 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htlib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3
pangolin	2.3.8
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.0.0
tidyverse	1.2.1
ShortRead	1.34.2
GenomicAlignments	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
GenomeInfoDb	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1