

COVID-19 subject 427

2021-01-27

The table below provides a summary of subject samples for which sequencing data is available.

The experiments column shows the number of sequencing experiments performed for each specimen.


Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with $> 90\%$ sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (≥ 5 reads)
VSP0510-2	single experiment	NA	NP-OP	2020-12-04	29.91	B.1.234	99.8%	99.7%

Variants shared across samples

The heat map below shows how variants (reference genome USA-WA1-2020) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.



NP-OP
2020-12-04

241 intergenic	14834
936 orf1ab T224I	14945
1912 orf1ab silent	7055
2258 orf1ab V665I	3194
2644 orf1ab silent	22566
3037 orf1ab silent	8504
5094 orf1ab H1610R	1924
6441 orf1ab K2059R	7465
8140 orf1ab silent	8214
8782 orf1ab silent	6799
9204 orf1ab D2980G	10402
14408 orf1ab P314L	11080
17010 orf1ab silent	25650
18060 orf1ab silent	5480
20268 orf1ab silent	483
23403 S D614G	21299
28144 orf8 S84L	6441
28854 N S194L	2037
29445 N T391I	795

Base change


- Expected
- A
- T
- C
- G
- N
- Ins/Del
- No data

VSP0510-2


Analyses of individual experiments and composite results

VSP0510-2 | 2020-12-04 | NP-OP | 427no | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htlib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htlib 1.10.2-57-gf58a6f3
pangolin	2.1.7
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.0.0
tidyverse	1.2.1
ShortRead	1.34.2
GenomicAlignments	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
GenomeInfoDb	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1