COVID-19 subject UPHS-0409

2021-06-23

The table below provides a summary of subject samples for which sequencing data is available.
The experiments column shows the number of sequencing experiments performed for each specimen.
Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

Experiment	Type	Genomes	Sample type	Sample date	Largest contig (KD)	Lineage	Reference read coverage	Reference read coverage (>= 5 reads)
VSP1535-1	single experiment	NA	NA	2021-03-22	4.82	NA	96.0%	86.2%

Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in $> 50\%$ of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.




NA
2021−03−22
820 ORF1ab silent
913 ORF1ab silent
3267 ORF1ab T1001I
5388 ORF1ab A1708D
11288 ORF1ab del 9
14408 ORF1ab P314L
14676 ORF1ab silent
15279 ORF1ab silent
16176 ORF1ab silent
23063 S N501Y
23271 S A570D
23403 S D614G
24506 S S982A
24914 S D1118H
25785 ORF3a W131C
27972 ORF8 Q27stop
28048 ORF8 R52I
28095 ORF8 K68stop
28111 ORF8 Y73C
28271 intergenic del 1
28280 N D3L
28281 N D3L
28282 N D3L
29272 N silent
29764 intergenic
32
41
24
23
17
18
15
34
47
16
26
28
13
72
17
28
25
24
26
23
19
19
21
33
53
VSP1535−1
Base change
Expected
A
T
C
G
N
Ins/Del
No data

Analyses of individual experiments and composite results

VSP1535-1 | 2021-03-22 | NA | UPHS-0409 | genomes | single experiment

The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



Software environment

Software/R package	Version
R	3.4.0
bwa	0.7.17-r1198-dirty
samtools	1.10 Using htslib 1.10
bcftools	1.10.2-34-g1a12af0-dirty Using htslib 1.10.2-57-gf58a6f3
pangolin	3.1.3
genbankr	1.4.0
optparse	1.6.0
forcats	0.3.0
stringr	1.4.0
dplyr	0.8.1
purrr	0.2.5
readr	1.1.1
tidyr	0.8.1
tibble	2.1.2
ggplot2	3.3.3
tidyverse	1.2.1
ShortRead	1.34.2
GenomicAlignments	1.12.2
SummarizedExperiment	1.6.5
DelayedArray	0.2.7
matrixStats	0.54.0
Biobase	2.36.2
Rsamtools	1.28.0
GenomicRanges	1.28.6
GenomeInfoDb	1.12.3
Biostrings	2.44.2
XVector	0.16.0
IRanges	2.10.5
S4Vectors	0.14.7
BiocParallel	1.10.1
BiocGenerics	0.22.1