# COVID-19 subject UPHS-1011

2021-06-23


The table below provides a summary of subject samples for which sequencing data is available. The experiments column shows the number of sequencing experiments performed for each specimen. Experiment specific analyses are shown at the end of this report. Lineages are called with the Pangolin software tool (Rambaut et al 2020) for genomes with > 90% sequence coverage.

Table 1. Sample summary.

| Experiment | Туре              | Genomes | Sample type | Sample date | Largest contig<br>(KD) | Lineage | Reference read<br>coverage | Reference read coverage (>= 5 reads) |
|------------|-------------------|---------|-------------|-------------|------------------------|---------|----------------------------|--------------------------------------|
| VSP2223-1  | single experiment | NA      | Saline      | 2021-04-10  | 7.35                   | NA      | 88.7%                      | 88.6%                                |

#### Variants shared across samples

The heat map below shows how variants (reference genome /home/common/SARS-CoV-2-Philadelphia/Wuhan-Hu-1) are shared across subject samples where the percent variance is colored. Variants are called if a variant position is covered by 5 or more reads, the alternative base is found in > 50% of read pairs and the variant yields a PHRED score > 20. Gray tiles denote positions where the variant was not the major variant or no variants were found. The relative base compositions of each experiment used to calculate tiles are shown in the following plot where the total number of position reads are shown atop of each plot.




#### Saline 2021-04-10

|                        | 2021-04-10 |
|------------------------|------------|
| 1488 ORF1ab G408V      | 1021       |
| 1594 ORF1ab silent     | 544        |
| 2110 ORF1ab silent     | 4332       |
| 3037 ORF1ab silent     | 704        |
| 3267 ORF1ab T1001I     | 1719       |
| 4470 ORF1ab I1402T     | 2341       |
| 5388 ORF1ab A1708D     | 6132       |
| 5986 ORF1ab silent     | 2501       |
| 6954 ORF1ab I2230T     | 4216       |
| 11288 ORF1ab del 9     | 3421       |
| 11502 ORF1ab G3746V    | 948        |
| 11693 ORF1ab V3810F    | 4716       |
| 13073 ORF1ab del 2     | 2105       |
| 14120 ORF1ab P218L     | 4096       |
| 14408 ORF1ab P314L     | 4639       |
| 14676 ORF1ab silent    | 4400       |
| 15279 ORF1ab silent    | 8163       |
| 16176 ORF1ab silent    | 3256       |
| 19390 ORF1ab P1975S    | 6156       |
| 21765 S del 6          | 854        |
| 21991 S del 3          | 1102       |
| 23063 S N501Y          | 338        |
| 23271 S A570D          | 4530       |
| 23403 S D614G          | 4591       |
| 23604 S P681H          | 1837       |
| 23709 S T716I          | 1316       |
| 24506 S S982A          | 1261       |
| 24914 S D1118H         | 3663       |
| 27247 ORF6 silent      | 4839       |
| 27972 ORF8 Q27stop     | 21846      |
| 28048 ORF8 R52I        | 13722      |
| 28095 ORF8 K68stop     | 13283      |
| 28111 ORF8 Y73C        | 10157      |
| 28271 intergenic del 1 | 1894       |
| 28280 N D3L            | 1137       |
| 28281 N D3L            | 1137       |
| 28282 N D3L            | 1243       |
| 28326 N G18V           | 2261       |
| 28881 N R203K          | 975        |
| 28882 N R203K          | 970        |
| 28883 N G204R          | 971        |
| 28977 N S235F          | 1217       |
|                        | 3–1        |
|                        | ς.         |


### Analyses of individual experiments and composite results

#### VSP2223-1 | 2021-04-10 | Saline | UPHS-1011 | genomes | single experiment


The plot below shows the number of reads covering each nucleotide position in the reference genome. Variants are shown as colored dots along the bottom of the plot and are color coded according by variant types: gray - transgenic, green - silent, gold - missense, red - nonsense, black - indel.



Excerpt from plot above focusing on reads coverage from 0 to 50 NT.



The longest five assembled contigs are shown below colored by their edit distance to the reference genome.



## Software environment

| Software/R package            | Version  |
|-------------------------------|--|
| R                             | 3.4.0  |
| bwa                           | 0.7.17-r1198-dirty   |
| samtools                      | 1.10 Using htslib 1.10                                     |
| bcftools                      | 1.10.2-34-g1a12af0-dirty Using htslib $1.10.2-57-gf58a6f3$ |
| pangolin                      | 3.1.3  |
| genbankr                      | 1.4.0  |
| optparse                      | 1.6.0  |
| forcats                       | 0.3.0  |
| stringr                       | 1.4.0  |
| dplyr                         | 0.8.1  |
| purrr                         | 0.2.5  |
| readr                         | 1.1.1  |
| tidyr                         | 0.8.1  |
| tibble                        | 2.1.2  |
| ggplot2                       | 3.3.3  |
| tidyverse                     | 1.2.1  |
| ShortRead                     | 1.34.2   |
| GenomicAlignments             | 1.12.2   |
| ${\bf Summarized Experiment}$ | 1.6.5  |
| DelayedArray                  | 0.2.7  |
| matrixStats                   | 0.54.0   |
| Biobase                       | 2.36.2   |
| Rsamtools                     | 1.28.0   |
| GenomicRanges                 | 1.28.6   |
| GenomeInfoDb                  | 1.12.3   |
| Biostrings                    | 2.44.2   |
| XVector                       | 0.16.0   |
| IRanges                       | 2.10.5   |
| S4Vectors                     | 0.14.7   |
| BiocParallel                  | 1.10.1   |
| BiocGenerics                  | 0.22.1   |