

New Exhibits for AEJ Revisions

February 17, 2026

Contents

1	TWFE Plots	2
2	Repeated Cross Section Plots	7
3	Decadal 2SLS plots, Panel	11
4	Decadal 2SLS plots, Split	13

List of Figures

1	TWFE, No controls	3
2	TWFE, year X sum of shares controls	4
3	TWFE, year X population growth controls	5
4	TWFE, year X census region FEs	6
5	Repeated Cross Section, Δ munis P.C.	8
6	Repeated Cross Section, $\frac{n_{70}-n_{40}}{Pop_{40}}$	9
7	Repeated Cross Section, Population Dilution Term	9
8	Repeated Cross Section, Population Growth	10
9	Decadal 2SLS, C.Goodman Munis	14
10	Decadal 2SLS, CoG Munis	15
11	Decadal 2SLS, School Districts	16
12	Decadal 2SLS, Special Districts	17
13	Decadal 2SLS, Main City Share	18

List of Tables

1	Panel IV	12
---	--------------------	----

1 TWFE Plots

These figures are the β_τ 's from the TWFE regression

$$y_{it} = \sum_{\tau=1910}^{2010} \beta_\tau \text{Above Med GM}_i \times 1\{t = \tau\} + \gamma_i + \delta_t + \epsilon_{it} \quad (1)$$

Where δ_t and γ_i are decade and CZ fixed effects. Sometimes controls are included.




Figure 1: TWFE, No controls




Figure 2: TWFE, year X sum of shares controls




Figure 3: TWFE, year X population growth controls




Figure 4: TWFE, year X census region FEs

2 Repeated Cross Section Plots

These figures are the β_t 's from running the 2SLS regression of $t - 10$ to t changes y_{it} on 1940-70 GM.

$$y_{it} = \alpha_i + \beta_t GM_i + \mathbf{X}_i' \zeta_t + \epsilon_{it} \quad (2)$$

Where X_i are the covariates in our main specification.




Figure 5: Repeated Cross Section, Δ munis P.C.




Figure 6: Repeated Cross Section, $\frac{n_{70}-n_{40}}{Pop_{40}}$




Figure 7: Repeated Cross Section, Population Dilution Term




Figure 8: Repeated Cross Section, Population Growth

3 Decadal 2SLS plots, Panel

This table comes from creating a separate X and IV for each decade 1940-50, 1950-60, and 1960-1970, then performing the 2SLS regression as a stacked panel.

$$y_{it} = \beta GM_{it} + \mathbf{X}_i' \zeta_t + \delta_t + \epsilon_{it} \tag{3}$$

Table 1: Panel IV

	C. Goodman		Census of Governments		
	Municipalities		School districts	Special Districts	Census
	(1)	(2)	(3)	(4)	Main City Share (5)
Panel A: First Stage $\widehat{GM}_t$					
$\widehat{GM}_t$	0.458 (0.306)	0.458 (0.306)	0.476 (0.302)	0.458 (0.306)	0.458 (0.306)
F-Stat	2.24	2.24	2.49	2.24	2.24
Panel B: IV 1940-70					
GM_t	0.026*** (0.010)	0.028*** (0.010)	0.677*** (0.233)	0.077** (0.035)	-2.543*** (0.678)
Observations	390	390	354	390	390

4 Decadal 2SLS plots, Split

These figures come from creating a separate X and IV for each decade 1940-50, 1950-60, and 1960-1970, then performing the 2SLS regression separately for each year.

$$y_{it} = \beta_t GM_{it} + \mathbf{X}_i' \zeta_t + \delta_t + \epsilon_{it} \quad (4)$$




Figure 9: Decadal 2SLS, C.Goodman Munis




Figure 10: Decadal 2SLS, CoG Munis




Figure 11: Decadal 2SLS, School Districts




Figure 12: Decadal 2SLS, Special Districts




Figure 13: Decadal 2SLS, Main City Share