

ELEMENTARY COMBINATORICS & PROBABILITY

Lecture 12

Expectation of a random variable

LOGISTICS

- Today
 - Graded assignment 5 (pts)
- Tomorrow
 - **FINAL EXAM (25 pts)**
 - ~ 2.5 - 3h
 - Start: 09:30
 - Extra assignment (10 – 15 pts)

LAST TIME

- Probability mass function
- Cumulative distribution function
- Special distributions

TODAY

- Review special distributions
- Expectation and variance
- Review

WARM-UP

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

$$P(AB) \quad P(A) \cdot P(B)$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

$$0.29 = P(AB) \quad P(A) \cdot P(B)$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

$$0.29 = P(AB) \quad P(A) \cdot P(B) = 0.73 \cdot 0.48 = 0.3504$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

$$0.29 = P(AB) \neq P(A) \cdot P(B) = 0.73 \cdot 0.48 = 0.3504 \rightarrow$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Are A and B independent?

$$0.29 = P(AB) \neq P(A) \cdot P(B) = 0.73 \cdot 0.48 = 0.3504 \rightarrow$$

A and B aren't independent.

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) =$$

$$P(B|A) =$$

$$P(A|B) =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) =$$

$$P(B|A) =$$

$$P(A|B) =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.73 + 0.48 - 0.29 = 0.92$$

$$P(B|A) =$$

$$P(A|B) =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.73 + 0.48 - 0.29 = 0.92$$

$$P(B|A) = \frac{P(AB)}{P(A)} =$$

$$P(A|B) =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.73 + 0.48 - 0.29 = 0.92$$

$$P(B|A) = \frac{P(AB)}{P(A)} = \frac{0.29}{0.73}$$

$$P(A|B) =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.73 + 0.48 - 0.29 = 0.92$$

$$P(B|A) = \frac{P(AB)}{P(A)} = \frac{0.29}{0.73}$$

$$P(A|B) = \frac{P(AB)}{P(B)} =$$

TWO EVENTS

- For two events A and B , $P(A) = 0.73$, $P(B) = 0.48$ and $P(A \cap B) = 0.29$. Find...

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.73 + 0.48 - 0.29 = 0.92$$

$$P(B|A) = \frac{P(AB)}{P(A)} = \frac{0.29}{0.73}$$

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{0.29}{0.48}$$

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

$$P(A) = 0.02, \quad P(B) = 0.1$$

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

$$P(A) = 0.02, \quad P(B) = 0.1$$

A and B are independent

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

$$P(A) = 0.02, \quad P(B) = 0.1$$

A and B are independent \rightarrow

$$P(AB) =$$

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

$$P(A) = 0.02, \quad P(B) = 0.1$$

A and B are independent \rightarrow

$$P(AB) = P(A) \cdot P(B) =$$

INDEPENDENT DISEASES

- Disease A occurs in 2% of the population, while 10% suffer from a more common disease B . Assuming that these two conditions are independent, what is the probability that a random person has both?

$$P(A) = 0.02, \quad P(B) = 0.1$$

A and B are independent \rightarrow

$$P(AB) = P(A) \cdot P(B) = 0.02 \cdot 0.1 = 0.002$$

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?
- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) =$$

- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} =$$

- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) =$$

- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.5^6 \cdot 0.5^4$$

- ... that you will get at least one Tails?

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.5^6 \cdot 0.5^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) =$$

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.5^6 \cdot 0.5^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) = 1 - P(0T) =$$

FAIR COIN

- You are tossing a fair coin 10 times. What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = \frac{1}{2^{10}} = \left(\frac{1}{2}\right)^4 \cdot \left(\frac{1}{2}\right)^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.5^6 \cdot 0.5^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) = 1 - P(0T) = 1 - 0.5^{10}$$

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?
- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) =$$

- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?
- ... that you will get at least one Tails?

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) =$$

- ... that you will get at least one Tails?

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.8^6 \cdot 0.2^4$$

- ... that you will get at least one Tails?

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.8^6 \cdot 0.2^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) =$$

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.8^6 \cdot 0.2^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) = 1 - P(0T) =$$

BIASED COIN

- You are tossing a biased coin with $p_{Heads} = 0.8$ ten times.
What is the probability...
- ... that you will get 4 Tails in a row, followed by 6 Heads in a row?

$$P(TTTTHHHHHH) = 0.2^4 \cdot 0.8^6$$

- ... that in total you will get 4 Tails and 6 Heads?

$$P(6H, 4T) = C(10, 6) \cdot 0.8^6 \cdot 0.2^4$$

- ... that you will get at least one Tails?

$$P(\geq 1T) = 1 - P(0T) = 1 - 0.8^{10}$$

BERNOULLI DISTRIBUTION

- Consider a random experiment with two possible outcomes:
“success” (with probability p) or “failure” (with probability $1 - p$)
 - tossing a coin: H or T;
 - a new child: a boy or a girl;
 - you take an exam: pass or fail.
- Consider a random variable $X \sim \text{Bernoulli}(p)$

x	0	1
$P(X = x)$	$1 - p$	p

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

Find the probability that the first person he encounters will be able to speak English.

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

Find the probability that the first person he encounters will be able to speak English.

X – random variable: $X = 1$ – person speaks English, $X = 0$ – doesn't.

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

Find the probability that the first person he encounters will be able to speak English.

X – random variable: $X = 1$ – person speaks English, $X = 0$ – doesn't.

$$X \sim \text{Bernoulli}(\quad) \rightarrow$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

Find the probability that the first person he encounters will be able to speak English.

X – random variable: $X = 1$ – person speaks English, $X = 0$ – doesn't.

$$X \sim \text{Bernoulli}(0.3) \rightarrow$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

Find the probability that the first person he encounters will be able to speak English.

X – random variable: $X = 1$ – person speaks English, $X = 0$ – doesn't.

$$X \sim \text{Bernoulli}(0.3) \rightarrow P(X = 1) = p = 0.3$$

BINOMIAL DISTRIBUTION

- A random variable Y is said to follow Binomial distribution with parameters n and p if its PMF is given by

$$P_Y(k) = P(Y = k) = \begin{cases} C(n, k) \cdot p^k (1 - p)^{n-k}, & k = 0, 1, \dots, n \\ 0, & \text{otherwise} \end{cases}$$

- Represents the number of successes in a series of n independent Bernoulli trials, each of which results in success with probability p .

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(n = \quad p = \quad)$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

$$P(Y \geq 1) =$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

$$P(Y \geq 1) = P(Y = 1) + P(Y = 2) + P(Y = 3) + P(Y = 4) =$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

$$\begin{aligned} P(Y \geq 1) &= P(Y = 1) + P(Y = 2) + P(Y = 3) + P(Y = 4) = \\ &= 1 - P(Y = 0) = \end{aligned}$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

$$\begin{aligned} P(Y \geq 1) &= P(Y = 1) + P(Y = 2) + P(Y = 3) + P(Y = 4) = \\ &= 1 - P(Y = 0) = 1 - C(4, 0) \cdot 0.3^0 \cdot 0.7^4 = \end{aligned}$$

A TOURIST

- An English-speaking tourist visits a country in which 30% of the population speaks English. He needs to ask someone directions.

The tourist sees four local people standing at a bus stop. Find the probability that at least one of them will be able to speak English.

Y – the number of people (out of 4) who speak English.

$Y \sim \text{Binomial}(4, 0.3)$

$$\begin{aligned} P(Y \geq 1) &= P(Y = 1) + P(Y = 2) + P(Y = 3) + P(Y = 4) = \\ &= 1 - P(Y = 0) = 1 - C(4, 0) \cdot 0.3^0 \cdot 0.7^4 = 1 - 0.7^4 \end{aligned}$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

$$Y \sim$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = \quad p = \quad)$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y – the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

$$P(Y = 0) =$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

$$P(Y = 0) = C(5, 0) \cdot 0.25^0 \cdot 0.75^5 =$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Find the probability that there will be no red-flowered plants in the produced five offspring.

Random variable Y — the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

$$P(Y = 0) = C(5, 0) \cdot 0.25^0 \cdot 0.75^5 = 0.75^5 \sim 0.237$$

EXPECTED VALUE

EXPECTED VALUE

- Consider the following random variable:

x	-1	0	1
$P(x)$	1/3	1/3	1/3

EXPECTED VALUE

- Consider the following random variable:

x	-1	0	1
$P(x)$	$1/3$	$1/3$	$1/3$

- Which value would you expect to get *on average*?

EXPECTED VALUE

- Consider the following random variable:

x	-1	0	1
$P(x)$	1/3	1/3	1/3

- Which value would you expect to get *on average*?
- And now?

x	-1	0	1
$P(x)$	0.1	0.1	0.8

EXPECTED VALUE

- Let X be a discrete random variable with a finite range $R_X = \{x_1, \dots, x_n\}$. The expected value of X is defined as:

$$EX = \sum_{x_k \in R_X} x_k \cdot P(X = x_k)$$

EXPECTED VALUE

- Let X be a discrete random variable with a finite range $R_X = \{x_1, \dots, x_n\}$. The expected value of X is defined as:

$$EX = \sum_{x_k \in R_X} x_k \cdot P(X = x_k)$$

- Different notations: $EX, E[X], \mu_X, \dots$

EXPECTED VALUE

• $EX =$

x	-1	0	1
$P(x)$	1/3	1/3	1/3

• $EY =$

x	-1	0	1
$P(x)$	0.1	0.1	0.8

EXPECTED VALUE

- $EX = -1 \cdot \frac{1}{3} + 0 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} = 0$

x	-1	0	1
$P(x)$	1/3	1/3	1/3

- $EY =$

x	-1	0	1
$P(x)$	0.1	0.1	0.8

EXPECTED VALUE

- $EX = -1 \cdot \frac{1}{3} + 0 \cdot \frac{1}{3} + 1 \cdot \frac{1}{3} = 0$

x	-1	0	1
$P(x)$	1/3	1/3	1/3

- $EY = -1 \cdot 0.1 + 0 \cdot 0.1 + 1 \cdot 0.8 = 0.7$

x	-1	0	1
$P(x)$	0.1	0.1	0.8

BERNOULLI

- Consider $X \sim \text{Bernoulli}(p)$. What is $EX = ?$

BERNOULLI

- Consider $X \sim \text{Bernoulli}(p)$. What is $EX = ?$

x	0	1
$P(X = x)$	$1 - p$	p

$$EX =$$

BERNOULLI

- Consider $X \sim \text{Bernoulli}(p)$. What is $EX = ?$

x	0	1
$P(X = x)$	$1 - p$	p

$$EX = 0 \cdot (1 - p) + 1 \cdot p = p$$

PROPERTIES OF EXPECTED VALUE

- Let X be a random variable with $EX = \mu_X$ and Y be a random variable such that $Y = aX + b$. Then

$$EY = E(aX + b) = aEX + b = a\mu_X + b$$

PROPERTIES OF EXPECTED VALUE

- Let X be a random variable with $EX = \mu_X$ and Y be a random variable such that $Y = aX + b$. Then

$$EY = E(aX + b) = aEX + b = a\mu_X + b$$

- For any set of random variables X_1, \dots, X_n

$$E(X_1 + \dots + X_n) = EX_1 + \dots + EX_n$$

EXAMPLE

- Let X be a random variable with $EX = 10$.
- Let $Y = 5X + 1$

$$E(Y) =$$

- Let $Z = 3X - 2$

$$E(Z) =$$

EXAMPLE

- Let X be a random variable with $EX = 10$.
- Let $Y = 5X + 1$

$$E(Y) = 5 \cdot 10 + 1 = 51$$

- Let $Z = 3X - 2$

$$E(Z) =$$

EXAMPLE

- Let X be a random variable with $EX = 10$.
- Let $Y = 5X + 1$

$$E(Y) = 5 \cdot 10 + 1 = 51$$

- Let $Z = 3X - 2$

$$E(Z) = 3 \cdot 10 - 2 = 28$$

EXAMPLE

- Let X be a random variable with $EX = 10$ and Y be a random variable with $EX = 100$.

What is the expected value of $5X - Y + 50$?

EXAMPLE

- Let X be a random variable with $EX = 10$ and Y be a random variable with $EX = 100$.

What is the expected value of $5X - Y + 50$?

$$E(5X - Y + 50) =$$

EXAMPLE

- Let X be a random variable with $EX = 10$ and Y be a random variable with $EY = 100$.

What is the expected value of $5X - Y + 50$?

$$E(5X - Y + 50) = 5EX - EY + 50 =$$

EXAMPLE

- Let X be a random variable with $EX = 10$ and Y be a random variable with $EY = 100$.

What is the expected value of $5X - Y + 50$?

$$E(5X - Y + 50) = 5EX - EY + 50 = 50 - 100 + 50 = 0$$

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.

$$X_i \sim \quad EX_i =$$

TOSSING A COIN

- You are tossing a coin n times. How many heads to you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i =$

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i = p$

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i = p$

Then the total number of tosses is

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i = p$

Then the total number of tosses is $X_1 + \dots + X_n$

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i = p$

Then the total number of tosses is $X_1 + \dots + X_n$

$$E(X_1 + \dots + X_n) =$$

TOSSING A COIN

- You are tossing a coin n times. How many heads do you expect to see on average, if the probability of Heads is p ?

Let $X_i, i = 1 \dots 100$ be the outcome of the i -th toss.
 $X_i \sim \text{Bernoulli}(p), \quad EX_i = p$

Then the total number of tosses is $X_1 + \dots + X_n$

$$E(X_1 + \dots + X_n) = EX_1 + \dots + EX_n = np$$

SO,...

- The expected value of a Bernoulli random variable with parameter p is p :

$$X \sim \text{Bernoulli}(p) \quad \rightarrow \quad EX = p$$

- The expected value of a Binomial random variable with parameters (n, p) is np :

$$Y \sim \text{Binomial}(n, p) \quad \rightarrow \quad EY = np$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

Let random variable Y denote the number of red-flowered offspring.

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

Let random variable Y denote the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

Let random variable Y denote the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

Let random variable Y denote the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

$$EY = np =$$

FLOWERS

- Cross-fertilizing a red and a white flower produces red flowers 25% of the time. You cross-fertilize five pairs of red and white flowers.

Of the five cross-fertilized offspring, how many red-flowered plants do you expect?

Let random variable Y denote the number of red-flowered offspring.

$$Y \sim \text{Binomial}(n = 5, p = 0.25)$$

$$EY = np = 5 \cdot 0.25 = 1.25$$

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX =$$

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}$$

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}, \quad EX^2 =$$

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}, \quad EX^2 =$$

- Let $Y = X^2$. Distribution of Y :

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}, \quad EX^2 =$$

- Let $Y = X^2$. Distribution of Y :

y	1
$P(y)$	1

EXPECTATION OF X^2

- Consider the following random variable X :

x	-1	1
$P(x)$	1/3	2/3

$$EX = -1 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \frac{1}{3}, \quad EX^2 = EY = 1$$

- Let $Y = X^2$. Distribution of Y :

y	1
$P(y)$	1

PROPERTIES OF EXPECTED VALUE 2

- Let X be a random variable with range $R_X = \{x_1, \dots, x_n\}$ and $Y = g(X)$. Then

$$EY = E(g(X)) = \sum_{x_k \in R_X} g(x_k) \cdot P(X = x_k)$$

VARIANCE

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \quad , \quad EY =$$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0, \quad EY =$$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0,$$

$$EY = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0, \quad EY = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$$

But X and Y are very different....

VARIANCE

- The **variance** of a random variable X with $EX = \mu_X$ is defined as

$$\text{Var}(X) = E(X - EX)^2 = (EX)^2 - E(X^2)$$

VARIANCE

- The **variance** of a random variable X with $EX = \mu_X$ is defined as

$$\text{Var}(X) = E(X - EX)^2 = (EX)^2 - E(X^2)$$

- “How often does X take values far from its mean?”

VARIANCE

$$\textit{Var}(X) \geq 0$$

VARIANCE

$$\textit{Var}(X) \geq 0$$

because by definition it's the expected value of
 $(X - \mu_X)^2 \geq 0$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0,$$

$$EY = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$$

$$\text{Var}(X) =$$

$$\text{Var}(Y) =$$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0, \quad EY = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$$

$$\text{Var}(X) = E(X - EX)^2 = EX^2 = \frac{2}{3} \cdot 10^4$$

$$\text{Var}(Y) =$$

VARIANCE

- Consider the following two random variables X and Y :

x	-100	0	100
$P(X = x)$	1/3	1/3	1/3

y	-1	1
$P(Y = y)$	1/2	1/2

$$EX = \frac{1}{3} \cdot (-100) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 100 = 0, \quad EY = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$$

$$\text{Var}(X) = E(X - EX)^2 = EX^2 = \frac{2}{3} \cdot 10^4$$

$$\text{Var}(Y) = E(Y - EY)^2 = E(Y^2) = 1$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX =$$

$$Var(X) =$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) =$$

$$Var(X) =$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{21}{6} = \frac{7}{2}$$

$$Var(X) =$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{21}{6} = \frac{7}{2}$$

$$Var(X) = (EX)^2 - E(X^2) =$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{21}{6} = \frac{7}{2}$$

$$Var(X) = (EX)^2 - E(X^2) =$$

$$EX^2 =$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{21}{6} = \frac{7}{2}$$

$$Var(X) = (EX)^2 - E(X^2) =$$

$$EX^2 = \frac{1}{6}(1 + 4 + 9 + 16 + 25 + 36) = \frac{91}{6}$$

EXAMPLE

- You roll a fair dice once, random variable X – the outcome.

$$EX = \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{21}{6} = \frac{7}{2}$$

$$Var(X) = (EX)^2 - E(X^2) = \frac{49}{4} - \frac{91}{6} \sim 2.92$$

$$EX^2 = \frac{1}{6}(1 + 4 + 9 + 16 + 25 + 36) = \frac{91}{6}$$

VARIANCE OF A LINEAR COMBINATION

$$\text{Var}(aX + b) =$$

VARIANCE OF A LINEAR COMBINATION

$$\begin{aligned} \text{Var}(aX + b) &= \\ &= E\left(aX + b - E(aX + b)\right)^2 = \end{aligned}$$

VARIANCE OF A LINEAR COMBINATION

$$\begin{aligned} \text{Var}(aX + b) &= \\ &= E\left(aX + b - E(aX + b)\right)^2 = \\ &= E(aX + b - aEX - b)^2 = E(aX - aEX)^2 = \\ &= a^2 E(X - EX)^2 = \end{aligned}$$

VARIANCE OF A LINEAR COMBINATION

$$\begin{aligned} \text{Var}(aX + b) &= \\ &= E\left(aX + b - E(aX + b)\right)^2 = \\ &= E(aX + b - aEX - b)^2 = E(aX - aEX)^2 = \\ &= a^2 E(X - EX)^2 = \\ &= a^2 \text{Var}(X) \end{aligned}$$

EXAMPLE

- Let X be a random variable with $\text{Var}(X) = 10$.
- Let $Y = 5X + 1$

$$\text{Var}(Y) =$$

- Let $Z = -3X - 2$

$$\text{Var}(Z) =$$

EXAMPLE

- Let X be a random variable with $\text{Var}(X) = 10$.
- Let $Y = 5X + 1$

$$\text{Var}(Y) = \text{Var}(5X + 1) =$$

- Let $Z = -3X - 2$

$$\text{Var}(Z) =$$

EXAMPLE

- Let X be a random variable with $\text{Var}(X) = 10$.
- Let $Y = 5X + 1$

$$\text{Var}(Y) = \text{Var}(5X + 1) = 25\text{Var}(X) = 250$$

- Let $Z = -3X - 2$

$$\text{Var}(Z) =$$

EXAMPLE

- Let X be a random variable with $\text{Var}(X) = 10$.
- Let $Y = 5X + 1$

$$\text{Var}(Y) = \text{Var}(5X + 1) = 25\text{Var}(X) = 250$$

- Let $Z = -3X - 2$

$$\text{Var}(Z) = \text{Var}(-3X - 2) =$$

EXAMPLE

- Let X be a random variable with $\text{Var}(X) = 10$.
- Let $Y = 5X + 1$

$$\text{Var}(Y) = \text{Var}(5X + 1) = 25\text{Var}(X) = 250$$

- Let $Z = -3X - 2$

$$\text{Var}(Z) = \text{Var}(-3X - 2) = 9\text{Var}(X) = 90$$

STANDARD DEVIATION

- What are the measurement units of EX ? Same as those of X .
- What are the measurement units of $\text{Var}(X)$? $[\text{units of } X]^2$
 - Difficult to interpret.
- Standard deviation:

$$\text{std}(X) = \sqrt{\text{Var}(X)}$$