

# **ELEMENTARY COMBINATORICS & PROBABILITY**

## Lecture 9

Conditional probability, Bayes' rule

# LAST TIME

- Conditional probability
- The law of total probability
- Bayes' rule
- (Python) Testing for a rare disease

# TODAY

- More examples
- Independent events
- (Maybe) modelling in Python

# WARM-UP

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. A single ball is selected at random.
- **What is the probability that it's red?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. A single ball is selected at random.
- **What is the probability that it's red?**

$$P(R) =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. A single ball is selected at random.
- **What is the probability that it's red?**

$$P(R) = \frac{2}{2 + 3} = \frac{2}{5} = 0.4$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- ... **they're both red?**
- .
- ... **they're of different colors?**



# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- ... **they're both red?**  
*You can consider it both ordered or unordered*  
.
- ... **they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- ... **they're both red?**  
*You can consider it both ordered or unordered*

$$P(RR) =$$

- ... **they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**  
*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} =$$

- **... they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} =$$

- **... they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} = \frac{1}{10}$$

- **... they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} = \frac{1}{10}$$

- **... they're of different colors?**

$$P(E) =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} = \frac{1}{10}$$

- **... they're of different colors?**

$$P(E) = \frac{C(2, 1) \cdot 2 \cdot 3}{5 \cdot 4} =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} = \frac{1}{10}$$

- **... they're of different colors?**

$$P(E) = \frac{C(2, 1) \cdot 2 \cdot 3}{5 \cdot 4} = \frac{C(2, 1) \cdot C(3, 1)}{C(5, 2)} =$$



# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected at random. What is the probability that...
- **... they're both red?**

*You can consider it both ordered or unordered*

$$P(RR) = \frac{2 \cdot 1}{5 \cdot 4} = \frac{C(2, 2)}{C(5, 2)} = \frac{1}{10}$$

- **... they're of different colors?**

$$P(E) = \frac{C(2, 1) \cdot 2 \cdot 3}{5 \cdot 4} = \frac{C(2, 1) \cdot C(3, 1)}{C(5, 2)} = \frac{3}{5}$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...
- ... they're both red?
- ... they're of different colors?

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...
- ... **they're both red?**

$$P(RR) =$$

- ... **they're of different colors?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...

- ... they're both red?

$$P(RR) = \frac{2 \cdot 2}{5 \cdot 5} =$$

- ... they're of different colors?

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...

- ... they're both red?

$$P(RR) = \frac{2 \cdot 2}{5 \cdot 5} = \frac{4}{25}$$

- ... they're of different colors?

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...

- ... they're both red?

$$P(RR) = \frac{2 \cdot 2}{5 \cdot 5} = \frac{4}{25}$$

- ... they're of different colors?

$$P(E) =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...

- ... they're both red?

$$P(RR) = \frac{2 \cdot 2}{5 \cdot 5} = \frac{4}{25}$$

- ... they're of different colors?

$$P(E) = \frac{C(2,1) \cdot 2 \cdot 3}{5 \cdot 5} =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Two balls are selected one by one. The first ball is returned to the box before the second one is selected. What is the probability that...
- ... **they're both red?**

$$P(RR) = \frac{2 \cdot 2}{5 \cdot 5} = \frac{4}{25}$$

- ... **they're of different colors?**

$$P(E) = \frac{C(2,1) \cdot 2 \cdot 3}{5 \cdot 5} = \frac{12}{25}$$



# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... all of them are red?
- ... 2 of them are red and the other one is blue?

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **all of them are red?**

$$P(RRR) =$$

- ... **2 of them are red and the other one is blue?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **all of them are red?**

$$P(RRR) = \frac{2 \cdot 2 \cdot 2}{5^3} =$$

- ... **2 of them are red and the other one is blue?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- **... all of them are red?**

$$P(RRR) = \frac{2 \cdot 2 \cdot 2}{5^3} = \frac{8}{125}$$

- **... 2 of them are red and the other one is blue?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... all of them are red?

$$P(RRR) = \frac{2 \cdot 2 \cdot 2}{5^3} = \frac{8}{125}$$

- ... 2 of them are red and the other one is blue?

$$P(E) =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- **... all of them are red?**

$$P(RRR) = \frac{2 \cdot 2 \cdot 2}{5^3} = \frac{8}{125}$$

- **... 2 of them are red and the other one is blue?**

$$P(E) = \frac{C(3,1) \cdot 3 \cdot 2^2}{5^3} =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Three balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- **... all of them are red?**

$$P(RRR) = \frac{2 \cdot 2 \cdot 2}{5^3} = \frac{8}{125}$$

- **... 2 of them are red and the other one is blue?**

$$P(E) = \frac{C(3,1) \cdot 3 \cdot 2^2}{5^3} = \frac{36}{125}$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Four balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **1 ball is red and 3 balls are blue?**
- ... **2 balls are red and 2 balls are blue?**



# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Four balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **1 ball is red and 3 balls are blue?**

$$P(E) =$$

- ... **2 balls are red and 2 balls are blue?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Four balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **1 ball is red and 3 balls are blue?**

$$P(E) = \frac{C(4,1) \cdot 2 \cdot 3^3}{5^4}$$

- ... **2 balls are red and 2 balls are blue?**

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Four balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **1 ball is red and 3 balls are blue?**

$$P(E) = \frac{C(4,1) \cdot 2 \cdot 3^3}{5^4}$$

- ... **2 balls are red and 2 balls are blue?**

$$P(E) =$$

# BALLS IN A BOWL

- There're 2 red balls and 3 blue balls in a box. Four balls are selected one by one. Each ball is returned to the box before the next one is selected. What is the probability that...
- ... **1 ball is red and 3 balls are blue?**

$$P(E) = \frac{C(4,1) \cdot 2 \cdot 3^3}{5^4}$$

- ... **2 balls are red and 2 balls are blue?**

$$P(E) = \frac{C(4,2) \cdot 2^2 \cdot 3^2}{5^4}$$

# PROBLEM SET 7

Conditional probability, the law of total probability

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} =$$

$$P(3H \text{ and } H **) = \quad , \quad P(H **) =$$



# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} =$$

$$P(3H \text{ and } H **) = \frac{1}{2^3}, \quad P(H **) =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} =$$

$$P(3H \text{ and } H **) = \frac{1}{2^3}, \quad P(H **) = \frac{2^2}{2^3} =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} =$$

$$P(3H \text{ and } H **) = \frac{1}{2^3}, \quad P(H **) = \frac{2^2}{2^3} = \frac{1}{2}$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first flip came up heads?

$$P(3H|H **) = \frac{P(3H \text{ and } H **)}{P(H **)} = \frac{2}{2^3} = \frac{1}{4}$$

$$P(3H \text{ and } H **) = \frac{1}{2^3}, \quad P(H **) = \frac{2^2}{2^3} = \frac{1}{2}$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} =$$

$$P(3H \text{ and } HH *) = \quad , \quad P(HH *) =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} =$$

$$P(3H \text{ and } HH *) = \frac{1}{2^3}, \quad P(HH *) =$$



# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} =$$

$$P(3H \text{ and } HH *) = \frac{1}{2^3}, \quad P(HH *) = \frac{2}{2^3} =$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} =$$

$$P(3H \text{ and } HH *) = \frac{1}{2^3}, \quad P(HH *) = \frac{2}{2^3} = \frac{1}{4}$$

# COINS

- A coin is flipped three times. What is the probability of getting three heads given that the first two flips resulted in two heads?

$$P(3H|HH *) = \frac{P(3H \text{ and } HH *)}{P(HH *)} = \frac{4}{2^3} = \frac{1}{2}$$

$$P(3H \text{ and } HH *) = \frac{1}{2^3}, \quad P(HH *) = \frac{2}{2^3} = \frac{1}{4}$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T ***) =$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T *** ) = \frac{P(2H \text{ and } T *** )}{P(T *** )} =$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T *** ) = \frac{P(2H \text{ and } T *** )}{P(T *** )} =$$

$$P(2H \text{ and } T *** ) = \quad , \quad P(T *** ) =$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T *** ) = \frac{P(2H \text{ and } T *** )}{P(T *** )} =$$

$$P(2H \text{ and } T *** ) = \frac{3}{2^4}, \quad P(T *** ) =$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T *** ) = \frac{P(2H \text{ and } T *** )}{P(T *** )} =$$

$$P(2H \text{ and } T *** ) = \frac{3}{2^4}, \quad P(T *** ) = \frac{2^3}{2^4} =$$



# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T *** ) = \frac{P(2H \text{ and } T *** )}{P(T *** )} =$$

$$P(2H \text{ and } T *** ) = \frac{3}{2^4}, \quad P(T *** ) = \frac{2^3}{2^4} = \frac{1}{2}$$

# COINS

- A coin is flipped 4 times. What is the probability that heads occurs twice given that the first flip is tails?

$$P(2H|T ***) = \frac{P(2H \text{ and } T ***)}{P(T ***)} = \frac{3}{8}$$

$$P(2H \text{ and } T ***) = \frac{3}{2^4}, \quad P(T ***) = \frac{2^3}{2^4} = \frac{1}{2}$$

# THE LAW OF TOTAL PROBABILITY

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$P(P) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$P(P) = P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) =$$



# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \\ &= \frac{1}{3} \cdot 0.21 + 0.1 \cdot (1 - 0.21) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \\ &= \frac{1}{3} \cdot 0.21 + 0.1 \cdot (1 - 0.21) = 0.07 + 0.079 = 0.149 \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$

# BAYES' RULE

# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

$$P(D) = 0.01$$

# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

$$P(D) = 0.01$$

- A 90% - accurate test has been developed to detect the disease

# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

$$P(D) = 0.01$$

- A 90% - accurate test has been developed to detect the disease

$$P(+|D) = 0.9, \quad P(-|no\ D) = 0.9$$



# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

$$P(D) = 0.01$$

- A 90% - accurate test has been developed to detect the disease

$$P(+|D) = 0.9, \quad P(-|no\ D) = 0.9$$

- You test positive. How much should you be worried?

# A TEST FOR DISEASE

- Imagine that a rare disease affects 1% of the population

$$P(D) = 0.01$$

- A 90% - accurate test has been developed to detect the disease

$$P(+|D) = 0.9, \quad P(-|no\ D) = 0.9$$

- You test positive. How much should you be worried?

$$P(D \mid +) = ?$$

# HOW RELIABLE THE TEST IS?

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9$$

REALITY / TEST RESULTS	POSITIVE	NEGATIVE
<b>HEALTHY</b> $\approx 92\%$	<b>FALSE POSITIVES</b> 10% of 99% healthy people = 9.9% of the population	<b>TRUE NEGATIVES</b> 90% of 99% healthy people = 89.1% of the population
<b>ILL</b> $\approx 8\%$	<b>TRUE POSITIVES</b> 90% of 1% ill people = 0.9% of the population	<b>FALSE NEGATIVES</b> 10% of 1% ill people = 0.1% of the population

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ \mid D) = P(- \mid \text{no } D) = 0.9, \quad P(D \mid +) = ?$$

$$P(D \mid +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) =$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) =$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) = 0.009$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) = 0.009$$

$$P(+) =$$



# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) = 0.009$$

$$P(+ ) = P(+ | D) \cdot P(D) + P(+ | \text{no } D) \cdot P(\text{no } D) =$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} =$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) = 0.009$$

$$\begin{aligned} P(+) &= P(+ | D) \cdot P(D) + P(+ | \text{no } D) \cdot P(\text{no } D) = \\ &= 0.9 \cdot 0.01 + 0.1 \cdot 0.99 = 0.108 \end{aligned}$$

# SAME THING, BUT IN TERMS OF PROBABILITY

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9, \quad P(D | +) = ?$$

$$P(D | +) = \frac{P(+ \text{ and } D)}{P(+)} = \frac{\mathbf{0.009}}{\mathbf{0.108}} \approx \mathbf{0.083}$$

$$P(+ | D) = \frac{P(+ \text{ and } D)}{P(D)} \rightarrow P(+ \text{ and } D) = P(+ | D) \cdot P(D) = 0.009$$

$$\begin{aligned} P(+) &= P(+ | D) \cdot P(D) + P(+ | \text{no } D) \cdot P(\text{no } D) = \\ &= 0.9 \cdot 0.01 + 0.1 \cdot 0.99 = 0.108 \end{aligned}$$

# HOW RELIABLE THE TEST IS?

$$P(D) = 0.01, \quad P(+ | D) = P(- | \text{no } D) = 0.9$$

REALITY / TEST RESULTS	POSITIVE	NEGATIVE
<b>HEALTHY</b> $\approx 92\%$	<b>FALSE POSITIVES</b> 10% of 99% healthy people = 9.9% of the population	<b>TRUE NEGATIVES</b> 90% of 99% healthy people = 89.1% of the population
<b>ILL</b> $\approx 8\%$	<b>TRUE POSITIVES</b> 90% of 1% ill people = 0.9% of the population	<b>FALSE NEGATIVES</b> 10% of 1% ill people = 0.1% of the population

# BAYES' RULE

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{P(B|A)}{P(B)} \cdot P(A)$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$P(A|blue) =$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$P(A|blue) = \frac{P(blue|A) \cdot P(A)}{P(blue)} =$$



# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$\begin{aligned} P(A|blue) &= \frac{P(blue|A) \cdot P(A)}{P(blue)} = \\ &= \frac{P(blue|A) \cdot P(A)}{P(blue|A) \cdot P(A) + P(blue|B) \cdot P(B)} = \end{aligned}$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$\begin{aligned} P(A|blue) &= \frac{P(blue|A) \cdot P(A)}{P(blue)} = \\ &= \frac{P(blue|A) \cdot P(A)}{P(blue|A) \cdot P(A) + P(blue|B) \cdot P(B)} = \frac{0.4 \cdot 0.5}{0.4 \cdot 0.5 + 0.2 \cdot 0.5} = \\ &= \frac{0.2}{0.3} = \frac{2}{3} \end{aligned}$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$  — a coin with 2 H is selected       $P(E_H) =$

$E_T$  — a coin with 2 T is selected       $P(E_T) =$

$E_F$  — a fair coin is selected       $P(E_F) =$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$$E_H \text{ — a coin with 2 H is selected} \qquad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \qquad P(E_T) =$$

$$E_F \text{ — a fair coin is selected} \qquad P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) =$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9},$$



# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) = \frac{P(H|E_F) \cdot P(E_F)}{P(H)} =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$$\begin{array}{ll} E_H - \text{a coin with 2 H is selected} & P(E_H) = \frac{3}{3+4+2} = \frac{1}{3} \\ E_T - \text{a coin with 2 T is selected} & P(E_T) = \frac{4}{3+4+2} = \frac{4}{9} \\ E_F - \text{a fair coin is selected} & P(E_F) = \frac{2}{3+4+2} = \frac{2}{9} \end{array}$$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) = \frac{P(H|E_F) \cdot P(E_F)}{P(H)} = \frac{\frac{1}{2} \cdot \frac{2}{9}}{\frac{4}{9}} = \frac{1}{4}$$

# EXAM

- Exam is divided into two parts. The probability of getting B or better on Part 1 is 0.8, the probability of getting B or better on Part 2 is 0.6, and the probability of getting B or better on both parts is 0.5. What is the probability of getting B or better on Part 2 if you got B or better on Part 1?

# EXAM

- Exam is divided into two parts. The probability of getting B or better on Part 1 is 0.8, the probability of getting B or better on Part 2 is 0.6, and the probability of getting B or better on both parts is 0.5. What is the probability of getting B or better on Part 2 if you got B or better on Part 1?

$$P(E_2|E_1) =$$

# EXAM

- Exam is divided into two parts. The probability of getting B or better on Part 1 is 0.8, the probability of getting B or better on Part 2 is 0.6, and the probability of getting B or better on both parts is 0.5. What is the probability of getting B or better on Part 2 if you got B or better on Part 1?

$$P(E_2|E_1) = \frac{P(E_1 \& E_2)}{P(E_1)} =$$

# EXAM

- Exam is divided into two parts. The probability of getting B or better on Part 1 is 0.8, the probability of getting B or better on Part 2 is 0.6, and the probability of getting B or better on both parts is 0.5. What is the probability of getting B or better on Part 2 if you got B or better on Part 1?

$$P(E_2|E_1) = \frac{P(E_1 \& E_2)}{P(E_1)} = \frac{0.5}{0.8} = \frac{5}{8}$$

# A MAGICIAN

- A magician has five coins in his pocket. Four of the coins are fair but the other is a biased coin and has two heads. He pulls a coin at random from his pocket and flips it five times. It comes up heads all five times. What is the probability that he selected a fair coin from his pocket?



# A MAGICIAN

- A magician has five coins in his pocket. Four of the coins are fair but the other is a biased coin and has two heads. He pulls a coin at random from his pocket and flips it five times. It comes up heads all five times. What is the probability that he selected a fair coin from his pocket?

$$P(F|5H) =$$

# A MAGICIAN

- A magician has five coins in his pocket. Four of the coins are fair but the other is a biased coin and has two heads. He pulls a coin at random from his pocket and flips it five times. It comes up heads all five times. What is the probability that he selected a fair coin from his pocket?

$$P(F|5H) = \frac{P(5H|F)P(F)}{P(5H)} =$$

# A MAGICIAN

- A magician has five coins in his pocket. Four of the coins are fair but the other is a biased coin and has two heads. He pulls a coin at random from his pocket and flips it five times. It comes up heads all five times. What is the probability that he selected a fair coin from his pocket?

$$P(F|5H) = \frac{P(5H|F)P(F)}{P(5H)} = \frac{P(5H|F)P(F)}{P(5H|F)P(F) + P(5H|B)P(B)} =$$

# A MAGICIAN

- A magician has five coins in his pocket. Four of the coins are fair but the other is a biased coin and has two heads. He pulls a coin at random from his pocket and flips it five times. It comes up heads all five times. What is the probability that he selected a fair coin from his pocket?

$$\begin{aligned} P(F|5H) &= \frac{P(5H|F)P(F)}{P(5H)} = \frac{P(5H|F)P(F)}{P(5H|F)P(F) + P(5H|B)P(B)} = \\ &= \frac{0.5^5 \cdot 0.8}{0.5^5 \cdot 0.8 + 1 \cdot 0.2} \end{aligned}$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

$$P(A) = \quad , \quad P(B) = \quad , \quad P(F|A) = \quad , \quad P(F|B) =$$



# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

$$P(A) = 0.7, \quad P(B) = \quad, \quad P(F|A) = \quad, \quad P(F|B) =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

$$P(A) = 0.7, \quad P(B) = 0.3, \quad P(F|A) = \quad , \quad P(F|B) =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

$$P(A) = 0.7, \quad P(B) = 0.3, \quad P(F|A) = 0.2, \quad P(F|B) =$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$P(A|F) = \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} =$$

$$P(A) = 0.7, \quad P(B) = 0.3, \quad P(F|A) = 0.2, \quad P(F|B) = 0.1$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$\begin{aligned} P(A|F) &= \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} = \\ &= \frac{0.2 \cdot 0.7}{0.2 \cdot 0.7 + 0.1 \cdot 0.3} = \end{aligned}$$

$$P(A) = 0.7, \quad P(B) = 0.3, \quad P(F|A) = 0.2, \quad P(F|B) = 0.1$$

# TEACHING

- Methods A and B are available for teaching a skill. The failure rate is 20% for A and 10% for B. Method B is used only 30% of the time, and A is used the other 70%. A worker was taught the skill by one of the methods but failed to learn it. What is the probability that she was taught by method A?

$$\begin{aligned} P(A|F) &= \frac{P(F|A) \cdot P(A)}{P(F)} = \frac{P(F|A) \cdot P(A)}{P(F|A) \cdot P(A) + P(F|B) \cdot P(B)} = \\ &= \frac{0.2 \cdot 0.7}{0.2 \cdot 0.7 + 0.1 \cdot 0.3} = \frac{14}{17} \approx 0.82 \end{aligned}$$

$$P(A) = 0.7, \quad P(B) = 0.3, \quad P(F|A) = 0.2, \quad P(F|B) = 0.1$$

# INDEPENDENT EVENTS

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) =$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**



# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} =$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

$$P(2H|H ***) =$$

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

$$P(2H|H ***) = \frac{P(2H \text{ and } H ***)}{P(H ***)} =$$

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

$$P(2H|H ***) = \frac{P(2H \text{ and } H ***)}{P(H ***)} = \frac{3/2^4}{2^3/2^4} = \frac{3}{8}$$

# COINS

- When a coin is flipped four times, what is the probability that ...
- ... **heads comes up exactly twice?**

$$P(2H) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

- ... **heads comes up exactly twice given that heads comes up on the first flip?**

$$P(2H|H ***) = \frac{P(2H \text{ and } H ***)}{P(H ***)} = \frac{3/2^4}{2^3/2^4} = \frac{3}{8}$$

# INDEPENDENT EVENTS

- Two events  $A$  and  $B$  from the same sample space  $S$  are independent if

$$P(AB) = P(A) \cdot P(B)$$

# INDEPENDENT EVENTS

- Two events  $A$  and  $B$  from the same sample space  $S$  are independent if

$$P(AB) = P(A) \cdot P(B)$$

- Note that this means that

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{P(A) \cdot P(B)}{P(B)} = P(A)$$



# INDEPENDENT EVENTS

- Two events  $A$  and  $B$  from the same sample space  $S$  are independent if

$$P(AB) = P(A) \cdot P(B)$$

- Note that this means that

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{P(A) \cdot P(B)}{P(B)} = P(A)$$

$$P(B|A) = \frac{P(AB)}{P(A)} = \frac{P(A) \cdot P(B)}{P(A)} = P(B)$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6},$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6},$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) =$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

$$P(EF) = P(E) \cdot P(F) = \frac{1}{36}$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

$$P(EF) = P(E) \cdot P(F) = \frac{1}{36} \rightarrow E \& F \text{ are independent!}$$



# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) =$$

$$P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) =$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4}$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4} \neq P(E) \cdot P(F) = \frac{5}{2^5}$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4} \neq P(E) \cdot P(F) = \frac{5}{2^5}$$

$\rightarrow E$  &  $F$  aren't independent

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) =$$

$$P(F) =$$

$$P(EF) =$$



# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) =$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) = \frac{3}{2^3} = P(E) \cdot P(F)$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) = \frac{3}{2^3} = P(E) \cdot P(F)$$

→  $E$  &  $F$  are independent!

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) =$$



# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) = P(Sat) \cdot P(Sun) =$$

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) = P(Sat) \cdot P(Sun) = \frac{1}{4}$$

# MONTY HALL

# MONTY HALL

