

ELEMENTARY COMBINATORICS & PROBABILITY

Probability of an event

LAST TIME

- Random experiments
- Sample space
- Events
- Probability of an event

TODAY

- Problem set 6
- More pen-and-paper exercises
- Python exercises

WARM-UP

PROBABILITY OF AN EVENT

- Consider a random experiment where all outcomes are equally likely.
- Probability $P(E)$ of an event E is such an experiment is

$$P(E) = \frac{\# \text{ ways } E \text{ can happen}}{\# \text{ possible outcomes}}$$

- Let E^C be the complementary event. Then


$$P(E^C) = 1 - P(E)$$

- Let F be some other event. Then

$$P(E \cup F) = P(E) + P(F) - P(E \cap F)$$


ROULETTE

- American roulette: 38 sectors
 - 18 red
 - 18 black
 - 2 green (0 and 00)




ROULETTE

- American roulette: 38 sectors
 - 18 red
 - 18 black
 - 2 green (0 and 00)
- What is the probability to win if you
 - Bet on red
 - Bet on black
 - Bet on green




ROULETTE

- American roulette: 38 sectors
 - 18 red
 - 18 black
 - 2 green (0 and 00)
- What is the probability to win if you
 - Bet on red
$$P(\text{win}) = 18/38 \approx 0.474$$
 - Bet on black
 - Bet on green




ROULETTE

- American roulette: 38 sectors
 - 18 red
 - 18 black
 - 2 green (0 and 00)
- What is the probability to win if you
 - Bet on red
$$P(\text{win}) = 18/38 \approx 0.474$$
 - Bet on black
$$P(\text{win}) = 18/38 \approx 0.474$$
 - Bet on green



ROULETTE

- American roulette: 38 sectors
 - 18 red
 - 18 black
 - 2 green (0 and 00)
- What is the probability to win if you
 - Bet on red
$$P(\text{win}) = 18/38 \approx 0.474$$
 - Bet on black
$$P(\text{win}) = 18/38 \approx 0.474$$
 - Bet on green
$$P(\text{win}) = 2/38 \approx 0.05$$



PROBLEM SET 6

Selected problems

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$|E| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$|E| = |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = 15 + 10 + 6 = 31 \\ &\text{ways to choose 2 same-color balls} \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = 15 + 10 + 6 = 31 \\ &\text{ways to choose 2 same-color balls} \end{aligned}$$

$$P(E) =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = 15 + 10 + 6 = 31 \\ &\text{ways to choose 2 same-color balls} \end{aligned}$$

$$P(E) = 31/105$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$$|S| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$|E| =$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$$|E| = |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$$\begin{aligned} |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 5 + 5 \cdot 4 + 4 \cdot 3 = \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$$\begin{aligned} |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 5 + 5 \cdot 4 + 4 \cdot 3 = 30 + 20 + 12 = 62 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

$$P(E) =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$$\begin{aligned} |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 5 + 5 \cdot 4 + 4 \cdot 3 = 30 + 20 + 12 = 62 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

$$P(E) = 62/210 = 31/105$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$|S| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$|S| = 15 \cdot 15 = 225$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$|S| = 15 \cdot 15 = 225$$
$$|E| = |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$\begin{aligned} |S| &= 15 \cdot 15 = 225 \\ |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 6 + 5 \cdot 5 + 4 \cdot 4 = 36 + 25 + 16 = 77 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$\begin{aligned} |S| &= 15 \cdot 15 = 225 \\ |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 6 + 5 \cdot 5 + 4 \cdot 4 = 36 + 25 + 16 = 77 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

$$P(E) =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color if the first ball is returned to the bowl before the second ball is selected?**

Grabbing 2 balls ne-by-one (with replacement):

$$\begin{aligned} |S| &= 15 \cdot 15 = 225 \\ |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 6 + 5 \cdot 5 + 4 \cdot 4 = 36 + 25 + 16 = 77 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

$$P(E) = 77/225$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$$|S| =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$$|E| =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

$$P(E) =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

$$P(E) = \frac{2}{36} = \frac{1}{18} \approx 0.056$$

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.
What is the probability of winning?

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls:

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball:

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations:

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations: $C(53, 5) \cdot 42 = 120,526,770$

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations: $C(53, 5) \cdot 42 = 120,526,770$

Number of winning combinations:

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations: $C(53, 5) \cdot 42 = 120,526,770$

Number of winning combinations: 1

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations: $C(53, 5) \cdot 42 = 120,526,770$

Number of winning combinations: 1

Probability of winning:

LOTTERY

- In a lottery, 5 balls are selected at random from a collection of 53 numbered white balls, and one ball is selected at random from a collection of 42 numbered red balls. A lottery ticket lists the numbers of 5 white balls (in any order) as well as the number of the red ball.

What is the probability of winning?

Ways of choosing 5 white balls: $C(53, 5)$

Ways of choosing 1 red ball: $C(42, 1) = 42$

Total number of combinations: $C(53, 5) \cdot 42 = 120,526,770$

Number of winning combinations: 1

Probability of winning: $1/[C(53, 5) \cdot 42] \approx 0.0000000083$

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| =$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| =$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| =$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) =$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- **... in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- **... in a 4-child family, there are 2 boys and 2 girls**

$$|S| =$$

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- **... in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- **... in a 4-child family, there are 2 boys and 2 girls**

$$|S| = 2^4$$

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- ... **in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- ... **in a 4-child family, there are 2 boys and 2 girls**

$$|S| = 2^4$$

$$|E| = C(4, 2) = 6 \text{ different families with 2 boys and 2 girls}$$

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- **... in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- **... in a 4-child family, there are 2 boys and 2 girls**

$$|S| = 2^4$$

$$|E| = C(4, 2) = 6 \text{ different families with 2 boys and 2 girls}$$

$$P(E) =$$

BOYS AND GIRLS

- Assuming equal outcomes, calculate the probability that ...
- **... in a 3-child family, there are 2 boys and 1 girl**

$$|S| = 2 \cdot 2 \cdot 2 = 8 \text{ possible}$$

$$|E| = C(3, 2) \cdot 1 \cdot 1 \cdot 1 = 3 \text{ different families with 2 boys and 1 girl}$$

$$P(E) = 3/8$$

- **... in a 4-child family, there are 2 boys and 2 girls**

$$|S| = 2^4$$

$$|E| = C(4, 2) = 6 \text{ different families with 2 boys and 2 girls}$$

$$P(E) = 3/8$$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- ... **it doesn't contain digit 7?**
- ... **it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- ... **it doesn't contain digit 7?**
 $|S| =$
- ... **it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| =$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| =$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| =$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| =$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits
 $|P| =$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits
 $|P| = 91/101 \approx 0.9$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- ... **all girls are standing next to each other?**
- ... **all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- ... **all girls are standing next to each other?**
 $|S| =$
- ... **all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| =$
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) =$$

- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| =$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| = 10!$ possible permutations of 10 kids

$|E| =$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| = 10!$ possible permutations of 10 kids

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together
$$P(E) =$$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together
$$P(E) = \frac{4! \cdot 6! \cdot 2!}{10!} \approx 0.0096$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$P(E_3 \cup E_7) =$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$P(E_3 \cup E_7) = P(E_3) + P(E_7) - P(E_3 \cap E_7) =$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$\begin{aligned} P(E_3 \cup E_7) &= P(E_3) + P(E_7) - P(E_3 \cap E_7) = \\ &= \frac{33}{100} + \frac{14}{100} - \frac{4}{100} = \frac{43}{100} = 0.43 \end{aligned}$$

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

ways to distribute candies

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

ways to distribute candies so that each kid gets at least one.

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

$P(E) =$

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

$$P(E) = \frac{C(14, 6)}{C(21, 6)} = \frac{3003}{54264} = 0.055$$

PROBABILITY AS LIMITING FREQUENCY

MORE EXAMPLES

PLAYING CARDS

- 52 cards in a deck
- 4 suits (types):
 - clubs 
 - diamonds 
 - hearts 
 - spades 
- 13 cards of each type

	2	3	4	5	6	7	8	9	10	J	Q	K	A
♥													
♦		3♦											
♣										J♣			
♠													

POKER

- In poker, each player is dealt (supplied with) 5 of the 52 cards, called a **hand**. How many different hands are there?

POKER

- In poker, each player is dealt (supplied with) 5 of the 52 cards, called a **hand**. How many different hands are there?

$$C(52, 5) = \frac{52!}{5! \cdot 47!} = 2,598,960$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

$$P(E) =$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3 ♥, 5 ♦, 6 ♠, 10 ♣ and Q ♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

$$P(E) = \frac{1}{2,598,960} \approx 0.000000385$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

$$P(E) =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

$$P(E) = \frac{5148}{2,598,960} \approx 0.00198$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

$$P(E) =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

$$P(E) = \frac{123552}{2,598,960} \approx 0.048$$