

ELEMENTARY COMBINATORICS & PROBABILITY

Lecture 1

Sets
Basic counting

TODAY

- Introduction
 - about this course;
 - about me.
- What is combinatorics (and why it's worth studying).
- Basic counting principles.
- Examples.


ABOUT ME

- EVGENIYA Korneva



evgeniakorneva@gmail.com

ABOUT ME

- EVGENIYA Korneva  evgeniakorneva@gmail.com
- Education:
 - 2015 - Bachelor of Applied Mathematics
(Moscow, Russia)
 - 2016 - Master of Artificial Intelligence
(Leuven, Belgium)



NATIONAL RESEARCH
UNIVERSITY

KU LEUVEN

ABOUT ME

- EVGENIYA Korneva  evgeniakorneva@gmail.com
- Education:
 - 2015 - Bachelor of Applied Mathematics
(Moscow, Russia)
 - 2016 - Master of Artificial Intelligence
(Leuven, Belgium)
 - 2016 - ... Doctoral researcher at KU Leuven



NATIONAL RESEARCH
UNIVERSITY



ABOUT ME

• EVGENIYA Korneva  evgeniakorneva@gmail.com

• Education:

• 2015 - Bachelor of Applied Mathematics
(Moscow, Russia)

• 2016 - Master of Artificial Intelligence
(Leuven, Belgium)

• 2016 - ... Doctoral researcher at KU Leuven



📍 Leuven, Belgium

LOGISTICS

- Class: 09:00 – 12:20

LOGISTICS

- Class: 09:00 – 12:20
- Week 1: Combinatorics.
- Weeks 2 & 3: Probability.

LOGISTICS

- Class: 09:00 – 12:20
- Week 1: Combinatorics.
- Weeks 2 & 3: Probability.
- Evaluation:
 - 5 graded assignments
 - 2 interim exams (March 22 & 29)
 - Final exam (April 2)

LOGISTICS

- Class: 09:00 – 12:20
- Week 1: Combinatorics.
- Weeks 2 & 3: Probability.
- Evaluation:
 - 5 graded assignments *5% each*;
 - 2 interim exams (March 22 & 29) *25% each*;
 - Final exam (April 2), *25%*.

LET'S START!

WHAT IS COMBINATORICS

- Fine art of counting.

WHAT IS COMBINATORICS




WHAT IS COMBINATORICS

- We can easily count by enumeration:

WHAT IS COMBINATORICS

- We can easily count by enumeration:




WHAT IS COMBINATORICS

- We can easily count by enumeration:



WHAT IS COMBINATORICS


- We can easily count by enumeration:



- But enumeration is not always easy.

MOTIVATING EXAMPLE: LICENSE PLATES

MOTIVATING EXAMPLE: LICENSE PLATES




- A Russian license plate:
3 letters + 3 digits + regional code

MOTIVATING EXAMPLE: LICENSE PLATES



- A Russian license plate:
3 letters + 3 digits + regional code
- 12 letters and 10 digits can be used.

MOTIVATING EXAMPLE: LICENSE PLATES




- A Russian license plate:
3 letters + 3 digits + regional code
- 12 letters and 10 digits can be used.
- How many combinations are there per region?

MOTIVATING EXAMPLE: LICENSE PLATES



- A Russian license plate:
 - 3 letters + 3 digits + regional code
- 12 letters and 10 digits can be used.
- How many combinations are there per region?
 - ~ 1.7 million unique plates

MOTIVATING EXAMPLE: LICENSE PLATES



- A Russian license plate:
3 letters + 3 digits + regional code
- 12 letters and 10 digits can be used.
- How many combinations are there per region?
 - ~ 1.7 million unique plates
- Moscow has **nine** regional codes.
 - ~ 1.7 million \rightarrow ~15.5 million unique plates.

MOTIVATING EXAMPLE: PHONE NUMBERS

MOTIVATING EXAMPLE: PHONE NUMBERS

Japan is running out of phone numbers, so it's making longer ones

Its current 11-digit numbers could run out by 2022

By Jon Porter | @JonPorty | May 16, 2019, 8:45am EDT

f t SHARE



Photo by Amelia Holowaty Krales / The Verge

Source: [The Verge](#)

Source: [Independent](#)



Local calls using area codes seen as a better alternative to adding another digit

WHAT IS COMBINATORICS

- We can easily count by enumeration:



- But enumeration is not always easy.

**Combinatorics = fine art of counting
(without *actually* enumerating).**

COMBINATORICS: WHAT ELSE FOR?

COMBINATORICS: WHAT ELSE FOR?

- Computer science: determining the time and storage required to solve a problem.

COMBINATORICS: WHAT ELSE FOR?

- Computer science: determining the time and storage required to solve a problem.
- Basis of probability theory
 - *we'll see it later in the course.*

COMBINATORICS: WHAT ELSE FOR?

- Computer science: determining the time and storage required to solve a problem.
- Basis of probability theory
 - *we'll see it later in the course.*
- Some proof techniques in mathematics rely on combinatorics.

SET THEORY

SETS

- We almost always have a collection of things we are trying to study.

SETS

- We almost always have a collection of things we are trying to study.
- **Set** is a collection of *distinct* objects in which the *order doesn't matter*.

SETS

- Set is a collection of *distinct* objects in which the *order doesn't matter*

SETS

- Set is a collection of distinct objects in which the order doesn't matter

SETS

- Set is a collection of distinct objects in which the order doesn't matter
 - {apple, peach} is a set

SETS

- Set is a collection of distinct objects in which the order doesn't matter
 - {apple, peach} is a set
 - {apple, apple, peach} is **not** a set

SETS

- Set is a collection of distinct objects in which the order doesn't matter
 - {apple, peach} is a set
 - {apple, apple, peach} is **not** a set
 - {apple, peach, pear} is the same as {peach, apple, pear}

SOME BASIC NOTATION

SOME BASIC NOTATION

- Set A contains elements a , b and c : $A = \{a, b, c\}$

SOME BASIC NOTATION

- Set A contains elements a , b and c : $A = \{a, b, c\}$
- Element a is in the set A : $a \in A$

SOME BASIC NOTATION

- Set A contains elements a , b and c : $A = \{a, b, c\}$
- Element a is in the set A : $a \in A$
- Set B contains all even numbers:

$$B = \{x \mid x = 2n, n = 0, 1, 2, \dots\}$$

SOME BASIC NOTATION

- Set A contains elements a , b and c : $A = \{a, b, c\}$
- Element a is in the set A : $a \in A$
- Set B contains all even numbers:

$$B = \{x \mid x = 2n, n = 0, 1, 2, \dots\}$$

- Element 5 doesn't belong to set B : $5 \notin B$

FINITE AND INFINITE SETS

- Sets can contain finite number of elements
 - digits: $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$
 - Latin characters: $\{A, B, C, \dots, X, Y, Z\}$
 - all possible 10-digit phone numbers

FINITE AND INFINITE SETS

- Sets can contain finite number of elements
 - digits: $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$
 - Latin characters: $\{A, B, C, \dots, X, Y, Z\}$
 - all possible 10-digit phone numbers
- But sets can also be infinite:
 - natural numbers \mathbb{N} : $\{1, 2, 3, \dots\}$
 - integer numbers \mathbb{Z} : $\{\dots, -2, -1, 0, 1, 2, \dots\}$

FINITE AND INFINITE SETS

- Sets can contain finite number of elements
 - digits: $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$
 - Latin characters: $\{A, B, C, \dots, X, Y, Z\}$
 - all possible 10-digit phone numbers
- But sets can also be infinite:
 - natural numbers \mathbb{N} : $\{1, 2, 3, \dots\}$
 - integer numbers \mathbb{Z} : $\{\dots, -2, -1, 0, 1, 2, \dots\}$

EMPTY SET

- A set with no elements:

EMPTY SET

- A set with no elements:

$$A = \{\}, \quad A = \emptyset$$

CARDINALITY OF A SET

- Cardinality = “how large the set is”.

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.
 - $A = \{\text{apple, banana, peach}\}, \quad |A| = 3$

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.
 - $A = \{\text{apple, banana, peach}\}, \quad |A| = 3$
 - B – a set of Latin characters,

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.
 - $A = \{\text{apple, banana, peach}\}, \quad |A| = 3$
 - B – a set of Latin characters, $|B| = 26$

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.
 - $A = \{\text{apple, banana, peach}\}, \quad |A| = 3$
 - B – a set of Latin characters, $|B| = 26$
 - $C = \{\}$ – an empty set,

CARDINALITY OF A SET

- Cardinality = “how large the set is”.
- Finite sets: cardinality = number of elements.
 - $A = \{\text{apple, banana, peach}\}, \quad |A| = 3$
 - B – a set of Latin characters, $|B| = 26$
 - $C = \{\}$ – an empty set, $|C| = 0$

SUBSET OF A SET

- If each element of a set A is also an element of a set B , we say that A is a subset of B :

$$A \subseteq B$$

SUBSET OF A SET

- If each element of a set A is also an element of a set B , we say that A is a subset of B :

$$A \subseteq B$$

- Example:

SUBSET OF A SET

- If each element of a set A is also an element of a set B , we say that A is a subset of B :

$$A \subseteq B$$

- Example:

A – a set of all numbers between 0 and 100 divisible by 3.

B – a set of all numbers between 0 and 100.

SUBSET OF A SET

- If each element of a set A is also an element of a set B , we say that A is a subset of B :

$$A \subseteq B$$


- Example:

A – a set of all numbers between 0 and 100 divisible by 3.

B – a set of all numbers between 0 and 100.

$$A \subseteq B$$

VISUALIZATION



OPERATIONS WITH SETS

OPERATIONS ON SETS: UNION

- The **union** of A and B is

$$A \cup B = \{x \mid x \in A \text{ or } x \in B \}$$

OPERATIONS ON SETS: UNION

- The **union** of A and B is

$$A \cup B = \{x \mid x \in A \text{ or } x \in B \}$$

- Example:

OPERATIONS ON SETS: UNION

- The **union** of A and B is

$$A \cup B = \{x \mid x \in A \text{ or } x \in B \}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \cup B = \{ \quad \quad \quad \}$

OPERATIONS ON SETS: UNION

- The **union** of A and B is

$$A \cup B = \{x \mid x \in A \text{ or } x \in B \}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \cup B = \{\text{apple, banana, orange, peach}\}$

OPERATIONS ON SETS: INTERSECTION

- The **intersection** of A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B \}$$

OPERATIONS ON SETS: INTERSECTION

- The **intersection** of A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B \}$$

- Example:

OPERATIONS ON SETS: INTERSECTION

- The **intersection** of A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B \}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

OPERATIONS ON SETS: INTERSECTION

- The **intersection** of A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B \}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \cap B = \{ \quad \quad \}$

OPERATIONS ON SETS: INTERSECTION

- The **intersection** of A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B \}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \cap B = \{\text{banana}\}$

OPERATIONS ON SETS: INTERSECTION

- Sets A and B are called **disjoint** if they don't have a common element:

$$A \cap B = \emptyset$$

OPERATIONS ON SETS: INTERSECTION

- Sets A and B are called **disjoint** if they don't have a common element:

$$A \cap B = \emptyset$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, peach}\}$

OPERATIONS ON SETS: INTERSECTION

- Sets A and B are called **disjoint** if they don't have a common element:

$$A \cap B = \emptyset$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, peach}\}$

$$A \cap B = \emptyset$$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \setminus B = \{ \quad \}$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \setminus B = \{\text{apple}\}$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

$A = \{\text{apple, banana}\}$

$B = \{\text{orange, banana, peach}\}$

$A \setminus B = \{\text{apple}\}$

$B \setminus A = \{ \quad \quad \quad \}$

OPERATIONS ON SETS: DIFFERENCE

- The **set difference** of A and B is

$$A \setminus B = A - B = \{x \mid x \in A \text{ and } x \notin B\}$$

- Example:

$A = \{\text{apple, banana}\}$


$B = \{\text{orange, banana, peach}\}$

$A \setminus B = \{\text{apple}\}$


$B \setminus A = \{\text{orange, peach}\}$

OPERATIONS WITH SETS: VENN'S DIAGRAMS


OPERATIONS WITH SETS: VENN'S DIAGRAMS




OPERATIONS WITH SETS: VENN'S DIAGRAMS




OPERATIONS WITH SETS: VENN'S DIAGRAMS




OPERATIONS WITH SETS: VENN'S DIAGRAMS



OPERATIONS WITH SETS: VENN'S DIAGRAMS



OPERATIONS WITH SETS: VENN'S DIAGRAMS




ABSORPTION LAW

$$A \cap (A \cup B) = \dots ?$$


ABSORPTION LAW

$$A \cap (A \cup B) = \dots ?$$




ABSORPTION LAW

$$A \cap (A \cup B) = \dots?$$



ABSORPTION LAW

$$A \cap (A \cup B) = A$$



COMPLEMENT OF A SET


- Let U be the universal set.
 - “*a set of everything*”, whatever that is in the given context.

COMPLEMENT OF A SET


- Let be U the universal set.
 - “*a set of everything*”, whatever that is in the given context.
- **Complement** of A is

$$A^c = A = U \setminus A$$

COMPLEMENT OF A SET: EXAMPLE




COMPLEMENT OF A SET: EXAMPLE




COMPLEMENT OF A SET: EXAMPLE

U Numbers between 0 and 100



COMPLEMENT OF A SET: EXAMPLE


U Numbers between 0 and 100




$$A^c = ?$$

COMPLEMENT OF A SET: EXAMPLE

U Numbers between 0 and 100



COMPLEMENT OF A SET: EXAMPLE



SHORT QUIZ

1. $A \cup \emptyset =$

SHORT QUIZ

1. $A \cup \emptyset = A$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset =$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset = \emptyset$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset = \emptyset$

3. $A \cup U =$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset = \emptyset$

3. $A \cup U = U$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset = \emptyset$

3. $A \cup U = U$

4. $A \cap U =$

SHORT QUIZ

1. $A \cup \emptyset = A$

2. $A \cap \emptyset = \emptyset$

3. $A \cup U = U$

4. $A \cap U = A$

OTHER LAWS

Name	Rule
Commutative laws	$A \cap B = B \cap A, \quad A \cup B = B \cup A$
Associative laws	$A \cap (B \cap C) = (A \cap B) \cap C$ $A \cup (B \cup C) = (A \cup B) \cup C$
Distributive laws	$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
De Morgan's laws	$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c$
Complement laws	$A \cap A^c = \emptyset, \quad A \cup A^c = U$
Double complement law	$(A^c)^c = A$
Idempotent laws	$A \cap A = A, \quad A \cup A = A$
Absorption laws	$A \cap (A \cup B) = A, \quad A \cup (A \cap B) = A$
Dominance laws	$A \cap \emptyset = \emptyset, \quad A \cup U = U$
Identity laws	$A \cup \emptyset = A, \quad A \cap U = A$

CARTESIAN PRODUCT

- Let A and B be two sets.
- **Cartesian product** of two sets A and B is a set, containing all sequences, where the first element is from set A and the second is from set B :

$$A \times B = \{(a, b) \mid a \in A, b \in B\}$$

CARTESIAN PRODUCT: EXAMPLE

- $D = \{\text{wine, beer, water}\}$
- $F = \{\text{spaghetti, burgers}\}$

CARTESIAN PRODUCT: EXAMPLE

- $D = \{\text{wine, beer, water}\}$
- $F = \{\text{spaghetti, burgers}\}$

$$D \times F = \{$$
$$\}$$

CARTESIAN PRODUCT: EXAMPLE

- $D = \{\text{wine, beer, water}\}$
- $F = \{\text{spaghetti, burgers}\}$

$$D \times F = \{(\text{wine, spaghetti}),$$
$$(\text{wine, burgers}),$$
$$\}$$

CARTESIAN PRODUCT: EXAMPLE

- $D = \{\text{wine, beer, water}\}$
- $F = \{\text{spaghetti, burgers}\}$

$$D \times F = \{(\text{wine, spaghetti}),$$
$$(\text{wine, burgers}),$$
$$(\text{beer, spaghetti}),$$
$$(\text{beer, burgers}),$$
$$\}$$

CARTESIAN PRODUCT: EXAMPLE

- $D = \{\text{wine, beer, water}\}$
- $F = \{\text{spaghetti, burgers}\}$

$$D \times F = \{(\text{wine, spaghetti}),$$
$$(\text{wine, burgers}),$$
$$(\text{beer, spaghetti}),$$
$$(\text{beer, burgers}),$$
$$(\text{water, spaghetti}),$$
$$(\text{water, burgers})\}$$

BACK TO COUNTING...

ADDITION PRINCIPLE

(aka Rule of sum)

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
 - Sushi:

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
Dominos, Pizza Hut or King Slice (3 restaurants)
 - Sushi:

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
Dominos, Pizza Hut or King Slice (3 restaurants)
 - Sushi:
Wabi Sabi or Sakura (2 restaurants)

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
Dominos, Pizza Hut or King Slice (3 restaurants)
 - Sushi:
Wabi Sabi or Sakura (2 restaurants)
- How many options are there in total?

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
Dominos, Pizza Hut or King Slice (3 restaurants)
 - Sushi:
Wabi Sabi or Sakura (2 restaurants)
- How many options are there in total?
3 pizza places **+** **2** sushi places =

RULE OF SUM: EXAMPLE

- We want to order dinner for tonight:
 - Pizza:
Dominos, Pizza Hut or King Slice (3 restaurants)
 - Sushi:
Wabi Sabi or Sakura (2 restaurants)
- How many options are there in total?
3 pizza places **+** **2** sushi places = **3 + 2 = 5** options in total

RULE OF SUM

- If there are **a** ways of doing something and **b** number of ways of doing another thing and we can not do both at the same time, then there are **a + b** ways to choose one of the actions.

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

$$|A \cup B| =$$

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

$$|A \cup B| = |A| + |B|$$

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

$$|A \cup B| = |A| + |B|$$

- **Generalized rule of sum:**

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

$$|A \cup B| = |A| + |B|$$

- **Generalized rule of sum:**

If $A_1 \cap A_2 \cap \dots \cap A_n = \emptyset$, then

RULE OF SUM (SET THEORY)

- If A and B are disjoint sets ($A \cap B = \emptyset$), then

$$|A \cup B| = |A| + |B|$$

- **Generalized rule of sum:**

If $A_1 \cap A_2 \cap \dots \cap A_n = \emptyset$, then

$$|A_1 \cup A_2 \cup \dots \cup A_n| = |A_1| + |A_2| + \dots + |A_n|$$

MULTIPLICATION PRINCIPLE

(aka Rule of product)

RULE OF PRODUCT INFORMALLY

- If there are **a** ways of doing something and **b** ways of doing another thing, then there are **$a \cdot b$** ways of performing both actions.

RULE OF PRODUCT: EXAMPLE 1

- You want to order pizza:
 - first, choose the type of crust:
 - second, choose one topping:

RULE OF PRODUCT: EXAMPLE 1

- You want to order pizza:
 - first, choose the type of crust:
thin or thick (2 choices);
 - second, choose one topping:

RULE OF PRODUCT: EXAMPLE 1

- You want to order pizza:
 - first, choose the type of crust:
thin or thick (2 choices);
 - second, choose one topping:
cheese, pepperoni, or sausage (3 choices).

RULE OF PRODUCT: EXAMPLE 1

- You want to order pizza:
 - first, choose the type of crust:
thin or thick (2 choices);
 - second, choose one topping:
cheese, pepperoni, or sausage (3 choices).
- How many different pizzas are there?

RULE OF PRODUCT: EXAMPLE 1

- You want to order pizza:
 - first, choose the type of crust:
thin or thick (2 choices);
 - second, choose one topping:
cheese, pepperoni, or sausage (3 choices).
- How many different pizzas are there?


2 ways of choosing crust **x** **3** ways of choosing topping =

RULE OF PRODUCT: EXAMPLE 1


- You want to order pizza:
 - first, choose the type of crust:
thin or thick (2 choices);
 - second, choose one topping:
cheese, pepperoni, or sausage (3 choices).
- How many different pizzas are there?

2 ways of choosing crust **x** **3** ways of choosing topping =
= 2 x 3 = 6 different pizzas


RULE OF PRODUCT: EXAMPLE 1




RULE OF PRODUCT: EXAMPLE 1



RULE OF PRODUCT: EXAMPLE 1



RULE OF PRODUCT: EXAMPLE 1



RULE OF PRODUCT (SET THEORY)

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

$$|A \times B| =$$

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

$$|A \times B| = |A| \cdot |B|$$

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

$$|A \times B| = |A| \cdot |B|$$

- **Generalized product rule:**

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

$$|A \times B| = |A| \cdot |B|$$

- **Generalized product rule:**

For any sets A_1, A_2, \dots, A_n

RULE OF PRODUCT (SET THEORY)

- For any two sets A and B

$$|A \times B| = |A| \cdot |B|$$

- **Generalized product rule:**

For any sets A_1, A_2, \dots, A_n

$$|A_1 \times A_2 \times \dots \times A_n| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_n|$$

RULE OF PRODUCT: EXAMPLE 2

- One can chose from the following meals during the day:
 - **B**reakfast = {pancakes, eggs, bagel}
 - **L**unch = {salad, soup, bagel}
 - **D**inner = {soup, pasta, burgers}

RULE OF PRODUCT: EXAMPLE 2

- One can choose from the following meals during the day:
 - **B**reakfast = {pancakes, eggs, bagel}
 - **L**unch = {salad, soup, bagel}
 - **D**inner = {soup, pasta, burgers}

How many different day menus can one have?

RULE OF PRODUCT: EXAMPLE 2

- One can chose from the following meals during the day:
 - **B**reakfast = {pancakes, eggs, bagel}
 - **L**unch = {salad, soup, bagel}
 - **D**inner = {soup, pasta, burgers}

How many different day menus can one have?

$$|B| \cdot |L| \cdot |D| = 3 \cdot 3 \cdot 3 = 27$$

RULE OF PRODUCT: EXAMPLE 3



- License plate:
3 letters + 3 digits + regional code
- How many unique plates are there per regional code?

RULE OF PRODUCT: EXAMPLE 3



- License plate:
3 letters + 3 digits + regional code
- How many unique plates are there per regional code?
- 12 letters and 10 digits can be used:

RULE OF PRODUCT: EXAMPLE 3



- License plate:
3 letters + 3 digits + regional code
- How many unique plates are there per regional code?
- 12 letters and 10 digits can be used:

$$12 \cdot 12 \cdot 12 \cdot 10 \cdot 10 \cdot 10$$

RULE OF PRODUCT: EXAMPLE 3



- License plate:
3 letters + 3 digits + regional code
- How many unique plates are there per regional code?
- 12 letters and 10 digits can be used:

$$12 \cdot 12 \cdot 12 \cdot 10 \cdot 10 \cdot 10 = 1\,728\,000 \text{ unique plates}$$

MORE EXAMPLES

Sum and product rules in action

SUM & PRODUCT RULES IN PROGRAMMING

```
for i in range(n):  
    print('hello')
```

```
for j in range(m):  
    print('hello')
```

```
for i in range(n):  
    for j in range(m):  
        print('hello')
```

SUM & PRODUCT RULES IN PROGRAMMING

RULE OF SUM

```
for i in range(n):  
    print('hello')
```

```
for j in range(m):  
    print('hello')
```

```
for i in range(n):  
    for j in range(m):  
        print('hello')
```

SUM & PRODUCT RULES IN PROGRAMMING

RULE OF SUM

```
for i in range(n):  
    print('hello')
```

```
for j in range(m):  
    print('hello')
```

```
for i in range(n):  
    for j in range(m):  
        print('hello')
```

hello will be printed **$m + n$** times

SUM & PRODUCT RULES IN PROGRAMMING

RULE OF SUM

```
for i in range(n):  
    print('hello')  
  
for j in range(m):  
    print('hello')
```

hello will be printed **$m + n$** times

RULE OF PRODUCT

```
for i in range(n):  
    for j in range(m):  
        print('hello')
```


SUM & PRODUCT RULES IN PROGRAMMING

RULE OF SUM

```
for i in range(n):  
    print('hello')  
  
for j in range(m):  
    print('hello')
```

hello will be printed **$m + n$** times

RULE OF PRODUCT

```
for i in range(n):  
    for j in range(m):  
        print('hello')
```

hello will be printed **$m * n$** times

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?
- Solution:

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?

- Solution:

*	*	*	*	*	*
---	---	---	---	---	---

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?

- Solution:

*	*	*	*	*	*
---	---	---	---	---	---

Each position:

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?

- Solution:

*	*	*	*	*	*
---	---	---	---	---	---

Each position: 0 or 1 (2 choices)

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?
- Solution:

*	*	*	*	*	*
---	---	---	---	---	---

Each position: 0 or 1 (2 choices)

Product rule:

NUMBER OF 6-BIT STRINGS

- Example of a 6-bit string:

0	1	1	1	0	1
---	---	---	---	---	---

- How many such strings are there?

- Solution:

*	*	*	*	*	*
---	---	---	---	---	---

Each position: 0 or 1 (2 choices)

Product rule:

$$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^6 = 64 \text{ different 6-bit strings.}$$

NUMBER OF 6-BIT STRINGS

- How many such strings are there that start with *10*?

NUMBER OF 6-BIT STRINGS

- How many such strings are there that start with *10*?
- Solution:

1	0	*	*	*	*
---	---	---	---	---	---

NUMBER OF 6-BIT STRINGS

- How many such strings are there that start with *10*?
- Solution:

1	0	*	*	*	*
---	---	---	---	---	---

Positions 1 & 2: 10

Other positions: 0 or 1 (2 choices)

NUMBER OF 6-BIT STRINGS

- How many such strings are there that start with *10*?
- Solution:

1	0	*	*	*	*
---	---	---	---	---	---

Positions 1 & 2: 10

Other positions: 0 or 1 (2 choices)

Product rule:

NUMBER OF 6-BIT STRINGS

- How many such strings are there that start with *10*?
- Solution:

1	0	*	*	*	*
---	---	---	---	---	---

Positions 1 & 2: 10

Other positions: 0 or 1 (2 choices)

Product rule:

$$1 \cdot 1 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^4 = 16$$


different 6-bit strings that start with **10**.

NUMBER OF 6-BIT STRINGS

- How many such strings are there that **don't** start with *10*?


NUMBER OF 6-BIT STRINGS

- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



NUMBER OF 6-BIT STRINGS


- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



SOLUTION 1 |strings that **don't** start with *10*| =

NUMBER OF 6-BIT STRINGS


- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



SOLUTION 1 |strings that **don't** start with *10*| =
|strings that start with *00*| + |strings that start with *01*| +
+ |strings that start with *11*|

NUMBER OF 6-BIT STRINGS

- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**




SOLUTION 1 |strings that **don't** start with *10*| =
|strings that start with *00*| + |strings that start with *01*| +
+ |strings that start with *11*|

$$2^4 + 2^4 + 2^4 = 16 + 16 + 16 = 48$$

NUMBER OF 6-BIT STRINGS


- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



SOLUTION 2 |strings that **don't** start with *10*| =

NUMBER OF 6-BIT STRINGS


- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



SOLUTION 2 $|\text{strings that **don't** start with } 10| =$
 $|\text{all strings}| - |\text{strings that **do** start with } 10|$

NUMBER OF 6-BIT STRINGS

- How many such strings are there that **don't** start with *10*?
- Solution: **not 10**



SOLUTION 2

$$\begin{aligned} & |\text{strings that **don't** start with } 10| = \\ & |\text{all strings}| - |\text{strings that **do** start with } 10| \end{aligned}$$

$$2^6 - 2^4 = 64 - 16 = 48$$

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?
- Solution:

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?
- Solution:

1. *Sum rule:*

$$N = N_6 + N_5 + N_4 + N_3 + N_2 + N_1, \quad N_i = \# \text{ bit strings of length } i$$

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?
- Solution:

1. *Sum rule:*

$$N = N_6 + N_5 + N_4 + N_3 + N_2 + N_1, \quad N_i = \# \text{ bit strings of length } i$$

$$N_i =$$

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?
- Solution:

1. *Sum rule:*

$$N = N_6 + N_5 + N_4 + N_3 + N_2 + N_1, \quad N_i = \# \text{ bit strings of length } i$$

2. *Product rule:*

$$N_i = 2^i$$

NUMBER OF BIT STRINGS OF LENGTH 6 OR LESS

- How many bit strings are there of length 6 or less?
- Solution:

1. *Sum rule:*

$$N = N_6 + N_5 + N_4 + N_3 + N_2 + N_1, \quad N_i = \# \text{ bit strings of length } i$$

2. *Product rule:*

$$N_i = 2^i$$

$$N = 2^6 + 2^5 + 2^4 + 2^3 + 2^2 + 2^1 = 126$$

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.
- 0 elements:
- 1 element:
- 2 elements:
- 3 elements:

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.

- 0 elements:

$\{\}$

- 1 element:

- 2 elements:

- 3 elements:

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.

- 0 elements:

$\{\}$

- 1 element:

$\{a\}$ $\{b\}$ $\{c\}$

- 2 elements:

- 3 elements:

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.

- 0 elements:

$\{\}$

- 1 element:

$\{a\}$ $\{b\}$ $\{c\}$

- 2 elements:

$\{a, b\}$ $\{a, c\}$ $\{b, c\}$

- 3 elements:

SUBSETS OF A SET

- Consider a set $\{a, b, c\}$. How many subsets does it have? List them all.

- 0 elements:

$\{\}$

- 1 element:

$\{a\}$ $\{b\}$ $\{c\}$

- 2 elements:

$\{a, b\}$ $\{a, c\}$ $\{b, c\}$

- 3 elements:

$\{a, b, c\}$

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?
- **Solution:**

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

- **Solution:**

Each element can either be included in the subset (1) or not (0):

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

- **Solution:**

Each element can either be included in the subset (1) or not (0):

0	1	0	...	0	0	1
---	---	---	-----	---	---	---

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

- **Solution:**

Each element can either be included in the subset (1) or not (0):

0	1	0	...	0	0	1
---	---	---	-----	---	---	---

2 options per each of the n element (***just like bit-strings!***).

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

- **Solution:**

Each element can either be included in the subset (1) or not (0):

0	1	0	...	0	0	1
---	---	---	-----	---	---	---

2 options per each of the n element (***just like bit-strings!***).

Product rule:

NUMBER OF SUBSETS OF A SET

- A 3-element set has 8 subsets.
- How many subsets does a set of n elements have?

- **Solution:**

Each element can either be included in the subset (1) or not (0):

0	1	0	...	0	0	1
---	---	---	-----	---	---	---

2 options per each of the n element (***just like bit-strings!***).

Product rule: $2 \cdot 2 \cdot \dots \cdot 2 = 2^n$ subsets

EXAMPLE: NUMBER OF PASSWORDS

- A valid password contains 6 to 8 symbols
 - first symbol is a letter (upper- or lowercase);
 - other symbols are letters (upper- or lowercase) or digits.
- How many different passwords are there?

EXAMPLE: NUMBER OF PASSWORDS

- A valid password contains 6 to 8 symbols
 - first symbol is a letter (upper- or lowercase);
 - other symbols are letters (upper- or lowercase) or digits.
- How many different passwords are there?
- Solution:

EXAMPLE: NUMBER OF PASSWORDS

- A valid password contains 6 to 8 symbols
 - first symbol is a letter (upper- or lowercase);
 - other symbols are letters (upper- or lowercase) or digits.
- How many different passwords are there?
- Solution:
Sum rule:

EXAMPLE: NUMBER OF PASSWORDS

- A valid password contains 6 to 8 symbols
 - first symbol is a letter (upper- or lowercase);
 - other symbols are letters (upper- or lowercase) or digits.
- How many different passwords are there?

- Solution:

Sum rule:

$$N = P_6 + P_7 + P_8 \qquad P_i - \# \text{ passwords of length } i$$

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter

Other symbols:

upper- or lowercase letter or digit

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

$$P_6 =$$

$$P_7 =$$

$$P_8 =$$

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

Product rule: $P_6 =$ $P_7 =$ $P_8 =$

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

Product rule: $P_6 = 52 \cdot 62^5$ $P_7 =$ $P_8 =$

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

Product rule: $P_6 = 52 \cdot 62^5$ $P_7 = 52 \cdot 62^6$ $P_8 = 52 \cdot 62^7$

EXAMPLE: NUMBER OF PASSWORDS

- Solution:

Sum rule: $N = P_6 + P_7 + P_8$ P_i – # passwords of length i

First symbol:

upper- or lowercase letter – $2 \cdot 26 = 52$ options

Other symbols:

upper- or lowercase letter or digit – $52 + 10 = 62$ options

Product rule: $P_6 = 52 \cdot 62^5$ $P_7 = 52 \cdot 62^6$ $P_8 = 52 \cdot 62^7$

$$N = 52 \cdot 62^5 + 52 \cdot 62^6 + 52 \cdot 62^7$$

OTHER USEFUL TRICKS

COUNTING ONE THING BY COUNTING ANOTHER

- How to count the number of people in the class?

COUNTING ONE THING BY COUNTING ANOTHER

- How to count the number of people in the class?
 - Count heads

COUNTING ONE THING BY COUNTING ANOTHER

- How to count the number of people in the class?
 - Count heads
 - Count their laptops

COUNTING ONE THING BY COUNTING ANOTHER

- How to count the number of people in the class?
 - Count heads
 - Count their laptops
 - Count ears and divide by 2 😊
(assuming that everyone has exactly two ears)



NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?
- Solution:

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?
- Solution:
Every person (n) shakes hands with everyone else:

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?

- Solution:

Every person (n) shakes hands with everyone else ($n-1$):

$n(n - 1)$ handshakes?

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?

- Solution:

Every person (n) shakes hands with everyone else ($n-1$):

$$n(n - 1) \text{ handshakes?}$$

- We counted each handshake twice!

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?

- Solution:

Every person (n) shakes hands with everyone else ($n-1$):

$$n(n - 1) \text{ handshakes?}$$

- We counted each handshake twice!

Fixing it:

NUMBER OF HANDSHAKES

- There are n people in a room. Everyone shakes hands with everyone. How many handshakes are there?

- Solution:

Every person (n) shakes hands with everyone else ($n-1$):

$$n(n - 1) \text{ handshakes?}$$

- We counted each handshake twice!

Fixing it: $\frac{n(n-1)}{2}$

LET'S PRACTICE

https://docs.google.com/document/d/1XM0qoghJNkjnrVL3QVb_cSYRnfz1qdf7CLK6yFezF18/edit?usp=sharing

TO SUM UP

- The basics of set theory.
- Basic counting principles:
 - sum rule;
 - product rule;
 - combining the rules.
- Other tricks
 - using complement of a set;
 - overcounting.