

# **ELEMENTARY COMBINATORICS & PROBABILITY**

Lecture 11

Random variables

# LAST TIME

- Random variables
- Probability mass function
- Cumulative distribution function

# TODAY

- Review PMFs and CDFs
- Special distributions

# PMF AND CDF

- For a discrete random variable  $X$  with range  $R_X = \{x_1, x_2, \dots\}$  **probability mass function (PMF)** is defined as

$$P_X(x) = \begin{cases} P(X = x), & x \in R_X \\ 0, & x \notin R_X \end{cases}$$

# PMF AND CDF

- For a discrete random variable  $X$  with range  $R_X = \{x_1, x_2, \dots\}$  **probability mass function (PMF)** is defined as

$$P_X(x) = \begin{cases} P(X = x), & x \in R_X \\ 0, & x \notin R_X \end{cases}$$

- **Cumulative distribution function (CDF)** of a random variable  $X$  is defined as follows:

$$F_X(x) = P(X \leq x) \quad \forall x \in \mathbb{R}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \left\{ \right.$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} & x = 0 \\ & x = 1 \\ & x = 3 \\ \textit{otherwise} \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ & x = 1 \\ & x = 2 \\ & \text{otherwise} \end{cases}$$



# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ & x = 2 \\ & \textit{otherwise} \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ \textit{otherwise} & \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 3 \\ 0, & \textit{otherwise} \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.25, & 0 \leq x < 1 \\ 0.75, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0 & x < 0 \\ 0.25 & 0 \leq x < 1 \\ 0.75 & 1 \leq x < 2 \\ 1 & x \geq 2 \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.25, & 0 \leq x < 1 \\ 0.75, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.25, & 0 \leq x < 1 \\ 0.75, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.25, & 0 \leq x < 1 \\ 0.75, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$



# PMF AND CDF: EXAMPLE

- $X$  – total number of heads after two tosses of a coin.

$$P_X(x) = P(X = x) = \begin{cases} 0.25, & x = 0 \\ 0.5, & x = 1 \\ 0.25, & x = 2 \\ 0, & \text{otherwise} \end{cases}$$

$$F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.25, & 0 \leq x < 1 \\ 0.75, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{ \quad \}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \left\{ \begin{array}{l} \\ \\ \end{array} \right. \quad F_X(x) = P(X \leq x) = \left\{ \begin{array}{l} \\ \\ \end{array} \right.$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} x = 0 \\ x = 1 \\ x = 2 \\ otherwise \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ & x = 1 \\ & x = 2 \\ otherwise \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ & x = 2 \\ \textit{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} \end{cases}$$



# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} & x < 0 \\ 0 \leq x < 1 \\ 1 \leq x < 2 \\ x \geq 2 \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.1, & 0 \leq x < 1 \\ 0.7, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.1, & 0 \leq x < 1 \\ 0.7, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.1, & 0 \leq x < 1 \\ 0.7, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# BALLS IN A BOWL

- A bowl contains **two red balls** and **three blue balls**. Three balls are selected at random from the bowl. Let random variable  $X$  denote the number of red balls among them.

$$R_X = \{0, 1, 2\}$$

$$P_X(x) = P(X = x) = \begin{cases} 0.1, & x = 0 \\ 0.6, & x = 1 \\ 0.3, & x = 2 \\ 0, & \text{otherwise} \end{cases} \quad F_X(x) = P(X \leq x) = \begin{cases} 0, & x < 0 \\ 0.1, & 0 \leq x < 1 \\ 0.7, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.



# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{ \quad \quad \quad \}$$

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$						
$P(X = x)$						

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$						

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$				$\frac{1 + 2 \cdot 3}{36}$		

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$			$\frac{1 + 2 \cdot 3}{36}$		

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$	$\frac{1 + 2}{36}$		$\frac{1 + 2 \cdot 3}{36}$		

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$	$\frac{1 + 2}{36}$	$\frac{1 + 2 \cdot 2}{36}$	$\frac{1 + 2 \cdot 3}{36}$		



# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$	$\frac{1 + 2}{36}$	$\frac{1 + 2 \cdot 2}{36}$	$\frac{1 + 2 \cdot 3}{36}$	$\frac{1 + 2 \cdot 4}{36}$	

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$	$\frac{1 + 2}{36}$	$\frac{1 + 2 \cdot 2}{36}$	$\frac{1 + 2 \cdot 3}{36}$	$\frac{1 + 2 \cdot 4}{36}$	$\frac{1 + 2 \cdot 5}{36}$

# ROLLING DICE

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- Distribution of  $X$ :

$x$	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \left\{ \begin{array}{l} \end{array} \right.$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} x < 1 \\ 1 \leq x < 2 \\ 2 \leq x < 3 \\ 3 \leq x < 4 \\ 4 \leq x < 5 \\ 5 \leq x < 6 \\ x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1 & 1 \leq x < 2 \\ 2 & 2 \leq x < 3 \\ 3 & 3 \leq x < 4 \\ 4 & 4 \leq x < 5 \\ 5 & 5 \leq x < 6 \\ 6 & x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ & 2 \leq x < 3 \\ & 3 \leq x < 4 \\ & 4 \leq x < 5 \\ & 5 \leq x < 6 \\ & x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ 4/36, & 2 \leq x < 3 \\ & 3 \leq x < 4 \\ & 4 \leq x < 5 \\ & 5 \leq x < 6 \\ & x \geq 6 \end{cases}$$



# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ 4/36, & 2 \leq x < 3 \\ 9/36, & 3 \leq x < 4 \\ & 4 \leq x < 5 \\ & 5 \leq x < 6 \\ & x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ 4/36, & 2 \leq x < 3 \\ 9/36, & 3 \leq x < 4 \\ 16/36, & 4 \leq x < 5 \\ & 5 \leq x < 6 \\ & x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$

- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ 4/36, & 2 \leq x < 3 \\ 9/36, & 3 \leq x < 4 \\ 16/36, & 4 \leq x < 5 \\ 25/36, & 5 \leq x < 6 \\ & x \geq 6 \end{cases}$$

# ROLLING DICE

$x$	1	2	3	4	5	6
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$


- You are rolling a pair of dice. Let random variable  $X$  denote the maximum of the two numbers you got.

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

- CDF of  $X$ :

$$F_X(x) = \begin{cases} 0, & x < 1 \\ 1/36, & 1 \leq x < 2 \\ 4/36, & 2 \leq x < 3 \\ 9/36, & 3 \leq x < 4 \\ 16/36, & 4 \leq x < 5 \\ 25/36, & 5 \leq x < 6 \\ 1, & x \geq 6 \end{cases}$$

# CDF OF A DISCRETE RANDOM VARIABLE



# PMF AND CDF: BASIC PROPERTIES

- Which basic properties does a PMF function have?
  - $0 \leq P(x) \leq 1$
  - All non-zero values sum up to one

# PMF AND CDF: BASIC PROPERTIES

- Which basic properties does a PMF function have?
  - $0 \leq P(x) \leq 1$
  - All non-zero values sum up to one
- Which basic properties does a CDF function have?
  - $0 \leq F(x) \leq 1$
  - $F(x)$  is non-decreasing

# ROLLING DICE: PMF AND CDF

Google Classroom -> Programming exercise



# **SPECIAL DISTRIBUTIONS**

# BERNOULLI DISTRIBUTION

- Consider a random experiment with two possible outcomes:  
“success” (with probability  $p$ ) or “failure” (with probability  $1 - p$ )
  - tossing a coin: H or T;
  - a new child: a boy or a girl;
  - you take an exam: pass or fail.

# BERNOULLI DISTRIBUTION

- Consider a random experiment with two possible outcomes:  
“success” (with probability  $p$ ) or “failure” (with probability  $1 - p$ )
  - tossing a coin: H or T;
  - a new child: a boy or a girl;
  - you take an exam: pass or fail.
- Consider a random variable  $X$

$x$	0	1
$P(X = x)$		

# BERNOULLI DISTRIBUTION

- Consider a random experiment with two possible outcomes:  
“success” (with probability  $p$ ) or “failure” (with probability  $1 - p$ )
  - tossing a coin: H or T;
  - a new child: a boy or a girl;
  - you take an exam: pass or fail.
- Consider a random variable  $X$

$x$	0	1
$P(X = x)$	$1 - p$	$p$

# BERNOULLI DISTRIBUTION

- Consider a random experiment with two possible outcomes:  
“success” (with probability  $p$ ) or “failure” (with probability  $1 - p$ )
  - tossing a coin: H or T;
  - a new child: a boy or a girl;
  - you take an exam: pass or fail.
- Consider a random variable  $X \sim \text{Bernoulli}(p)$

$x$	0	1
$P(X = x)$	$1 - p$	$p$

# BERNOULLI DISTRIBUTION

- A random variable  $X$  is said to be a Bernoulli random variable with parameter  $p$  if its PMF is given by

$$P_X(x) = P(X = x) = \begin{cases} p, & x = 1 \\ 1 - p, & x = 0 \\ 0, & \text{otherwise} \end{cases}$$

# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.

# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.
- $X$ - a random variable:  $X = 1$  if guessed correctly and  $X = 0$  otherwise.



# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.
- $X$ - a random variable:  $X = 1$  if guessed correctly and  $X = 0$  otherwise.

$$X \sim \text{Bernoulli}( \quad )$$

# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.
- $X$ - a random variable:  $X = 1$  if guessed correctly and  $X = 0$  otherwise.

$$X \sim \text{Bernoulli}(0.25)$$

# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.
- $X$ - a random variable:  $X = 1$  if guessed correctly and  $X = 0$  otherwise.

$$X \sim \text{Bernoulli}(0.25)$$

$$P_X(x) = P(X = x) =$$

# EXAMPLE: PASSING AN EXAM

- You need to answer a multiple-choice question (4 options, only 1 is correct).
- You don't know the answer, so you are randomly guessing.
- $X$ - a random variable:  $X = 1$  if guessed correctly and  $X = 0$  otherwise.

$$X \sim \text{Bernoulli}(0.25)$$

$$P_X(x) = P(X = x) = \begin{cases} 0.75, & x = 0 \\ 0.25, & x = 1 \\ 0, & \text{otherwise} \end{cases}$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X$  denotes if you get heads:

$$R_X = \{0,1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0,1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0,1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{ \quad \quad \quad \}$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0,1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{0, 1, 2, \dots, n\}$$



# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0, 1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{0, 1, 2, \dots, n\}$$

$$P(Y = k) =$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0, 1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{0, 1, 2, \dots, n\}$$

$$P(Y = k) = P(kH, (n - k)T) =$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0, 1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{0, 1, 2, \dots, n\}$$

$$P(Y = k) = P(kH, (n - k)T) = C(n, k) \cdot p^k (1 - p)^{n-k}$$

# BINOMIAL DISTRIBUTION

- You toss a coin once. Random variable  $X \sim \text{Bernoulli}(p)$  denotes if you get heads:

$$R_X = \{0, 1\}, \quad P(X = 0) = P(T) = 1 - p, \quad P(X = 1) = P(H) = p$$

- Imagine now that you toss a coin  $n$  times. Random variable  $Y$  is the total number of heads you get:

$$R_Y = \{0, 1, 2, \dots, n\}$$

$$P(Y = k) = P(kH, (n - k)T) = C(n, k) \cdot p^k (1 - p)^{n-k}$$

$$Y \sim \text{Binomial}(n, p)$$

# BINOMIAL DISTRIBUTION

- A random variable  $Y$  is said to follow Binomial distribution with parameters  $n$  and  $p$  if its PMF is given by

$$P_Y(k) = P(Y = k) = \begin{cases} C(n, k) \cdot p^k (1 - p)^{n-k}, & k = 0, 1, \dots, n \\ 0, & \text{otherwise} \end{cases}$$

# BINOMIAL DISTRIBUTION

- A random variable  $Y$  is said to follow Binomial distribution with parameters  $n$  and  $p$  if its PMF is given by

$$P_Y(k) = P(Y = k) = \begin{cases} C(n, k) \cdot p^k (1 - p)^{n-k}, & k = 0, 1, \dots, n \\ 0, & \text{otherwise} \end{cases}$$

- Represents the number of successes in a series of  $n$  independent Bernoulli trials, each of which results in success with probability  $p$ .

# EXAMPLE: EXAM

- You are randomly guessing the answers to 5 multiple choice questions (4 options, 1 correct in each).
- $Y \sim \text{Binomial}(5, 0.25)$  – number of correctly guessed answers.

$$P_Y(k) = C(n, k)p^k(1 - p)^{n-k} = \left\{ \right.$$

# EXAMPLE: EXAM

- You are randomly guessing the answers to 5 multiple choice questions (4 options, 1 correct in each).
- $Y \sim \text{Binomial}(5, 0.25)$  – number of correctly guessed answers.

$$P_Y(k) = C(n, k)p^k(1 - p)^{n-k} = \begin{cases} 0.75^5, & k = 0 \\ 5 \cdot 0.25 \cdot 0.75^4, & k = 1 \\ 10 \cdot 0.25^2 \cdot 0.75^3, & k = 2 \\ 10 \cdot 0.25^3 \cdot 0.75^2, & k = 3 \\ 5 \cdot 0.25^4 \cdot 0.75, & k = 4 \\ 0.25^5, & k = 5 \\ 0, & \text{otherwise} \end{cases}$$



# EXAMPLE: EXAM

- $Y \sim \text{Binomial}(5, 0.25)$  – number of correctly guessed answers.

$$P_Y(k) = C(n, k)p^k(1 - p)^{n-k} = \begin{cases} 0.75^5, & k = 0 \\ 5 \cdot 0.25 \cdot 0.75^4, & k = 1 \\ 10 \cdot 0.25^2 \cdot 0.75^3, & k = 2 \\ 10 \cdot 0.25^3 \cdot 0.75^2, & k = 3 \\ 5 \cdot 0.25^4 \cdot 0.75, & k = 4 \\ 0.25^5, & k = 5 \\ 0, & \text{otherwise} \end{cases}$$

- What are the chances of passing the test (= guessing 3 or more)?

$$P(Y \geq 3) =$$

# EXAMPLE: EXAM

- $Y \sim \text{Binomial}(5, 0.25)$  – number of correctly guessed answers.

$$P_Y(k) = C(n, k)p^k(1 - p)^{n-k} = \begin{cases} 0.75^5, & k = 0 \\ 5 \cdot 0.25 \cdot 0.75^4, & k = 1 \\ 10 \cdot 0.25^2 \cdot 0.75^3, & k = 2 \\ 10 \cdot 0.25^3 \cdot 0.75^2, & k = 3 \\ 5 \cdot 0.25^4 \cdot 0.75, & k = 4 \\ 0.25^5, & k = 5 \\ 0, & \text{otherwise} \end{cases}$$

- What are the chances of passing the test (= guessing 3 or more)?

$$P(Y \geq 3) = P_Y(3) + P_Y(4) + P_Y(5) \sim 0.1035$$