

ELEMENTARY COMBINATORICS & PROBABILITY

Lecture 5

Pigeonhole principle, review

LAST TIME

- Review permutations and combinations
- Tuples
 - Ordered sequence with repetitions
- Combinations with repetitions
 - In how many ways can we color k balls with n colors?

TODAY

- Problem set 4
 - Permutations, combinations, tuples, ...
- Pigeonhole principle
- Problem set 5

- Graded assignment 2 is out
- Deadline: Monday, March 22, 23:59 Barcelona time
- See Google classroom

REVIEW


DIFFERENT ARRANGEMENTS

- Imagine you have n objects. How can you arrange them?

	WITHOUT REPETITIONS	WITH REPETITIONS
ORDERED	<p><i>PERMUTATIONS</i></p> <p>Seating n people in a row</p> <p>$n!$</p>	<p><i>TUPLES</i></p> <p>Counting different n-bit strings are there?</p> <p>k^n</p>
NOT ORDERED	<p><i>COMBINATIONS</i></p> <p>Choosing k out of n different candies in a bag</p> <p>$C(n, k) = \frac{n!}{k!(n-k)!}$</p>	<p><i>COMBINATIONS with repetitions</i></p> <p>Distributing k identical candies among n kids</p> <p>$C(k+n-1, n-1)$</p>

COMBINATIONS WITH REPETITIONS


- How many ways are there to place 4 colored balls in a bag, when each ball should be either Red, Green, or Blue?



COMBINATIONS WITH REPETITIONS



- Where to put $3 - 1 = 2$ bars?

○|○|○○ |○|○○○ ○○○○|



COMBINATIONS WITH REPETITIONS

- Where to put $3 - 1 = 2$ bars?



$$C(4 + 3 - 1, 3 - 1) = C(6, 2) = \frac{6!}{2! 4!} = 15$$

COMBINATIONS WITH REPETITIONS

- How many ways are there to place k colored balls in a bag, when each ball can be of one of the n colours?

$$C(k + n - 1, n - 1)$$

BAGELS

- Chris is ordering bagels for three friends he's studying with, as well as one for himself. The bagel shop sells eight varieties of bagel. In how many ways can he choose the bagels to give to Jan, Tom, Olive, and himself?

BAGELS

- Chris is ordering bagels for three friends he's studying with, as well as one for himself. The bagel shop sells eight varieties of bagel. In how many ways can he choose the bagels to give to Jan, Tom, Olive, and himself?

It matters who gets which bagel → ordered arrangement.

BAGELS

- Chris is ordering bagels for three friends he's studying with, as well as one for himself. The bagel shop sells eight varieties of bagel. In how many ways can he choose the bagels to give to Jan, Tom, Olive, and himself?

It matters who gets which bagel → ordered arrangement.
Repetitions allowed.

BAGELS

- Chris is ordering bagels for three friends he's studying with, as well as one for himself. The bagel shop sells eight varieties of bagel. In how many ways can he choose the bagels to give to Jan, Tom, Olive, and himself?

It matters who gets which bagel → ordered arrangement.
Repetitions allowed.

$$8 \cdot 8 \cdot 8 \cdot 8 = 8^4$$

ways to choose the bagels

DOUGHNUTS, PART I

- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

DOUGHNUTS, PART I

- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

DOUGHNUTS, PART I

- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.
Repetitions are allowed

DOUGHNUTS, PART I

- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions

DOUGHNUTS, PART I


- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



DOUGHNUTS, PART I


- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



DOUGHNUTS, PART I


- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



$$C(8 + 5 - 1, 5 - 1) =$$

DOUGHNUTS, PART I


- Chris also went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there for Chris to fill this order?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



$$C(8 + 5 - 1, 5 - 1) = C(12, 4) = \frac{12!}{4!8!} = 495$$

ways to fill a box of 8 doughnuts

DOUGHNUTS, PART 1


This corresponds to the number of non-negative integer solutions of which equation?

A. $x_1 + x_2 + x_3 + x_4 = 12$

C. $x_1 + x_2 + \cdots + x_{11} + x_{12} = 4$

B. $x_1 + x_2 + x_3 + x_4 + x_5 = 8$

D. $x_1 + x_2 + \cdots + x_7 + x_8 = 5$



$$C(8 + 4 - 1, 4 - 1) = C(12, 4) = \frac{12!}{4!8!} = 495$$

ways to fill a box of 8 doughnuts

DOUGHNUTS, PART 1


This corresponds to the number of non-negative integer solutions of which equation?

A. $x_1 + x_2 + x_3 + x_4 = 12$

C. $x_1 + x_2 + \cdots + x_{11} + x_{12} = 4$

B. $x_1 + x_2 + x_3 + x_4 + x_5 = 8$

D. $x_1 + x_2 + \cdots + x_7 + x_8 = 5$



$$C(8 + 4 - 1, 4 - 1) = C(12, 3) = \frac{12!}{3!9!} = 220$$

ways to fill a box of 8 doughnuts

DOUGHNUTS, PART II

- In how many ways can 8 doughnuts (two chocolate, three maple and three vanilla ones) be given to 8 people?

DOUGHNUTS, PART II

- In how many ways can 8 doughnuts (two chocolate, three maple and three vanilla ones) be given to 8 people?

Doughnuts are assigned to different people ordered arrangement

DOUGHNUTS, PART II

- In how many ways can 8 doughnuts (two chocolate, three maple and three vanilla ones) be given to 8 people?

Doughnuts are assigned to different people ordered arrangement
Some objects are repeated

DOUGHNUTS, PART II

- In how many ways can 8 doughnuts (two chocolate, three maple and three vanilla ones) be given to 8 people?

Doughnuts are assigned to different people ordered arrangement
Some objects are repeated

$$\frac{8!}{2! \cdot 3! \cdot 3!}$$

unique ways to “order” repeating doughnuts

PROBLEM SET 4

Permutations and combinations

A PARTY – PART 1

- Five couples need to hold a meeting dedicated to the planning of the party. The meeting should consist of five people, one from each couple. How many possible ways do they have to pick people for the meeting?

A PARTY – PART 1

- Five couples need to hold a meeting dedicated to the planning of the party. The meeting should consist of five people, one from each couple. How many possible ways do they have to pick people for the meeting?

We need a representative from every couple:

A PARTY – PART 1

- Five couples need to hold a meeting dedicated to the planning of the party. The meeting should consist of five people, one from each couple. How many possible ways do they have to pick people for the meeting?

We need a representative from every couple:

$$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^5$$

A PARTY – PART 2

- Five couples need to pick three couples who will be responsible for bringing food for the party. How many ways are there to do this?

A PARTY – PART 2

- Five couples need to pick three couples who will be responsible for bringing food for the party. How many ways are there to do this?

Order doesn't matter

A PARTY – PART 2

- Five couples need to pick three couples who will be responsible for bringing food for the party. How many ways are there to do this?

Order doesn't matter

$$C(5,3) = \frac{5!}{3! 2!} = 10$$

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first chose three couples:

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first chose three couples: $C(5,3)$

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first choose three couples: $C(5,3)$

Now, let's choose one of the two people in each couple:

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first choose three couples: $C(5,3)$

Now, let's choose one of the two people in each couple: $2 \cdot 2 \cdot 2$

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first choose three couples: $C(5,3)$

Now, let's choose one of the two people in each couple: $2 \cdot 2 \cdot 2$

Putting it all together:

A PARTY – PART 3

- Five couples need to choose three people who will be responsible for cleaning up after the party. People from the same couple should not be picked for this. How many ways are there to do this?

Let's first choose three couples: $C(5,3)$

Now, let's choose one of the two people in each couple: $2 \cdot 2 \cdot 2$

Putting it all together:

$$C(5,3) \cdot 2 \cdot 2 \cdot 2 = 80$$

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

Order doesn't matter →

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

Order doesn't matter → counting combinations.

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

Order doesn't matter → counting combinations.

+ Product rule:

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

Order doesn't matter → counting combinations.

+ Product rule:

$$C(7,3) \cdot C(5,3) =$$

BOOK CLUB

- Alice has 7 textbooks and Bob has 5 textbooks. All textbooks are different. Alice gives Bob three of her books and Bob gives Alice three of his books. How many ways do they have to do it?

Order doesn't matter → counting combinations.

+ Product rule:

$$C(7,3) \cdot C(5,3) = \frac{7!}{3!4!} \cdot \frac{5!}{3!2!}$$

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

→ counting combinations
with repetitions

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

→ counting combinations
with repetitions

slices = “balls”

types of pizza = “colour”, category

PIZZA


- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

→ counting combinations
with repetitions

slices = “balls”

types of pizza = “colour”, category



PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

→ counting combinations
with repetitions

slices = “balls”

types of pizza = “colour”, category



$$C(10 + 5 - 1, 5 - 1) =$$

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.

Order doesn't matter
Repetitions are allowed

→ counting combinations
with repetitions

slices = “balls”

types of pizza = “colour”, category

○|○|○ ○ ○ ○|○|○ ○ ○

$$C(10 + 5 - 1, 5 - 1) = C(14, 4) = \frac{14!}{10!4!} = \frac{14 \cdot 13 \cdot 12 \cdot 11}{4!} = 1001$$

ways of eating 10 slices of pizza of 5 types

PIZZA


- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.



Alternative representation:

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.



Alternative representation:

$x_i \geq 0$ – number of slices of pizza of type i .

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.




Alternative representation:

$x_i \geq 0$ – number of slices of pizza of type i .

You are planning to eat 10 pieces, so

PIZZA

- There are 5 types of pizza. You are planning to eat 10 slices in total. In how many ways can you do this? Order in which you eat the slices doesn't matter.



Alternative representation:

$x_i \geq 0$ – number of slices of pizza of type i .

You are planning to eat 10 pieces, so

$$x_1 + x_2 + x_3 + x_4 + x_5 = 10$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces?

PIRATES

- In how many ways can 6 pirates split 50 gold pieces?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50, \quad x_i \geq 0$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50, \quad x_i \geq 0$$

50 gold pieces = 50 “balls”, 6 pirates = 6 “colours”, “categories”

PIRATES

- In how many ways can 6 pirates split 50 gold pieces?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50, \quad x_i \geq 0$$

50 gold pieces = 50 “balls”, 6 pirates = 6 “colours”, “categories”

$$C(50 + 6 - 1, 6 - 1) =$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50, \quad x_i \geq 0$$

50 gold pieces = 50 “balls”, 6 pirates = 6 “colours”, “categories”

$$C(50 + 6 - 1, 6 - 1) = C(55, 5) = \frac{55!}{5!50!} = \frac{55 \cdot 54 \cdot 53 \cdot 52 \cdot 51}{5 \cdot 4 \cdot 3 \cdot 2}$$

ways of distributing the gold without constraints

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50,$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50,$$
$$x_1 \geq 10, \quad x_2 \geq 8, \quad x_3 \geq 4, \quad x_5 \geq 2, \quad x_6 \geq 0$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50,$$
$$x_1 \geq 10, \quad x_2 \geq 8, \quad x_3 \geq 4, \quad x_5 \geq 2, \quad x_6 \geq 0$$

$$y_1 = x_1 - 10, \quad y_2 = x_2 - 8, \quad y_3 = x_3 - 4, \quad y_4 = x_4 - 2,$$
$$y_5 = x_5 - 2, \quad y_6 = x_6, \quad y_i \geq 0$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 50,$$
$$x_1 \geq 10, \quad x_2 \geq 8, \quad x_3 \geq 4, \quad x_5 \geq 2, \quad x_6 \geq 0$$

$$y_1 = x_1 - 10, \quad y_2 = x_2 - 8, \quad y_3 = x_3 - 6, \quad y_4 = x_4 - 4,$$
$$y_5 = x_5 - 2, \quad y_6 = x_6, \quad y_i \geq 0$$

$$y_1 + 10 + y_2 + 8 + y_3 + 6 + y_4 + 4 + y_5 + 2 + y_6 = 50$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 20, \quad y_i \geq 0$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 20, \quad y_i \geq 0$$

20 “balls”, 6 “colours”

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 20, \quad y_i \geq 0$$

20 “balls”, 6 “colours”

$$C(20 + 6 - 1, 6 - 1) =$$

PIRATES

- In how many ways can 6 pirates split 50 gold pieces if the first pirate needs at least 10 pieces, the second needs at least 8, the third needs at least 6, the fourth needs at least 4, and the fifth needs at least 2, and the sixth can get any amount?

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 20, \quad y_i \geq 0$$

20 “balls”, 6 “colours”

$$C(20 + 6 - 1, 6 - 1) = C(26, 5) = \frac{26!}{5!21!} = \frac{26 \cdot 25 \cdot 24 \cdot 23 \cdot 22}{5!}$$

ways of distributing the gold with constraints above

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. Each student should receive exactly one assignment. Assignments for different students should be different. How many ways are there to do it?

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. Each student should receive exactly one assignment. Assignments for different students should be different. How many ways are there to do it?

Different assignments for different students →

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. Each student should receive exactly one assignment. Assignments for different students should be different. How many ways are there to do it?

Different assignments for different students →
order matters, no repetitions

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. Each student should receive exactly one assignment. Assignments for different students should be different. How many ways are there to do it?

Different assignments for different students →
order matters, no repetitions

$$9 \cdot 8 \cdot 7 \cdot 6 = \frac{9!}{(9 - 4)!}$$

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

Different assignments for different students →

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

Different assignments for different students →
order matters.

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

Different assignments for different students →
order matters.

Let's look at this problem from the position of the assignments:

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

Different assignments for different students →
order matters.

Let's look at this problem from the position of the assignments:

$$4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 = 4^9$$

ways of distributing different assignments between different students

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

What would make it a “combinations with repetitions” problem?

STUDENTS AND ASSIGNMENTS

- There are 4 students and 9 different assignments. We need to distribute all the assignments among the students. Every student can receive an arbitrary number of assignments from 0 to 9, no assignment should be assigned to two people. How many ways are there to do it?

What would make it a “combinations with repetitions” problem?

- indistinguishable students (no ordering)
- indistinguishable assignments (repetitions)

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →
repetitions allowed, order doesn't matter →

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →
repetitions allowed, order doesn't matter →
combinations with repetitions

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →
repetitions allowed, order doesn't matter →
combinations with repetitions

15 candies = 15 balls, 7 kids = 7 colours, categories

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →
repetitions allowed, order doesn't matter →
combinations with repetitions

15 candies = 15 balls, 7 kids = 7 colours, categories

$$C(15 + 7 - 1, 7 - 1) =$$

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids?

Identical candies →
repetitions allowed, order doesn't matter →
combinations with repetitions

15 candies = 15 balls, 7 kids = 7 colours, categories

$$C(15 + 7 - 1, 7 - 1) = C(21, 6) = \frac{21!}{6! 15!} = 54264$$

ways of distributing the candies among the kids with no constraints

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

Let's give one candy to every kid to satisfy the condition.

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

Let's give one candy to every kid to satisfy the condition.
Now we need to distribute the rest:

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

Let's give one candy to every kid to satisfy the condition.
Now we need to distribute the rest:

$15 - 7 = 8$ candies = 8 balls, 7 kids = 7 colours, categories

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

Let's give one candy to every kid to satisfy the condition.
Now we need to distribute the rest:

$15 - 7 = 8$ candies = 8 balls, 7 kids = 7 colours, categories

$$C(8 + 7 - 1, 7 - 1)$$

CANDIES

- There are 15 identical candies. In how many ways can you distribute them among 7 kids so that each kid gets at least one candy?

Let's give one candy to every kid to satisfy the condition.
Now we need to distribute the rest:

$15 - 7 = 8$ candies = 8 balls, 7 kids = 7 colours, categories

$$C(8 + 7 - 1, 7 - 1) = C(14, 6) = \frac{14!}{6!8!} = 3003$$

ways of distributing the candies among the kid so that everyone gets
at least one candy

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter → combinations

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter → combinations

$$C(12,2) \cdot C(10,2) \cdot C(8,2) \cdot C(6,2) \cdot C(4,2) \cdot C(2,2) = \frac{12!}{(2!)^6}$$

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter → combinations

$$C(12,2) \cdot C(10,2) \cdot C(8,2) \cdot C(6,2) \cdot C(4,2) \cdot C(2,2) = \frac{12!}{(2!)^6}$$

But the groups are unordered!

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter → combinations

$$C(12,2) \cdot C(10,2) \cdot C(8,2) \cdot C(6,2) \cdot C(4,2) \cdot C(2,2) = \frac{12!}{(2!)^6}$$

But the groups are unordered!
We counted every split $6!$ times, fixing it:

WORKING GROUPS

- There are 12 students in the class. How many ways are there to split them into working groups of size 2 to work on the same assignment?

Order within groups doesn't matter → combinations

$$C(12,2) \cdot C(10,2) \cdot C(8,2) \cdot C(6,2) \cdot C(4,2) \cdot C(2,2) = \frac{12!}{(2!)^6}$$

But the groups are unordered!
We counted every split $6!$ times, fixing it:


$$\frac{12!}{6!(2!)^6} = 10395 \text{ possible splits}$$

THE PIGEONHOLE PRINCIPLE



THE PIGEONHOLE PRINCIPLE

If n or more pigeons are distributed among k pigeonholes and $n > k$, then at least one pigeonhole contains two or more pigeons.



BIRTHDAYS 1

- What should be the minimal number of people in the class so that we can *guarantee* that at least two students were born in the same month?

BIRTHDAYS 1

- What should be the minimal number of people in the class so that we can *guarantee* that at least two students were born in the same month?

Students = pigeons, months = holes

BIRTHDAYS 1

- What should be the minimal number of people in the class so that we can *guarantee* that at least two students were born in the same month?

Students = pigeons, months = holes

There are $k = 12$ months.

BIRTHDAYS 1

- What should be the minimal number of people in the class so that we can *guarantee* that at least two students were born in the same month?

Students = pigeons, months = holes

There are $k = 12$ months.

If there are $n > k$ students,
at least two of them are born in the same month.

BIRTHDAYS 1

- What should be the minimal number of people in the class so that we can *guarantee* that at least two students were born in the same month?

Students = pigeons, months = holes

There are $k = 12$ months.

If there are $n > k$ students,
at least two of them are born in the same month.

Therefore, there must be at least **13** students.

BIRTHDAYS 2

- How many people there should be in the room to guarantee that two people have birthday on the same day?

BIRTHDAYS 2

- How many people there should be in the room to guarantee that two people have birthday on the same day?

People = pigeons, days = holes

BIRTHDAYS 2

- How many people there should be in the room to guarantee that two people have birthday on the same day?

People = pigeons, days = holes

There are $k = 366$ days.

BIRTHDAYS 2

- How many people there should be in the room to guarantee that two people have birthday on the same day?

People = pigeons, days = holes

There are $k = 366$ days.

If there are $n > k$ people,
at least two of them are born in the same month.

BIRTHDAYS 2

- How many people there should be in the room to guarantee that two people have birthday on the same day?

People = pigeons, days = holes

There are $k = 366$ days.

If there are $n > k$ people,
at least two of them are born in the same month.

Therefore, there must be at least **367** people.

NUMBERS

- Prove that if seven distinct numbers are selected from $\{1, 2, \dots, 11\}$, then some two of these numbers sum to 12.

NUMBERS

- Prove that if seven distinct numbers are selected from $\{1, 2, \dots, 11\}$, then some two of these numbers sum to 12.

We can split the numbers into the following sets:

$$\{1, 11\}, \{2, 10\}, \{3, 9\}, \{4, 8\}, \{5, 7\}, \{6\}$$

NUMBERS

- Prove that if seven distinct numbers are selected from $\{1, 2, \dots, 11\}$, then some two of these numbers sum to 12.

We can split the numbers into the following sets:

$$\{1, 11\}, \{2, 10\}, \{3, 9\}, \{4, 8\}, \{5, 7\}, \{6\}$$

Those are our $k = 6$ holes.

NUMBERS

- Prove that if seven distinct numbers are selected from $\{1, 2, \dots, 11\}$, then some two of these numbers sum to 12.

We can split the numbers into the following sets:

$$\{1, 11\}, \{2, 10\}, \{3, 9\}, \{4, 8\}, \{5, 7\}, \{6\}$$

Those are our $k = 6$ holes.

$n = 7 > k$ numbers are selected.

NUMBERS

- Prove that if seven distinct numbers are selected from $\{1, 2, \dots, 11\}$, then some two of these numbers sum to 12.

We can split the numbers into the following sets:

$$\{1, 11\}, \{2, 10\}, \{3, 9\}, \{4, 8\}, \{5, 7\}, \{6\}$$


Those are our $k = 6$ holes.

$n = 7 > k$ numbers are selected.

At least two of them will belong to the same set, meaning that at least two of them will sum up to 12.

GENERALIZED PIGEONHOLE PRINCIPLE


- Given n items that fall into k different categories, if $n > m \cdot k$ for some positive integer m , then at least $m + 1$ of the items must fall into the same category.



BREAKOUT ROOMS

- Given n items that fall into k different categories, if $n > m \cdot k$ for some positive integer m , then at least $m + 1$ of the items must fall into the same category.

Example:




There are $n = 5$ people in this class.

$k = 2$ breakout rooms are created.

BREAKOUT ROOMS

- Given n items that fall into k different categories, if $n > m \cdot k$ for some positive integer m , then at least $m + 1$ of the items must fall into the same category.

Example:



There are $n = 5$ people in this class.


$k = 2$ breakout rooms are created.

$$n > 2 \cdot k = 4 \Rightarrow$$

BREAKOUT ROOMS

- Given n items that fall into k different categories, if $n > m \cdot k$ for some positive integer m , then at least $m + 1$ of the items must fall into the same category.

Example:



There are $n = 5$ people in this class.

$k = 2$ breakout rooms are created.

$$n > 2 \cdot k = 4 \Rightarrow$$

at least $2 + 1 = 3$ students will be put in the same breakout room.

TEST

- 30 students participated in an exam. The worst student in class got 13 answers wrong, while others made fewer mistakes. Show that there are at least 3 students who all made the same number of mistakes.

TEST

- 30 students participated in an exam. The worst student in class got 13 answers wrong, while others made fewer mistakes. Show that there are at least 3 students who all made the same number of mistakes.

Students = pigeons, number of mistakes = pigeonholes

TEST

- 30 students participated in an exam. The worst student in class got 13 answers wrong, while others made fewer mistakes. Show that there are at least 3 students who all made the same number of mistakes.

Students = pigeons, number of mistakes = pigeonholes

$$30 > 2 \cdot 13$$

TEST

- 30 students participated in an exam. The worst student in class got 13 answers wrong, while others made fewer mistakes. Show that there are at least 3 students who all made the same number of mistakes.

Students = pigeons, number of mistakes = pigeonholes

$$30 > 2 \cdot 13$$

According to the generalized pigeonhole principle, at least $2 + 1 = 3$ students must have received the same score.

FLOOR AND CEILING

- Useful functions:
 - Floor $\lfloor x \rfloor$ – the largest integer smaller than x .

FLOOR AND CEILING

- Useful functions:
 - Floor $\lfloor x \rfloor$ — the largest integer smaller than x .
 - Ceiling $\lceil x \rceil$ — the smallest integer larger than x .

FLOOR AND CEILING

- Useful functions:
 - Floor $\lfloor x \rfloor$ – the largest integer smaller than x .


$$\lfloor 5 \rfloor = 5, \quad \lfloor 2.5 \rfloor = 2, \quad \lfloor -3.3 \rfloor = -4$$

- Ceiling $\lceil x \rceil$ – the smallest integer larger than x .

$$\lceil 5 \rceil = 5, \quad \lceil 2.5 \rceil = 3, \quad \lceil -3.3 \rceil = -3$$

GENERALIZED PIGEONHOLE PRINCIPLE

- Given n items that fall into k different categories, then at least $\left\lceil \frac{n}{k} \right\rceil$ of the items must fall into the same category.



CONFERENCE ROOMS

- A conference room contains 8 tables and 107 chairs. What is the smallest possible number of chairs at the table having the most chairs?

CONFERENCE ROOMS

- A conference room contains 8 tables and 107 chairs. What is the smallest possible number of chairs at the table having the most chairs?

Pigeons: 107 chairs

Pigeonholes: 8 tables

CONFERENCE ROOMS

- A conference room contains 8 tables and 107 chairs. What is the smallest possible number of chairs at the table having the most chairs?

Pigeons: 107 chairs

Pigeonholes: 8 tables

$$\left\lceil \frac{107}{8} \right\rceil = 14$$

CONFERENCE ROOMS

- A conference room contains 8 tables and 107 chairs. What is the smallest possible number of chairs at the table having the most chairs?

Pigeons: 107 chairs

Pigeonholes: 8 tables

$$\left\lceil \frac{107}{8} \right\rceil = 14$$

Therefore, according to the generalized pigeonhole principle, at least 14 chairs are at the same table.

ODD OR EVEN

- Show that in any finite set of n numbers, at least half of the numbers are odd or at least half of them are even.

ODD OR EVEN

- Show that in any finite set of n numbers, at least half of the numbers are odd or at least half of them are even.

There are $k = 2$ “holes”: even and odd numbers.

ODD OR EVEN

- Show that in any finite set of n numbers, at least half of the numbers are odd or at least half of them are even.

There are $k = 2$ “holes”: even and odd numbers.

If n numbers are distributed between $k = 2$ holes, at least $\left\lceil \frac{n}{2} \right\rceil$ of them belong to the same hole \Leftrightarrow

ODD OR EVEN

- Show that in any finite set of n numbers, at least half of the numbers are odd or at least half of them are even.

There are $k = 2$ “holes”: even and odd numbers.

If n numbers are distributed between $k = 2$ holes, at least $\left\lceil \frac{n}{2} \right\rceil$ of them belong to the same hole \Leftrightarrow

At least $\left\lceil \frac{n}{2} \right\rceil$ are odd or $\left\lceil \frac{n}{2} \right\rceil$ are even.

PROBLEM SET 5

<https://docs.google.com/document/d/1Kf4NYeMPyABbbiVnm5sYKGgZOr3oK3XDznbb1Sf0oyY/edit?usp=sharing>

LOGISTICS INTERIM EXAM

- Monday, March 22
 - 09:00 – 10:00 (roughly): review
 - 10:20 – 12:20: exam
- Topics included:
 - Basic counting
 - Inclusion-exclusion
 - Permutations and combinations
 - Pigeonhole principle