

# INTRODUCTION TO STATISTICS

## LECTURE 8

# TODAY

- Confidence intervals based on normal data
  - for  $\mu$  when  $\sigma$  is known;
  - for  $\mu$  when  $\sigma$  is unknown;
  - for  $\sigma$ .

# TODAY

- Confidence intervals based on normal data
  - for  $\mu$  when  $\sigma$  is known;
  - for  $\mu$  when  $\sigma$  is unknown;
  - for  $\sigma$ .
- Large sample CI

# TODAY

- Confidence intervals based on normal data
  - for  $\mu$  when  $\sigma$  is known;
  - for  $\mu$  when  $\sigma$  is unknown;
  - for  $\sigma$ .
- Large sample CI
- Bernoulli data and polling

# A QUICK REMINDER

What we saw last time

# CI: DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (T_1, T_2)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

# CI: DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (T_1, T_2)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals:  $T_1$  and  $T_2$  are functions of random samples.

# CI: DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (T_1, T_2)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals:  $T_1$  and  $T_2$  are functions of random samples.
- $\theta$  is unknown, but fixed  
 $T_1$  and  $T_2$  are random



# CI: INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

# CI: INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.

# CI: INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.
- Common interpretation:

# CI: INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.
- Common interpretation:

*If I repeat the experiment many times, the interval will contain the true value of  $\theta$  95% of the time ( $\alpha=0.05$ ).*

# CI FOR NORMAL DATA

CI for  $\mu$ , *known*  $\sigma$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.  
How to construct a CI for  $\mu$ ?

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.  
How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.  
How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim$$



# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.  
How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N( \quad ) \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.  
How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \quad) \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0,1)$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

# CI FOR $\mu$ , KNOWN $\sigma$


$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :


$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left( \quad < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < \quad \right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$




# CI FOR $\mu$ , KNOWN $\sigma$







# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

$$P\left(\bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .
- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05,$$



# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} =$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .
- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} =$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} = 1.96$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} = 1.96$$

$$12 \pm \frac{3}{\sqrt{100}} \cdot 1.96$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.90 \Rightarrow \alpha = 0.1, \quad z_{1-0.05} = 1.64$$



# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.90 \Rightarrow \alpha = 0.1, \quad z_{1-0.05} = 1.64$$

$$5 \pm \frac{1}{\sqrt{100}} \cdot 1.64$$

# PRACTICE!

Google Classroom -> Lecture 8 -> z-intervals

# CI FOR NORMAL DATA

CI for  $\mu$ , *unknown*  $\sigma$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ?

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$



# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s}$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n-1) \text{ — Student distribution}$$

# STUDENT DISTRIBUTION



William Sealy Gosset

# STUDENT DISTRIBUTION


$$\frac{(\bar{X}_n - \mu)\sqrt{n}}{s} \sim t(n - 1)$$



William Sealy Gosset

# STUDENT DISTRIBUTION

$$\frac{(\bar{X}_n - \mu)\sqrt{n}}{s} \sim t(n - 1)$$



William Sealy Gosset

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

Before: 
$$P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$



# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

Before: 
$$P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now:

$$P\left(-t_{n-1, 1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{n-1, 1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

Before: 
$$P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(-t_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

Before: 
$$P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(-t_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

$$\text{Before: } P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(\bar{X} - \frac{s}{\sqrt{n}} t_{1-\alpha/2} < \mu < \bar{X} + \frac{s}{\sqrt{n}} t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05,$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} =$$



# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} = 2.093$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} = 2.093$$

$$\mu = 42 \pm \frac{6}{\sqrt{20}} \cdot 2.093$$

# PRACTICE!

Google Classroom -> Lecture 8 -> t-intervals

# CI FOR NORMAL DATA

CI for  $\sigma$ , unknown  $\mu$ ,  $\sigma$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ?

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\mu$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$\frac{(n-1)s^2}{\sigma^2}$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\mu$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1) -$$

Chi-squared distribution

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i.i.d. } N(0,1)$$

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i. i. d. } N(0,1)$$

$\Rightarrow$


$$Q = Z_1^2 + \dots + Z_n^2 \sim \chi^2(n - 1)$$

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i.i.d. } N(0,1)$$

$\Rightarrow$

$$Q = Z_1^2 + \dots + Z_n^2 \sim \chi^2(n-1)$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :


$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left( < \frac{(n-1)s^2}{\sigma^2} < \right) = 1 - \alpha$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

Chi Square Distribution: df = 8

$P(X < 3.49) = 10\%$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left(c_{\alpha/2} < \frac{(n-1)s^2}{\sigma^2} < c_{1-\alpha/2}\right) = 1 - \alpha$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\mu$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left(c_{\alpha/2} < \frac{(n-1)s^2}{\sigma^2} < c_{1-\alpha/2}\right) = 1 - \alpha$$

$$P\left(\frac{(n-1)s^2}{c_{1-\alpha/2}} < \sigma^2 < \frac{(n-1)s^2}{c_{\alpha/2}}\right) = 1 - \alpha$$

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

$$\left[ \frac{(n-1)s^2}{c_{1-\alpha/2}}; \frac{(n-1)s^2}{c_{\alpha/2}} \right]$$

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42$ ,  $s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

$$\left[ \frac{(n-1)s^2}{c_{1-\alpha/2}}; \frac{(n-1)s^2}{c_{\alpha/2}} \right]$$

$$\left[ \frac{19 \cdot 36}{32.85}; \frac{19 \cdot 36}{8.91} \right]$$

# LARGE SAMPLES

Using Central Limit Theorem

# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1)$$

# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1) \Rightarrow \mu \approx$$



# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1) \Rightarrow \mu \approx \bar{X} \pm \frac{s}{\sqrt{n}} z_{1-\alpha/2}$$

# PRACTICE!

Google Classroom -> Lecture 8 -> Large sample CI