



INTRODUCTION TO STATISTICS

LECTURE 10


MID-TERM RESULTS: AN OVERVIEW




MID-TERM RESULTS: AN OVERVIEW



MID-TERM RESULTS: AN OVERVIEW




MID-TERM RESULTS: AN OVERVIEW




- MIN and MAX scores:
 - 5.25 and 40
- Mean score:
 - 23 points

MID-TERM RESULTS: AN OVERVIEW



- MIN and MAX scores:
 - 5.25 and 40
- Mean score:
 - 23 points
- Median score:
 - 23.1 points

MID-TERM RESULTS: AN OVERVIEW



- MIN and MAX scores:
 - 5.25 and 40
- Mean score:
 - 23 points
- Median score:
 - 23.1 points
- Standard deviation:
 - 10.5 points

PLANS FOR THE WEEK

- MONDAY
 - CI recap
 - Hypothesis testing I
- TUESDAY
 - Hypothesis testing II
- WEDNESDAY
 - Covariance and correlation
 - Linear regression
- THURSDAY
 - Big recap
- FRIDAY
 - Final exam

PLANS FOR THE WEEK

- MONDAY

- CI recap
- Hypothesis testing I

- Assignment 4, part I is due 23:59
- Assignment 4, part II is out

- TUESDAY

- Hypothesis testing II

- WEDNESDAY

- Covariance and correlation
- Linear regression

- Assignment 4, part II is due 23:59
- Assignment 5 is out

- THURSDAY

- Big recap

- FRIDAY

- Final exam

- Assignment 5 is due 23:59

LAST TIME

- Confidence intervals:
 - z-interval;
 - t-interval;
 - χ^2 -interval;
 - some examples.
- Hypothesis testing
 - *The Lady Tasting Tea experiment;*
 - *an introduction to hypothesis testing.*

LAST TIME

- Confidence intervals:
 - z-interval;
 - t-interval;
 - χ^2 -interval;
 - some examples.


CONFIDENCE INTERVALS

A BRIEF RECAP


CI: A BRIEF RECAP

Is your data
normally
distributed?


CI: A BRIEF RECAP




CI: A BRIEF RECAP




CI: A BRIEF RECAP




CI: A BRIEF RECAP




CI: A BRIEF RECAP




CI: A BRIEF RECAP



CI: A BRIEF RECAP



CI: A BRIEF RECAP



QUICK QUIZ

CI: A BRIEF RECAP

A machine fills in wine bottles with a random amount of wine that follows normal distribution with unknown mean μ and unknown standard deviation σ .

You check the last 100 bottles and see that on average they were filled with $\bar{X} = 705$ ml of wine, with sample std $s = 3$ ml.

**Which interval would you use
to construct a 95%-CI for μ ?**

CI: A BRIEF RECAP

In a study on cholesterol levels a sample of 1000 patients was chosen.

The average plasma cholesterol levels subjects was $\bar{X} = 6$ mmol/L, with sample std $s = 0.4$ mmol/L.

**Which interval would you use
to construct a 95%-CI for the true mean?**

CI: A BRIEF RECAP

Height of a female student is a random variable following normal distribution with unknown mean μ and standard deviation $\sigma = 5$ cm.

The average height of 100 female students $\bar{X} = 165$ cm, and you sample std $s = 4$ cm

**Which interval would you use
to construct a 95%-CI for μ ?**

CI: A BRIEF RECAP

On a candy factory, a machine fills packs with random number of candies which follows normal distribution with unknown mean μ and unknown standard deviation σ .

In the last 50 packs, there were on average $\bar{X} = 90$ g of sweets, with sample std $s = 7$ g.

**Which interval would you use
to construct a 95%-CI for σ ?**

HYPOTHESIS TESTING

HYPOTHESIS TESTING

- Test whether a default assumption about the data is plausible.
- Key idea: you can't *prove* something, but you can disprove it.

HYPOTHESIS TESTING

- Test whether a default assumption about the data is plausible.

HYPOTHESIS TESTING

**ASSUMPTION:
ALL THE SWANS ARE WHITE**



HYPOTHESIS TESTING

**ASSUMPTION:
ALL THE SWANS ARE WHITE**

EVIDENCE:



HYPOTHESIS TESTING

**ASSUMPTION:
ALL THE SWANS ARE WHITE**



**EVIDENCE:
NOT CONVINCING...**



HYPOTHESIS TESTING

**ASSUMPTION:
ALL THE SWANS ARE WHITE**



**EVIDENCE:
NOW IT'S CONVINCING!**



HYPOTHESIS TESTING

ASSUMPTION:
ALL THE SWANS ARE WHITE



EVIDENCE:



HYPOTHESIS TESTING

**ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON**



HYPOTHESIS TESTING

**ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON**

EVIDENCE:




HYPOTHESIS TESTING

ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON



EVIDENCE:
STILL NO REASON TO BELIEVE THERE ARE
ALIENS




HYPOTHESIS TESTING

**ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON**



EVIDENCE:

STILL NO REASON TO BELIEVE THERE ARE
ALIENS




HYPOTHESIS TESTING

ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON



EVIDENCE:

THERE ARE ALIENS!!!




HYPOTHESIS TESTING

ASSUMPTION:
THERE'RE NO ALIENS ON THE MOON



EVIDENCE:

THERE ARE ALIENS!!!



HYPOTHESIS TESTING

- Note:
 - Default assumption = “status quo”.

HYPOTHESIS TESTING

- Note:
 - Default assumption = “status quo”.
 - Alternative: something we can find evidence for.

HOW TO TEST A HYPOTHESIS

BASIC INGREDIENTS

RUNNING EXAMPLE

How to check if a coin is a fair one?

RUNNING EXAMPLE

We flip a coin 10 times to test if it's a fair one.

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

$$H_1: p \neq 0.5 \text{ 'the coin is not fair'}$$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

$$H_1: p \neq 0.5 \text{ 'the coin is not fair'}$$

$$H_1: p > 0.5$$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

$$H_1: p \neq 0.5 \text{ 'the coin is not fair'}$$

$$H_1: p > 0.5 \text{ or } H_1: p < 0.5$$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

Two-sided: $H_1: p \neq 0.5$ 'the coin is not fair'

One-sided: $H_1: p > 0.5$ or $H_1: p < 0.5$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

- **The null hypothesis H_0** : *the default assumption for the model generating the data.*

$$H_0: p = 0.5 \text{ 'the coin is fair'}$$

- **The alternative hypothesis H_1** : *if we reject H_0 , we accept H_1 as the best explanation for the data.*

Two-sided: $H_1: p \neq 0.5$ 'the coin is not fair'

One-sided: $H_1: p > 0.5$ or $H_1: p < 0.5$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X — *number of H*

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X — *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X — number of H

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X — number of H

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(\quad)$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X — *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X – *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

X											
$P(X H_0)$											

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X – *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$											

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X – *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001										0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X – *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01								0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

- **Test statistic:** *computed from the data*

X – *number of H*

- **Null distribution:** *the probability distribution of X assuming H_0*

$X \sim \text{Binomial}(10, 0.5)$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- **Rejection region R:** if X is in the rejection region, we reject H_0 in favor of H_1 .

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.


$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- **Rejection region R:** if X is in the rejection region, we reject H_0 in favor of H_1 .
- **Significance level α :** $P(X \in R|H_0) \leq \alpha$
 - Typically chosen in advance (common values are 0.1, 0.05, 0.01.)

INGREDIENTS

- Distribution of the test statistic under H_0 , two-sided alternative:



INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

$$\alpha = 0.2$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

$$\alpha = 0.2$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

REMARKS

- **The null hypothesis** is the *cautious default*
 - we won't claim the coin is unfair unless we have compelling evidence.

REMARKS

- **The null hypothesis** is the *cautious default*
 - we won't claim the coin is unfair unless we have compelling evidence.
- **Rejection region:** data that is *extreme* under the null hypothesis = outcomes in the tail(s) of the null distribution.
 - depends on the significance level α of the test.

REMARKS

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair' $\alpha = 0.05$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

REMARKS

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair' $\alpha = 0.05$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- If we get 0, 1, 9, 10 H
 - Reject H_0

REMARKS

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair' $\alpha = 0.05$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- If we get 0, 1, 9, 10 H
 - Reject H_0
- If we get 2, 3, 4, 5, 6, 7, 8 H, then the test statistic is in the non-rejection region.

REMARKS

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair' $\alpha = 0.05$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- If we get 0, 1, 9, 10 H
 - Reject H_0
- If we get 2, 3, 4, 5, 6, 7, 8 H, then the test statistic is in the non-rejection region.
 - Interpretation: the data 'does not support rejecting the null hypothesis'.

REMARKS

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair' $\alpha = 0.05$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

- If we get 0, 1, 9, 10 H
 - Reject H_0
- If we get 2, 3, 4, 5, 6, 7, 8 H, then the test statistic is in the non-rejection region.
 - Interpretation: the data 'does not support rejecting the null hypothesis'.
 - **Never claim that the data proves the null hypothesis is true.**

EXPERIMENT

EXPERIMENT

- Toss a coin 10 times, record the number of heads you got.

EXPERIMENT

- Toss a coin 10 times, record the number of heads you got.
- Based on the result, do you reject the null hypothesis?

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair', $\alpha = 0.2$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INCORRECTLY REJECTING H_0

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

What is the probability to incorrectly reject H_0 ?

INCORRECTLY REJECTING H_0

$H_0: p = 0.5$ 'the coin is fair', $H_1: p \neq 0.5$ 'the coin is not fair'

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

What is the probability to incorrectly reject H_0 ?

$$P(\text{Reject } H_0 | H_0) = P(X \in R | H_0) = \alpha - \text{significance level}$$

ONE-SIDED ALTERNATIVES

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p > 0.5$ 'the biased towards H'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p > 0.5$ 'the biased towards H'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$, $H_1: p > 0.5$ 'the biased towards H'

$$\alpha = 0.1$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5$, $H_1: p > 0.5$ 'the biased towards H'

$$\alpha = 0.1$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p < 0.5$ 'the biased towards T'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p < 0.5$ 'the biased towards T'

$$\alpha = 0.05$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p < 0.5$ 'the biased towards T'

$$\alpha = 0.1$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

INGREDIENTS

We flip a coin 10 times to test if it's a fair one.

$H_0: p = 0.5,$ $H_1: p < 0.5$ 'the biased towards T'

$$\alpha = 0.1$$

X	0	1	2	3	4	5	6	7	8	9	10
$P(X H_0)$	0.001	0.01	0.044	0.117	0.205	0.246	0.205	0.117	0.044	0.01	0.001

Z-TEST

When the data is coming from normal distribution
and variance is known

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- We suspect that HarbourSpace students are different. How do we test it?

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- We suspect that HarbourSpace students are different. How do we test it?

$$H_0: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} = \mu_0$$

‘HarbourSpace students’ IQs are the same as those of the general population’

$$H_1: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} > \mu_0$$

‘HarbourSpace students’ IQs are higher than those of the general population’

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- We suspect that HarbourSpace students are different. How do we test it?

$$H_0: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} = \mu_0$$

‘HarbourSpace students’ IQs are the same as those of the general population’

$$H_1: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} \neq \mu_0$$

‘HarbourSpace students’ IQs aren’t the same as than those of the general population’

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- We suspect that HarbourSpace students are different. How do we test it?

$$H_0: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} = \mu_0$$

‘HarbourSpace students’ IQs are the same as those of the general population’

$$H_1: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} < \mu_0$$

‘HarbourSpace students’ IQs are lower than those of the general population’

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- We suspect that HarbourSpace students are different. How do we test it?

$$H_0: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} = \mu_0$$

'HarbourSpace students' IQs are the same as those of the general population'

$$H_1: X \sim N(\mu_{HS}, \sigma^2), \quad \mu_{HS} \neq \mu_0$$

'HarbourSpace students' IQs aren't the same as than those of the general population'

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Average IQ of n students: \bar{X}

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Average IQ of n students: \bar{X}
- Significance level α

z-test

- IQ follows $N(\mu_0, \sigma^2)$ where σ is known.
- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Average IQ of n students: \bar{X}
- Significance level α
- Can we reject H_0 ?

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim N(\mu_0, \sigma^2)$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim N(\mu_0, \sigma^2)$

\Rightarrow

$\sim N(0, 1)$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim N(\mu_0, \sigma^2)$

$$\Rightarrow \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim N(\mu_0, \sigma^2)$

$$\Rightarrow T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$

- Test statistic:

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Assuming H_0 , $X_i \sim N(\mu_0, \sigma^2)$


$$\Rightarrow T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$$

z-test

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0 \quad H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:

z-test


- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu \neq \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:



ONE-SIDED ALTERNATIVES


z-test: ONE-SIDED

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu > \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:




z-test: ONE-SIDED

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu > \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:




z-test: ONE-SIDED

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu < \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:



z-test: ONE-SIDED

- $H_0: X \sim N(\mu_{HS}, \sigma^2), \mu = \mu_0$ $H_1: X \sim N(\mu_{HS}, \sigma^2), \mu < \mu_0$
- Test statistic: $T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{\sigma} \sim N(0, 1)$
- Significance level α
- (non-) Rejection region:



Z-TEST: EXAMPLE 1

- Weight loss is described by $X \sim N(5, 7^2)$.
- $n = 30$ patients followed an experimental diet.
- Average weight loss $\bar{X} = 6.1$ kg.

Does the diet make any difference?

Z-TEST: EXAMPLE 2

- A bakery supplies loaves of bread to supermarkets. Not every loaf weighs the same: loaf weight $X \sim N(\mu_0, 0.1^2)$, supermarket expects $\mu_0 = 2$ kg.
- A supermarket draws a sample of $n = 20$ loaves: average weights is $\bar{X} = 1.97$ kg.
- The supermarket wants to be sure that the weights are, on average, not lower than 2 kg.

Is there evidence against this?

T-TEST

When the data is coming from normal distribution
and variance is unknown

T-TEST

- Hypothesis about μ from $X \sim N(\mu_0, \sigma^2)$, σ is unknown.

T-TEST

- Hypothesis about μ from $X \sim N(\mu_0, \sigma^2)$, σ is unknown.
- Unknown variance is estimated from the sample:

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

T-TEST

- Hypothesis about μ from $X \sim N(\mu_0, \sigma^2)$, σ is unknown.
- Unknown variance is estimated from the sample:

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- The test statistic is therefore:

$$T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{s} \sim$$

T-TEST

- Hypothesis about μ from $X \sim N(\mu_0, \sigma^2)$, σ is unknown.
- Unknown variance is estimated from the sample:

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- The test statistic is therefore:

$$T(X_1, \dots, X_n) = \frac{(\bar{X} - \mu_0)\sqrt{n}}{s} \sim t(n-1)$$

T-TEST: EXAMPLE 1

- A bakery supplies loaves of bread to supermarkets. Not every loaf weighs the same: loaf weight $X \sim N(\mu_0, \sigma^2)$, supermarket expects $\mu_0 = 2$ kg.
- A supermarket draws a sample of $n = 20$ loaves: average weights is $\bar{X} = 1.9668$ kg, $s^2 = 0.0927^2$
- The supermarket wants to be sure that the weights are, on average, not lower than 2 kg.

Is there evidence against this?

T-TEST: EXAMPLE 2

- Sales at your company: $X \sim N(\mu_0, \sigma^2)$, $\mu_0 = 100$ dollars per transaction.
- You hire new employees. Based of a sample of $n = 25$ salesmen, average sale is $\bar{X} = 130$ dollars, with sample standard deviation $s = 15$.

Are the new employees better than the old ones?
Test your hypothesis at $\alpha = 0.05$.

TO SUM UP

- Hypothesis testing ingredients
 - H_0 and H_1
 - Test statistic
 - Null distribution
 - Rejection region
 - Two- and one-sided alternatives

TO SUM UP

- Hypothesis testing ingredients
 - H_0 and H_1
 - Test statistic
 - Null distribution
 - Rejection region
 - Two- and one-sided alternatives
- One-sample tests:
 - Z-test: normal distribution, variance known
 - T-test: normal distribution, variance unknown