

# INTRODUCTION TO STATISTICS

## LECTURE 2

# LAST TIME

- DESCRIPTIVE STATISTICS
  - Data about **the whole population** is available.
  - Summarize it with
    - *summary statistics*;
    - *tables*;
    - *plots*.

# LAST TIME

- DESCRIPTIVE STATISTICS

- Data about **the whole population** is available.
- Summarize it with
  - *summary statistics*;
  - *tables*;
  - *plots*.

- INFERENCE STATISTICS

- Data about **a sample of the whole population** is available.
- Reason about the whole population.

# LAST TIME

- DESCRIPTIVE STATISTICS

- Data about **the whole population** is available.
- Summarize it with
  - *summary statistics;*
  - *tables;*
  - *plots.*

- INFERENCE STATISTICS

- Data about **a sample of the whole population** is available.
- Reason about the whole population.



# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*
- **How many men and women are there in the world?**

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*
- **How many men and women are there in the world?**

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands

<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>



# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?


<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands

<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>



**p = 0.5**

# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>


- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands

<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>



**p = 0.5**

# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?


$$N_{\text{men}} = X_1 + \dots + X_{100} \sim \text{Bi}(100, p)$$

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands

<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>



$$p = 0.5$$

# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?

$$N_{\text{men}} = X_1 + \dots + X_{100} \sim \text{Bi}(100, \mathbf{p})$$


$$E(N_{\text{men}}) = 100 * \mathbf{p} = 50$$

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands

<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>



$$\mathbf{p} = 0.5$$

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$

$$P(X = 1) = \mathbf{p} = ?$$

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*
- **How many vegetarians are there in the world?**

$X \sim \text{Bernoulli}(\mathbf{p})$

$P(X = 1) = \mathbf{p} = ?$

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*
- **How many vegetarians are there in the world?**
- **No precise data!** 😞

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

*What would you do?*



# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

- **How many vegetarians are there in the world?**

- **No precise data!** 😞

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

- *What is a good guess for the value of  $\mathbf{p}$ ?*

- **How many vegetarians are there in the world?**

- **No precise data!** 😞

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

- *What is a good guess for the value of  $\mathbf{p}$ ?*

$$\hat{\mathbf{p}} = 13 / 100 = 0.13$$

- **How many vegetarians are there in the world?**

- **No precise data!** 😞

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

## INFERENCE STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?
- How many vegetarians are there in the world?
- No precise data! ☹️

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

What would you do?

- What is a good guess for the value of  $\mathbf{p}$ ?

$$\hat{\mathbf{p}} = 13 / 100 = 0.13$$

...get the parameter estimate

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?

## (NON-)VEGETARIANS

### INFERENCE STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?
- **How many vegetarians are there in the world?**

$$X \sim \text{Bernoulli}(p)$$
$$P(X = 1) = p = ?$$

- No precise data! ☹️

What would you do?

- What is a good guess for the value of  $p$ ?

$$\hat{p} = 13 / 100 = 0.13$$

- Survey:

- 100 people:  $X_1 \dots X_{100}$
- 13 are vegetarians

...get the parameter estimate

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?

## (NON-)VEGETARIANS

INFERENCE  
STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?  
 $X \sim \text{Bernoulli}(p)$   
 $P(X = 1) = p = ?$
- **How many vegetarians are there in the world?**
  - No precise data! ☹️

What would you do?
- What is a good guess for the value of  $p$ ?
  - Survey:
    - 100 people:  $X_1 \dots X_{100}$
    - 13 are vegetarians

$\hat{p} = 13 / 100 = 0.13$

...get the parameter estimate

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?
- Would it be better if there were more people in the survey?

**(NON-)VEGETARIANS**

INFERENTIAL STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?  
 $X \sim \text{Bernoulli}(p)$   
 $P(X = 1) = p = ?$
- How many vegetarians are there in the world?
  - No precise data! ☹️
  - What would you do?
- What is a good guess for the value of  $p$ ?
  - Survey:
    - 100 people:  $X_1 \dots X_{100}$
    - 13 are vegetarians

$\hat{p} = 13 / 100 = 0.13$

...get the parameter estimate

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?
- Would it be better if there were more people in the survey?

## (NON-)VEGETARIANS

• If we take a random person  $X$ , what's the probability that it's a vegetarian?

$$X \sim \text{Bernoulli}(p)$$
$$P(X = 1) = p = ?$$

• What is a good guess for the value of  $p$ ?

$\hat{p} = 13 / 100 = 0.13$

...get the parameter estimate

## INFERENCE STATISTICS

• How many vegetarians are there in the world?

• No precise data! ☹️

What would you do?

• Survey:

- 100 people:  $X_1 \dots X_{100}$
- 13 are vegetarians

From the data...



# **THE METHOD OF MAXIMUM LIKELIHOOD**

A general framework for parameter estimation

# THE METHOD OF MAXIMUM LIKELIHOOD

A general framework for parameter estimation

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\boldsymbol{p}$ ?

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\boldsymbol{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\mathbf{p})$  – whether a person is a vegetarian (1) or not (0).
- $\mathbf{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\mathbf{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid \mathbf{p}) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\mathbf{p})$  – whether a person is a vegetarian (1) or not (0).
- $\mathbf{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\mathbf{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid \mathbf{p}) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid \mathbf{p}) * \dots * P(X_{13} = 1 \mid \mathbf{p}) * P(X_{14} = 0 \mid \mathbf{p}) * \dots * P(X_{100} = 0 \mid \mathbf{p}) =$$



# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(p)$  – whether a person is a vegetarian (1) or not (0).
- $p$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $p$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid p) * \dots * P(X_{13} = 1 \mid p) * P(X_{14} = 0 \mid p) * \dots * P(X_{100} = 0 \mid p) =$$

$$= p^{13}(1-p)^{87}$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(p)$  – whether a person is a vegetarian (1) or not (0).
- $p$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $p$ ? **Likelihood:**

$$L(p) = P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid p) * \dots * P(X_{13} = 1 \mid p) * P(X_{14} = 0 \mid p) * \dots * P(X_{100} = 0 \mid p) =$$

$$= p^{13}(1-p)^{87}$$

**Likelihood function** is the joint probability of realized sample given the parameters.

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?
  - We want the data that we obtained to be the most likely one.

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?
  - We want the data that we obtained to be the most likely one.
  - **Maximize  $L(p)$  w.r.t.  $p$ !**

# **CALCULUS 101**

- **How to optimize a function?**



# CALCULUS 101

- **How to optimize a function?**
  1. Compute its derivative.
  2. Set it to zero.
  3. Get the critical point(s).
  4. Chose the point of maximum.

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1 - x} = 0$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1 - x} = 0$$

Critical points:

$$x = 0, \quad x = 1, \quad x = \frac{1}{2}$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1-x} = 0$$

Critical points:

$$x = 0, \quad x = 1, \quad x = \frac{1}{2}$$



# MAXIMUM LIKELIHOOD


$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$f(x) = x*(1-x)$$



**$\log f(x)$**



# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) =$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) = \frac{13}{p} - \frac{87}{1-p} = 0$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) = \frac{13}{p} - \frac{87}{1-p} = 0$$

$$\hat{p} = \frac{13}{100}$$

# MAXIMUM LIKELIHOOD ESTIMATE

1. Write down the likelihood function:

$$L(\theta) = P(X_1, \dots, X_n \mid \theta) = \prod_{i=1}^n P(X_i \mid \theta)$$

2. Find its maximum w.r.t. the unknown parameter  $\theta$ :

$$\hat{\theta} = \operatorname{argmax}_{\theta} L(\theta) \text{ w.r.t. } \theta$$

(!) In many cases, it's easier to maximize **log-likelihood**:

$$\log L(\theta) = \log \prod_{i=1}^n P(X_i \mid \theta) = \sum_{i=1}^n \log P(X_i \mid \theta)$$

$$\hat{\theta} = \operatorname{argmax}_{\theta} \log L(\theta)$$

**Maximum Likelihood Estimate (MLE)** is the value which

...

**Maximum Likelihood Estimate (MLE)** is the value which maximizes the probability of observing the realized sample.

**BREAK**



# **MLE FOR SOME DISCRETE DISTRIBUTIONS**

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

- *(General case of the previous example)*
- You want to estimate the proportion of the vegetarians.

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

- *(General case of the previous example)*
- You want to estimate the proportion of the vegetarians.
- Survey: you ask  $N$  people whether they are vegetarians and get responses

$$X_1, X_2, \dots, X_N$$

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

- *(General case of the previous example)*
- You want to estimate the proportion of the vegetarians.
- Survey: you ask  $N$  people whether they are vegetarians and get responses

$$X_1, X_2, \dots, X_N$$

- What's the MLE of  $p$ ?

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$



# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{1}{N} \sum_{i=1}^N X_i$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{1}{N} \sum_{i=1}^N X_i$$


# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.



# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Suppose you want to model the number of phone calls a call center receives in an hour.

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.

# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.

- Suppose you want to model the number of phone calls a call center receives in an hour.
- You've been recording it for the past  $n$  hours:

$X_1, X_2, \dots, X_n$



# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.

- Suppose you want to model the number of phone calls a call center receives in an hour.
- You've been recording it for the past  $n$  hours:

$$X_1, X_2, \dots, X_n$$

- What's the MLE of  $\lambda$ ?

# MLE FOR POISSON DISTRIBUTION

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) =$$

# MLE FOR POISSON DISTRIBUTION

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) =$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) =$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0,$$



# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0, \quad \hat{\lambda} = \frac{1}{n} \sum_{i=1}^n X_i$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} \log L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0,$$

$$\hat{\lambda} = \frac{1}{n} \sum_{i=1}^n X_i$$

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	?	?	?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	?	?	?

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	$p$	$q$	$1-p-q$

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:
- We observed it  $N$  times:

<b>X</b>	1	2	3
<b>P(X)</b>	$p$	$q$	$1-p-q$

<b>X</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:
- We observed it  $N$  times:

<b>X</b>	1	2	3
<b>P(X)</b>	$p$	$q$	$1-p-q$

<b>X</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- How many parameters does such a distribution have?
- What is the MLE of the parameters?

# **CALCULUS 101**

- **How to optimize a function of several variables?**



# CALCULUS 101

- **How to optimize a function of several variables?**

1. Compute partial derivatives.
2. Set them to zero.
3. Solve the equations and get the critical points.

# CALCULUS 101

- **How to optimize a function of several variables?**

1. Compute partial derivatives.
2. Set them to zero.

3. Solve the equations and get the critical points.

$$f(x, y) = x^2 + 2xy - 2x - 4y$$

$$\frac{d}{dx}f(x, y) = 2x + 2y - 2 = 0$$

$$\frac{d}{dy}f(x, y) = 2x - 4 = 0$$

$$x^* = 2, \quad y^* = 1$$

# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3
<b>P(X)</b>	$p$	$q$	$1-p-q$

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) =$$



# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$



# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$



$$\log L(p, q) =$$

# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$

$\Leftrightarrow$

$$\log L(p, q) = n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$$

maximize  $\log L(p, q)$  w.r.t.  $p, q$

# MLE: EXAMPLE 2

<b>X</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

maximize  $n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$  w.r.t.  $p, q$

$$\frac{d}{dp} \log L(p, q) = \frac{n_1}{p} - \frac{n_3}{1 - p - q} = 0$$

$$\frac{d}{dq} \log L(p, q) = \frac{n_2}{q} - \frac{n_3}{1 - p - q} = 0$$

# MLE: EXAMPLE 2


<b>X</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

maximize  $n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$  w.r.t.  $p, q$

$$\frac{d}{dp} \log L(p, q) = \frac{n_1}{p} - \frac{n_3}{1 - p - q} = 0$$

$$\frac{d}{dq} \log L(p, q) = \frac{n_2}{q} - \frac{n_3}{1 - p - q} = 0$$



$$\hat{p} = \frac{n_1}{N}$$
$$\hat{q} = \frac{n_2}{N}$$




# BINOMIAL DISTRIBUTION

$X \sim \text{Bi}(n, p), \quad n = 1, 2, \dots, \quad 0 < p < 1$

$$P(X = k) = C_n^k p^k (1 - p)^{n-k}$$

$$E(X) = np, \quad \text{Var}(X) = np(1-p)$$

- Models the number of successes in a series of  $n$  independent Bernoulli trials, each of which has a success probability  $p$ .



# BINOMIAL DISTRIBUTION

$$X \sim \text{Bi}(n, p), \quad n = 1, 2, \dots, \quad 0 < p < 1$$

$$P(X = k) = C_n^k p^k (1 - p)^{n-k}$$

$$E(X) = np, \quad \text{Var}(X) = np(1-p)$$

- Models the number of successes in a series of  $n$  independent Bernoulli trials, each of which has a success probability  $p$ .

- You sell two types of sandwiches: chicken and vegetarian. Which one people like more?

- For the past  $N$  days, you were selling  $n=100$  sandwiches every day and recorded the number of the chicken ones:

$$X_1, X_2, \dots, X_N$$

- What is the MLE of the  $p$  parameter?

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = 0$$



# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{\sum_{i=1}^N X_i}{100N}$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N \left[ \log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p) \right]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{\sum_{i=1}^N X_i}{100N}$$

# BREAK

Task: find a coin 😊

# **RANDOMIZED RESPONSE**

*Asking embarrassing questions*

# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.

# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.
- BUT WHAT IF the question is very sensitive?
  - People won't tell the truth.



# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.
- BUT WHAT IF the question is very sensitive?
  - People won't tell the truth.
- EXAMPLE: *Do you find this course boring?*
  - How do I find out what my students *actually* think?

# STRATEGY: RANDOMIZED RESPONSE

- Watch the video about the randomized response strategy.
- In groups, discuss the strategy and complete the assignment.
- See Google Classroom.

# LET'S TRY THIS OUT!

- Toss a coin...

# LET'S TRY THIS OUT!

- Toss a coin...
- Now, answer one of the following questions:

# LET'S TRY THIS OUT!

- Toss a coin...
- Now, answer one of the following questions:

<b>If you got HEADS:</b>	DO YOU FIND THIS CLASS BORING?
<b>If you got TAILS:</b>	ARE YOU AT THE STATISTICS CLASS RIGHT NOW?

# TO SUM UP

- **Likelihood function** is the joint probability of realized sample given the parameters.
- **Maximum Likelihood Estimate (MLE)** is the value which maximizes the probability of observing the realized sample.