

INTRODUCTION TO STATISTICS

LECTURE 7

LAST TIME

- Normal distribution
 - probability of being in an interval;
 - some properties;
 - MLE for μ and σ .

LAST TIME

- Normal distribution
 - probability of being in an interval;
 - some properties;
 - MLE for μ and σ .
- Central limit theorem – an introduction.

TODAY

- Central limit theorem
 - revision;
 - practical exercise.

TODAY

- Central limit theorem
 - revision;
 - practical exercise.
- Properties of point estimators
 - bias, variance and consistency.

TODAY

- Central limit theorem
 - revision;
 - practical exercise.
- Properties of point estimators
 - bias, variance and consistency.
- Confidence intervals

CENTRAL LIMIT THEOREM

Why normal distribution is so important

CENTRAL LIMIT THEOREM

In the [video](#):


- Instead of measuring every single rabbit, weigh samples of size N and compute sample means.



CENTRAL LIMIT THEOREM

In the [video](#):

- Instead of measuring every single rabbit, weigh samples of size N and compute sample means.
- **Central limit theorem (informally):** the larger the N , the more “normal” the distribution of the sample averages is.



CENTRAL LIMIT THEOREM

Samples X_1, X_2, \dots, X_n :

- i.i.d.
- a *finite* mean μ and *finite* variance σ^2

CENTRAL LIMIT THEOREM

Samples X_1, X_2, \dots, X_n :

- i.i.d.
- a *finite* mean μ and *finite* variance σ^2

Let

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

CENTRAL LIMIT THEOREM

Samples X_1, X_2, \dots, X_n :

- i.i.d.
- a *finite* mean μ and *finite* variance σ^2

Let

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

Then

$$\bar{X}_n \approx N\left(\mu, \frac{\sigma^2}{n}\right)$$

CENTRAL LIMIT THEOREM

- $X \sim Po(5)$ - number of errors per computer program
 - $E(X) = Var(X) = 5$
- X_1, X_2, \dots, X_{125} - number of errors in the programs.

$$P(\bar{X}_n \leq 5.5) =$$

CENTRAL LIMIT THEOREM

- $X \sim Po(5)$ - number of errors per computer program
 - $E(X) = Var(X) = 5$
- X_1, X_2, \dots, X_{125} - number of errors in the programs.

$$\begin{aligned} P(\bar{X}_n \leq 5.5) &= \\ &= P\left(\frac{\sqrt{125}(\bar{X}_n - 5)}{\sqrt{5}} \leq \right) \approx \end{aligned}$$

CENTRAL LIMIT THEOREM

- $X \sim Po(5)$ - number of errors per computer program
 - $E(X) = Var(X) = 5$
- X_1, X_2, \dots, X_{125} - number of errors in the programs.

$$\begin{aligned} P(\bar{X}_n \leq 5.5) &= \\ &= P\left(\frac{\sqrt{125}(\bar{X}_n - 5)}{\sqrt{5}} \leq \frac{\sqrt{125}(5.5 - 5)}{\sqrt{5}}\right) \approx \end{aligned}$$

CENTRAL LIMIT THEOREM

- $X \sim Po(5)$ - number of errors per computer program
 - $E(X) = Var(X) = 5$
- X_1, X_2, \dots, X_{125} - number of errors in the programs.

$$\begin{aligned} P(\bar{X}_n \leq 5.5) &= \\ &= P\left(\frac{\sqrt{125}(\bar{X}_n - 5)}{\sqrt{5}} \leq \frac{\sqrt{125}(5.5 - 5)}{\sqrt{5}}\right) \approx \\ &\approx P(Z \leq 2.5) = \end{aligned}$$

CENTRAL LIMIT THEOREM

- $X \sim Po(5)$ - number of errors per computer program
 - $E(X) = Var(X) = 5$
- X_1, X_2, \dots, X_{125} - number of errors in the programs.

$$\begin{aligned} P(\bar{X}_n \leq 5.5) &= \\ &= P\left(\frac{\sqrt{125}(\bar{X}_n - 5)}{\sqrt{5}} \leq \frac{\sqrt{125}(5.5 - 5)}{\sqrt{5}}\right) \approx \\ &\approx P(Z \leq 2.5) = 0.9938 \end{aligned}$$

CLT IN ACTION

Google Classroom -> Lecture 7 -> Mean of means

PROPERTIES OF ESTIMATORS

Bias, variance and consistency

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

- $\hat{\theta} = T(X_1, \dots, X_n),$ $T(X)$ – estimator.

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$
- So far, we only constructed ML estimators.

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$

- So far, we only constructed ML estimators.
But *anything* can be an estimator:

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$

- So far, we only constructed ML estimators.

But *anything* can be an estimator:

$$\hat{\theta}_1 = \frac{1}{n} \sum_{i=1}^n X_i, \quad \hat{\theta}_2 = \frac{X_1 + X_n}{2}, \quad \hat{\theta}_3 = \max(X_1, \dots, X_n), \quad \dots$$

PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples X_1, X_2, \dots, X_n

Obtain estimate $\hat{\theta}$ of an unknown parameter θ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$

- So far, we only constructed ML estimators.
But *anything* can be an estimator:

$$\hat{\theta}_1 = \frac{1}{n} \sum_{i=1}^n X_i, \quad \hat{\theta}_2 = \frac{X_1 + X_n}{2}, \quad \hat{\theta}_3 = \max(X_1, \dots, X_n), \quad \dots$$

- We need to compare different estimators.

PROPERTIES OF ESTIMATORS

- Bias
- Variance
- Consistency

BIAS

An estimator $T(X)$ is unbiased of θ if

$$E(T(X)) = \theta$$

BIAS

An estimator $T(X)$ is unbiased of θ if

$$E(T(X)) = \theta$$

Bias is defined as

$$bias(T(X)) = \theta - E(T(X))$$

BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) =$$

BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i \text{ — (un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$

BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) =$$

BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot np = p$$

BIAS: EXAMPLE 2

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

BIAS: EXAMPLE 2

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{\mu}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot n\mu = \mu$$

BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \text{ -- (un)biased?}$$

BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 =$$

BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \left(\frac{1}{n} \sum_{i=1}^n X_i^2 \right)^2 =$$

BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \left(\frac{1}{n} \sum_{i=1}^n X_i^2 \right)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) =$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

$$E X_i^2 = ?$$

$$Var(X_i) = \sigma^2 =$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

$$E X_i^2 = ?$$

$$Var(X_i) = \sigma^2 = E X_i^2 - (E X_i)^2 =$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

$$Var(X_i) = \sigma^2 = EX_i^2 - (EX_i)^2 = EX_i^2 - \mu^2$$

BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

$$Var(X_i) = \sigma^2 = EX_i^2 - (EX_i)^2 = EX_i^2 - \mu^2$$

$$E(X_i^2) = \sigma^2 + \mu^2$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

$$E\bar{X}^2 = \frac{\sigma^2}{n} + \mu^2$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$ is a **biased** estimator for the parameter σ^2 of the Normal distribution.

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$ is a **biased** estimator for the parameter σ^2 of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$ is a **biased** estimator for the parameter σ^2 of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 = \frac{n}{(n-1)} \cdot S^2 =$$

BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$ is a **biased** estimator for the parameter σ^2 of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 = \frac{n}{(n-1)} \cdot S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

VARIANCE

Variance of an estimator $T(X)$ is defined as

$$Var(T(X)) = E\{T(X) - E(T(X))\}^2$$

VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) =$$

VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$

VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?

BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$Var(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$Var(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

but

$$Var \hat{\mu}_2 =$$

BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$\text{Var}(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$


$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

but


$$\text{Var}(\hat{\mu}_2) = \frac{\sigma^2}{n}$$

BIAS VS VARIANCE


1



2



3




4




BIAS VS VARIANCE


1



2



3




4




LOW BIAS
LOW VARIANCE

BIAS VS VARIANCE


1



2



3




4




BIAS VS VARIANCE


1



2



3




4




**LOW BIAS
HIGH VARIANCE**

BIAS VS VARIANCE


1



2



3




4




BIAS VS VARIANCE


1



2



3




4




**HIGH BIAS
LOW VARIANCE**

BIAS VS VARIANCE


1



2



3




4




BIAS VS VARIANCE


1



2



3




4



**HIGH BIAS
HIGH VARIANCE**


BIAS VS VARIANCE

1




**LOW BIAS
LOW VARIANCE**

2



**HIGH BIAS
LOW VARIANCE**

3



**LOW BIAS
HIGH VARIANCE**


4



**HIGH BIAS
HIGH VARIANCE**

BIAS VS VARIANCE


1



**LOW BIAS
LOW VARIANCE**




2



**HIGH BIAS
LOW VARIANCE**

3



**LOW BIAS
HIGH VARIANCE**


4



**HIGH BIAS
HIGH VARIANCE**

BIAS VS VARIANCE


1



**LOW BIAS
LOW VARIANCE**




2



**HIGH BIAS
LOW VARIANCE**

3



**LOW BIAS
HIGH VARIANCE**

4




**HIGH BIAS
HIGH VARIANCE**



BIAS-VARIANCE TRADE-OFF

- Impossible to simultaneously optimize bias and variance.
- Related to *under-* and *overfitting* in Machine Learning.



**LOW BIAS
LOW VARIANCE**

CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

Let T_1, T_2, \dots, T_n be a sequence of estimators for θ , $T_k = T(X_1, \dots, X_k)$.

CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

Let T_1, T_2, \dots, T_n be a sequence of estimators for θ , $T_k = T(X_1, \dots, X_k)$.

Then $\{T_n\}$ is consistent if $\forall \epsilon > 0$

$$\lim_{n \rightarrow \infty} P(|T_n - \theta| < \epsilon) = 1$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad Var(\hat{\mu}) = \sigma^2/n$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu,$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad Var(\hat{\mu}) = \sigma^2/n$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq$$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} =$$

Chebyshev's inequality:

$$P\left(|X - \mu| \geq k\sigma\right) \leq \frac{1}{k^2}$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} = \frac{\sigma^2}{n\epsilon^2}$$

CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose $\epsilon = \frac{c\sigma}{\sqrt{n}}$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} = \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \quad n \rightarrow \infty$$

PROPERTIES OF MAX. LIKELIHOOD EST.

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If $\hat{\theta}$ is MLE for θ than $g(\hat{\theta})$ is MLE for $g(\theta)$

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If $\hat{\theta}$ is MLE for θ than $g(\hat{\theta})$ is MLE for $g(\theta)$

3. MLE is **asymptotically efficient**:

PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If $\hat{\theta}$ is MLE for θ than $g(\hat{\theta})$ is MLE for $g(\theta)$

3. MLE is **asymptotically efficient**:
roughly speaking, among well-behaved estimators, it has the smallest variance, at least for large samples.

CONFIDENCE INTERVALS

MOTIVATION

- Point estimation provides a single “best guess”.

MOTIVATION

- Point estimation provides a single “best guess”.
- How confident can we be in our estimation?

MOTIVATION

- Point estimation provides a single “best guess”.
- How confident can we be in our estimation?
- CI proposes a range of plausible values.

DEFINITION

A $1 - \alpha$ confidence interval for a parameter θ is an interval $C_n = (a, b)$ such that $T_1 = t_1(X_1, \dots, X_n)$, $T_2 = t_2(X_1, \dots, X_n)$ and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

DEFINITION

A $1 - \alpha$ confidence interval for a parameter θ is an interval $C_n = (T_1, T_2)$ such that $T_1 = t_1(X_1, \dots, X_n)$, $T_2 = t_2(X_1, \dots, X_n)$ and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals: T_1 and T_2 are functions of random samples.

DEFINITION

A $1 - \alpha$ confidence interval for a parameter θ is an interval $C_n = (T_1, T_2)$ such that $T_1 = t_1(X_1, \dots, X_n)$, $T_2 = t_2(X_1, \dots, X_n)$ and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals: T_1 and T_2 are functions of random samples.
- θ is unknown, but fixed
 T_1 and T_2 are random

INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about θ since it's fixed.

INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about θ since it's fixed.
- Common interpretation:

INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about θ since it's fixed.
- Common interpretation:

If I repeat the experiment many times, the interval will contain the true value of θ 95% of the time ($\alpha=0.05$).

CI FOR MEAN (VARIANCE IS KNOWN)

CI FOR μ (σ IS KNOWN)

- Suppose that X_1, X_2, \dots, X_n - i.i.d. samples from a distribution with unknown mean μ and known variance σ^2 .

CI FOR μ (σ IS KNOWN)

- Suppose that X_1, X_2, \dots, X_n - i.i.d. samples from a distribution with unknown mean μ and known variance σ^2 .
- Point estimate for μ :

$$\mu = \bar{X} = \sum_{i=1}^n X_i$$

CI FOR μ (σ IS KNOWN)

- Suppose that X_1, X_2, \dots, X_n - i.i.d. samples from a distribution with unknown mean μ and known variance σ^2 .
- Point estimate for μ :

$$\mu = \bar{X} = \sum_{i=1}^n X_i$$

- How to compute a confidence interval?

CI FOR μ (σ IS KNOWN)

Central Limit Theorem:

CI FOR μ (σ IS KNOWN)

Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right)$

CI FOR μ (σ IS KNOWN)

Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X}-\mu)}{\sigma} \approx N(0, 1)$


CI FOR μ (σ IS KNOWN)

Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$

$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)


Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)


Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)


Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)


Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)


Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{\alpha/2} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{\alpha/2}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

Central Limit Theorem: $\bar{X} \approx N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} \approx N(0, 1)$



$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

$$P\left(\bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.95$

$$P\left(\bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.95$

$$P\left(5 - \frac{1}{10} z_{1-\frac{0.05}{2}} < \mu < 5 + \frac{1}{10} z_{1-\frac{0.05}{2}}\right) = 0.95$$

CI FOR μ (σ IS KNOWN)

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.95$

$$P\left(5 - \frac{1}{10} z_{1-\frac{0.05}{2}} < \mu < 5 + \frac{1}{10} z_{1-\frac{0.05}{2}}\right) = 0.95$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

CI FOR μ (σ IS KNOWN)

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.95$

$$P\left(5 - \frac{1}{10} z_{1-\frac{0.05}{2}} < \mu < 5 + \frac{1}{10} z_{1-\frac{0.05}{2}}\right) = 0.95$$

$$P(5 - 0.196 < \mu < 5 + 0.196) = 0.95$$

CI FOR μ (σ IS KNOWN)

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.95$

$$P\left(5 - \frac{1}{10} z_{1-\frac{0.05}{2}} < \mu < 5 + \frac{1}{10} z_{1-\frac{0.05}{2}}\right) = 0.95$$

$$P(5 - 0.196 < \mu < 5 + 0.196) = 0.95$$

$$(5 - 0.196 ; 5 + 0.196)$$

CI FOR μ (σ IS KNOWN)

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.90$

$$P\left(\bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

CI FOR μ (σ IS KNOWN)

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.90$

$$P\left(5 - \frac{1}{10} z_{1-\frac{0.1}{2}} < \mu < 5 + \frac{1}{10} z_{1-\frac{0.1}{2}}\right) = 0.90$$

CI FOR μ (σ IS KNOWN)

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

- Example:

$$n = 100, \quad \bar{X} = 5, \quad \sigma = 1$$

- Construct CI for μ at the level $1 - \alpha = 0.90$

$$P\left(5 - \frac{1}{10}z_{0.95} < \mu < 5 + \frac{1}{10}z_{0.95}\right) = 0.90$$

$$P(5 - 0.164 < \mu < 5 + 0.164) = 0.90$$

$$(5 - 0.164 ; 5 + 0.164)$$

TO SUM SUP

- Central Limit Theorem
- Properties of estimators
 - bias, variance, consistency;
 - properties of ML estimates.
- Confidence intervals

MID-TERM

- Tomorrow, Wednesday, December 9
- 09:00 – 12:00 (no class)
- Assignment will become available on Google Classroom
- You should submit by 12:00

Topics:

- Descriptive statistics
- Discrete distributions
- Continuous random variables (CDFs, PDFs, probabilities)
- Maximum likelihood (discrete and continuous)

MID-TERM

- Tomorrow, Wednesday, December 9
- 09:00 – 12:00 (no class)
- Assignment will become available on Google Classroom
- You should submit by 12:00