

PROBABILITY & STATISTICS

Lecture 2 – Basic probability review

LAST TIME

- Probability as limiting frequency
- Basic combinatorics review

TODAY

- Review of practice problems
- Probability of an event: more examples
- Axioms of probability

Practice problems review

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

2 freshmen, 1 sophomore and 1 junior

1 freshman, 2 sophomores and 1 junior

1 freshman, 1 sophomore and 2 juniors

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

2 freshmen, 1 sophomore and 1 junior

$$C(6,2) \cdot C(5,1) \cdot C(4,1) = N_{211}$$

1 freshman, 2 sophomores and 1 junior

1 freshman, 1 sophomore and 2 juniors

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

2 freshmen, 1 sophomore and 1 junior

$$C(6,2) \cdot C(5,1) \cdot C(4,1) = N_{211}$$

1 freshman, 2 sophomores and 1 junior

$$C(6,1) \cdot C(5,2) \cdot C(4,1) = N_{121}$$

1 freshman, 1 sophomore and 2 juniors

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

2 freshmen, 1 sophomore and 1 junior

$$C(6,2) \cdot C(5,1) \cdot C(4,1) = N_{211}$$

1 freshman, 2 sophomores and 1 junior

$$C(6,1) \cdot C(5,2) \cdot C(4,1) = N_{121}$$

1 freshman, 1 sophomore and 2 juniors

$$C(6,1) \cdot C(5,1) \cdot C(4,2) = N_{112}$$

COMMITTEE

- A total of 6 freshmen, 5 sophomores and 4 juniors have volunteered to serve on a 4-person committee. How many such committees are possible if at least one freshman, one sophomore and one junior must serve on the committee?

2 freshmen, 1 sophomore and 1 junior

$$C(6,2) \cdot C(5,1) \cdot C(4,1) = N_{211}$$

1 freshman, 2 sophomores and 1 junior

$$C(6,1) \cdot C(5,2) \cdot C(4,1) = N_{121}$$

1 freshman, 1 sophomore and 2 juniors

$$C(6,1) \cdot C(5,1) \cdot C(4,2) = N_{112}$$

$N_{211} + N_{121} + N_{112}$ committees with at least one student from each group

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?


DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

$$|\heartsuit| = ?$$

$$|\spadesuit| = ?$$

$$|\heartsuit \cap \spadesuit| = ?$$



A set of shuffles where ace of hearts **is** directly on top of king of hearts.



A set of shuffles where ace of spades **is** directly on top of king of spades.

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

shuffles where ace of hearts **is** on top of king of hearts

shuffles where ace of spades **is** on top of king of spades

shuffles where both is true

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

shuffles where ace of hearts **is** on top of king of hearts
51!

shuffles where ace of spades **is** on top of king of spades

shuffles where both is true

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

shuffles where ace of hearts **is** on top of king of hearts
51!

shuffles where ace of spades **is** on top of king of spades
51!

shuffles where both is true

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

shuffles where ace of hearts **is** on top of king of hearts
51!

shuffles where ace of spades **is** on top of king of spades
51!

shuffles where both is true
50!

DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

shuffles where ace of hearts **is** on top of king of hearts
51!

shuffles where ace of spades **is** on top of king of spades
51!

shuffles where both is true
50!

So, the number of possible shuffles is

$$52! - (51! + 51! - 50!) = 52! - 2 \cdot 51! + 50!$$


DECK OF CARDS

- How many shuffles are there of a deck of cards, such that ace of hearts is not directly on top of king of hearts, and ace of spades is not directly on top of king of spades?

$$|\heartsuit| = 51!$$

$$|\spadesuit| = 51!$$

$$|\heartsuit \cap \spadesuit| = 50!$$



A set of shuffles where ace of hearts **is** directly on top of king of hearts.



A set of shuffles where ace of spades **is** directly on top of king of spades.

DOUGHNUTS

- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

DOUGHNUTS

- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order
Repetitions

DOUGHNUTS


- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order doesn't matter.

Repetitions are allowed



Combinations with repetitions



DOUGHNUTS


- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order doesn't matter.

Repetitions are allowed



Combinations with repetitions



DOUGHNUTS


- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



$$C(\quad, 5 - 1) =$$

DOUGHNUTS


- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



$$C(8 + 5 - 1, 5 - 1) =$$

DOUGHNUTS


- You went to the doughnut store to buy a box of eight doughnuts. If the doughnut store has five varieties, how many ways are there to fill the box?

Order doesn't matter.

Repetitions are allowed

→

Combinations with repetitions



$$C(8 + 5 - 1, 5 - 1) = C(12, 4) = \frac{12!}{4!8!} = 495$$

ways to fill a box of 8 doughnuts

Back to probability

REMINDER: BASIC DEFINITIONS

- A **random experiment** is a process by which we observe something uncertain.
- The result of a random experiment is called the **outcome**.
- The set of all possible outcomes is called the **sample space** (denoted by S).

REMINDER: RELATIVE FREQUENCY

- Probability of an event is defined by the **limiting frequency** with which this event appears in a long series of similar experiments.
- Example:
If we flip a fair coin infinitely many times, it will come up heads half of the times.

Programming Exercise

Google Classroom -> Day 2

Axioms of probability

AXIOMS OF PROBABILITY

1. For any event E , $P(E) \geq 0$.

AXIOMS OF PROBABILITY

1. For any event E , $P(E) \geq 0$.

Example: rolling a die

$$E_1: \text{'get 1'} \qquad P(E_1) = \frac{1}{6}$$

$$E_2: \text{'get an even number'} \qquad P(E_2) = \frac{1}{2}$$

$$E_3: \text{'get 27'} \qquad P(E_3) = 0$$

AXIOMS OF PROBABILITY

2. Probability of the sample space S is $P(S) = 1$.

AXIOMS OF PROBABILITY

2. Probability of the sample space S is $P(S) = 1$.

Example: tossing a coin

$$S = \{H, T\}$$

$$E = S: \text{'get Heads or Tails'} \qquad P(E) = 1$$

AXIOMS OF PROBABILITY

3. If E_1, E_2, E_3, \dots are mutually exclusive events, then

$$P(E_1 \cup E_2 \cup E_3 \dots) = P(E_1) + P(E_2) + P(E_3) + \dots$$

AXIOMS OF PROBABILITY

3. If E_1, E_2, E_3, \dots are mutually exclusive events, then

$$P(E_1 \cup E_2 \cup E_3 \dots) = P(E_1) + P(E_2) + P(E_3) + \dots$$

- Example:

Candidates A, B, C and D are participating in the elections. Based on the polls, A has a 20% chance of winning and B has a 40% chance of winning. What is the probability that A **or** B will win?

AXIOMS OF PROBABILITY

3. If E_1, E_2, E_3, \dots are mutually exclusive events, then

$$P(E_1 \cup E_2 \cup E_3 \dots) = P(E_1) + P(E_2) + P(E_3) + \dots$$

- Example:

Candidates A, B, C and D are participating in the elections. Based on the polls, A has a 20% chance of winning and B has a 40% chance of winning. What is the probability that A **or** B will win?

$$P(E_A \cup E_B) =$$

AXIOMS OF PROBABILITY

3. If E_1, E_2, E_3, \dots are mutually exclusive events, then

$$P(E_1 \cup E_2 \cup E_3 \dots) = P(E_1) + P(E_2) + P(E_3) + \dots$$

- Example:

Candidates A, B, C and D are participating in the elections. Based on the polls, A has a 20% chance of winning and B has a 40% chance of winning. What is the probability that A **or** B will win?

$$P(E_A \cup E_B) = P(E_A) + P(E_B) = 0.2 + 0.4 = 0.6$$

Complementary events

COMPLEMENT OF AN EVENT

- A complement $E^C = \bar{E}$ of an event E is a set of all outcomes from the sample space S that don't belong to E .

COMPLEMENT OF AN EVENT

- A complement $E^C = \bar{E}$ of an event E is a set of all outcomes from the sample space S that don't belong to E .

Example 1: flipping a coin

$$S = \{H, T\}$$

$$E = \{H\} \quad \bar{E} =$$

COMPLEMENT OF AN EVENT

- A complement $E^C = \bar{E}$ of an event E is a set of all outcomes from the sample space S that don't belong to E .

Example 1: flipping a coin

$$S = \{H, T\}$$

$$E = \{H\} \qquad \bar{E} = \{T\}$$

COMPLEMENT OF AN EVENT

- A complement $E^C = \bar{E}$ of an event E is a set of all outcomes from the sample space S that don't belong to E .

Example 1: flipping a coin

$$S = \{H, T\}$$

$$E = \{H\} \quad \bar{E} = \{T\}$$

Example 2: rolling a die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2, 3\} \quad \bar{E} =$$

COMPLEMENT OF AN EVENT

- A complement $E^C = \bar{E}$ of an event E is a set of all outcomes from the sample space S that don't belong to E .

Example 1: flipping a coin

$$S = \{H, T\}$$

$$E = \{H\} \quad \bar{E} = \{T\}$$

Example 2: rolling a die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2, 3\} \quad \bar{E} = \{4, 5, 6\}$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2\}$$

$$P(E) =$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2\}$$

$$P(E) = \frac{2}{6} = \frac{1}{3}$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2\}$$

$$P(E) = \frac{2}{6} = \frac{1}{3}$$

$$E^C = \{3, 4, 5, 6\}$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2\}$$

$$P(E) = \frac{2}{6} = \frac{1}{3}$$

$$E^C = \{3, 4, 5, 6\}$$

$$P(E^C) = 1 - P(E) =$$

PROBABILITIES OF COMPLEMENTARY EVENTS

- If E is an event and E^C is its complement, then

$$P(E) = 1 - P(E^C)$$

- Example:

Rolling a fair die

$$S = \{1, 2, 3, 4, 5, 6\}$$

$$E = \{1, 2\}$$

$$P(E) = \frac{2}{6} = \frac{1}{3}$$

$$E^C = \{3, 4, 5, 6\}$$

$$P(E^C) = 1 - P(E) = 1 - \frac{1}{3} = \frac{2}{3}$$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| =$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

E^C : 'none of the bits is 0'

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

E^C : 'none of the bits is 0' $|E^C| =$ $P(E^C) =$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

E^C : 'none of the bits is 0' $|E^C| = 1$ $P(E^C) = \frac{1}{2^{10}}$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

E^C : 'none of the bits is 0' $|E^C| = 1$ $P(E^C) = \frac{1}{2^{10}}$

$\rightarrow P(E) =$

EXAMPLE: BIT STRINGS

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

S — all possible 10-bit strings. $|S| = 2^{10}$

E : 'at least one of the bits is 0' $P(E) = ?$

E^C : 'none of the bits is 0' $|E^C| = 1$ $P(E^C) = \frac{1}{2^{10}}$

$$\rightarrow P(E) = 1 - P(E^C) = 1 - 1/2^{10}$$

Union of events

PROBABILITY OF A UNION

- Let E_1 and E_2 be some events in a sample space S . Then

$$P(E_1 \cup E_2) =$$

PROBABILITY OF A UNION

- Let E_1 and E_2 be some events in a sample space S . Then

$$P(E_1 \cup E_2) = \frac{|E_1 \cup E_2|}{|S|} =$$

PROBABILITY OF A UNION

- Let E_1 and E_2 be some events in a sample space S . Then

$$P(E_1 \cup E_2) = \frac{|E_1 \cup E_2|}{|S|} = \frac{|E_1| + |E_2| - |E_1 \cap E_2|}{|S|} =$$

=

PROBABILITY OF A UNION

- Let E_1 and E_2 be some events in a sample space S . Then

$$P(E_1 \cup E_2) = \frac{|E_1 \cup E_2|}{|S|} = \frac{|E_1| + |E_2| - |E_1 \cap E_2|}{|S|} =$$

$$= \frac{|E_1|}{|S|} + \frac{|E_2|}{|S|} - \frac{|E_1 \cap E_2|}{|S|} =$$

PROBABILITY OF A UNION

- Let E_1 and E_2 be some events in a sample space S . Then

$$P(E_1 \cup E_2) = \frac{|E_1 \cup E_2|}{|S|} = \frac{|E_1| + |E_2| - |E_1 \cap E_2|}{|S|} =$$

$$= \frac{|E_1|}{|S|} + \frac{|E_2|}{|S|} - \frac{|E_1 \cap E_2|}{|S|} = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

DRAWING NAMES

A total of 20 students participate in a drawing in which one name is chosen at random. There are 6 seniors (2 men and 4 women), 5 juniors (3 men and 2 women) and 9 freshmen (5 men and 4 women). What is the probability that the name that is drawn belongs to **either a woman or to a junior**?

DRAWING NAMES

A total of 20 students participate in a drawing in which one name is chosen at random. There are 6 seniors (2 men and 4 women), 5 juniors (3 men and 2 women) and 9 freshmen (5 men and 4 women). What is the probability that the name that is drawn belongs to **either a woman or to a junior**?

$$P(W \cup J) = P(W) + P(J) - P(W \cap J) =$$

=

DRAWING NAMES

A total of 20 students participate in a drawing in which one name is chosen at random. There are 6 seniors (2 men and 4 women), 5 juniors (3 men and 2 women) and 9 freshmen (5 men and 4 women). What is the probability that the name that is drawn belongs to **either a woman or to a junior**?

$$P(W \cup J) = P(W) + P(J) - P(W \cap J) =$$

$$= \frac{|W|}{|S|} + \frac{|J|}{|S|} - \frac{|W \cap J|}{|S|} =$$

=

DRAWING NAMES

A total of 20 students participate in a drawing in which one name is chosen at random. There are 6 seniors (2 men and 4 women), 5 juniors (3 men and 2 women) and 9 freshmen (5 men and 4 women). What is the probability that the name that is drawn belongs to **either a woman or to a junior**?

$$\begin{aligned} P(W \cup J) &= P(W) + P(J) - P(W \cap J) = \\ &= \frac{|W|}{|S|} + \frac{|J|}{|S|} - \frac{|W \cap J|}{|S|} = \\ &= \frac{4 + 2 + 4}{20} + \frac{5}{20} - \frac{2}{20} = \end{aligned}$$

DRAWING NAMES

A total of 20 students participate in a drawing in which one name is chosen at random. There are 6 seniors (2 men and 4 women), 5 juniors (3 men and 2 women) and 9 freshmen (5 men and 4 women). What is the probability that the name that is drawn belongs to **either a woman or to a junior**?

$$\begin{aligned} P(W \cup J) &= P(W) + P(J) - P(W \cap J) = \\ &= \frac{|W|}{|S|} + \frac{|J|}{|S|} - \frac{|W \cap J|}{|S|} = \\ &= \frac{4 + 2 + 4}{20} + \frac{5}{20} - \frac{2}{20} = 0.5 + 0.25 - 0.1 = 0.65 \end{aligned}$$

More examples

PROBABILITY OF AN EVENT

- Consider a random experiment where all outcomes are equally likely.
- Probability $P(E)$ of an event E is such an experiment is

$$P(E) = \frac{\text{\# ways } E \text{ can happen}}{\text{\# possible outcomes}}$$

- Let E^C be the complementary event. Then

$$P(E^C) = 1 - P(E)$$

- Let F be some other event. Then

$$P(E \cup F) = P(E) + P(F) - P(E \cap F)$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$$|S| =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$$|E| =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

$$P(E) =$$

ROLLING DICE

- Two fair dice are rolled.
Compute the probability that the sum of the faces is 3.

$|S| = 6 \cdot 6 = 36$ different pairs of values are possible

$|E| = |\{(1, 2), (2, 1)\}| = 2$
of the combinations give 3 in total

$$P(E) = \frac{2}{36} = \frac{1}{18} \approx 0.056$$

Practice problems

Google Classroom -> Lecture 2

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$|E| = |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| =$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = 15 + 10 + 6 = 31 \\ &\text{ways to choose 2 same-color balls} \end{aligned}$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 1: combinations (*grabbing 2 balls at the same time*)

$$|S| = C(15, 2) = \frac{15!}{2!13!} = 105 \text{ ways to choose 2 balls out of 15}$$

$$\begin{aligned} |E| &= |E_{2 \text{ red}}| + |E_{2 \text{ blue}}| + |E_{2 \text{ green}}| = \\ &= C(6, 2) + C(5, 2) + C(4, 2) = 15 + 10 + 6 = 31 \\ &\text{ways to choose 2 same-color balls} \end{aligned}$$

$$P(E) = 31/105$$

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

BALLS FROM A BOWL

- A bowl contains 15 balls: 6 are red, 5 are blue and 4 are green. What is the probability that two balls selected at random from the bowl **have the same color?**

Method 2: 2-permutations (*grabbing 2 balls one-by-one*)

$|S| = 15 \cdot 14 = 210$ ways to pick 2 balls out of 15 one-by-one

$$\begin{aligned} |E| &= |E_{red,red}| + |E_{blue,blue}| + |E_{green,green}| = \\ &= 6 \cdot 5 + 5 \cdot 4 + 4 \cdot 3 = 30 + 20 + 12 = 62 \\ &\text{ways to pick two same-color balls one by one} \end{aligned}$$

$$P(E) = 62/210 = 31/105$$

NUMBERS

- What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

$$P(E_2 \cup E_5) =$$

NUMBERS

- What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

$$P(E_2 \cup E_5) = P(E_2) + P(E_5) - P(E_2 \cap E_5) =$$

=

NUMBERS

- What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

$$\begin{aligned} P(E_2 \cup E_5) &= P(E_2) + P(E_5) - P(E_2 \cap E_5) = \\ &= \frac{|E_2|}{|S|} + \frac{|E_5|}{|S|} - \frac{|E_2 \cap E_5|}{|S|} = \\ &= \end{aligned}$$

NUMBERS

- What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

$$\begin{aligned} P(E_2 \cup E_5) &= P(E_2) + P(E_5) - P(E_2 \cap E_5) = \\ &= \frac{|E_2|}{|S|} + \frac{|E_5|}{|S|} - \frac{|E_2 \cap E_5|}{|S|} = \\ &= \frac{50}{100} + \frac{20}{100} - \frac{10}{100} = 0.6 \end{aligned}$$

POKER

- In poker, each player is dealt (supplied with) 5 of the 52 cards, called a **hand**. How many different hands are there?

POKER

- In poker, each player is dealt (supplied with) 5 of the 52 cards, called a **hand**. How many different hands are there?

$$C(52, 5) = \frac{52!}{5! \cdot 47!} = 2,598,960$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

$$P(E) =$$

POKER

- What is the probability of being dealt a poker hand that consists of the 3♥, 5♦, 6♠, 10♣ and Q♦?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = 1$$

$$P(E) = \frac{1}{2,598,960} \approx 0.000000385$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

$$P(E) =$$

POKER

- A hand that consists of five cards of the same suit is called a **flush**. What is the probability of getting a flush?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(4, 1) \cdot C(13, 5) = 4 \cdot \frac{13!}{5! 8!} = 5148$$

$$P(E) = \frac{5148}{2,598,960} \approx 0.00198$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$|E| = C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

$$P(E) =$$

POKER

- A **pair** in poker are two cards of the same value (e.g., two twos or two eights). What is the probability of being dealt a hand of two pairs in poker?

$$|S| = C(52, 5) = 2,598,960$$

$$\begin{aligned} |E| &= C(13, 2) \cdot C(4, 2) \cdot C(4, 2) \cdot C(44, 1) = \\ &= 78 \cdot 6 \cdot 6 \cdot 44 = 123552 \end{aligned}$$

$$P(E) = \frac{123552}{2,598,960} \approx 0.048$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$P(E_3 \cup E_7) =$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$P(E_3 \cup E_7) = P(E_3) + P(E_7) - P(E_3 \cap E_7) =$$

NUMBERS

- What is the probability that a integer between 1 and 100 is divisible by 3 or 7?

$$\begin{aligned} P(E_3 \cup E_7) &= P(E_3) + P(E_7) - P(E_3 \cap E_7) = \\ &= \frac{33}{100} + \frac{14}{100} - \frac{4}{100} = \frac{43}{100} = 0.43 \end{aligned}$$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- ... **all girls are standing next to each other?**
- ... **all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- ... **all girls are standing next to each other?**
 $|S| =$
- ... **all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| =$
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) =$$

- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| =$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| = 10!$ possible permutations of 10 kids

$|E| =$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| = 10!$ possible permutations of 10 kids

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...
- **... all girls are standing next to each other?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 7!$ possible permutations where girls are together
$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$
- **... all the girls are next to each other and all the boys as well?**
 $|S| = 10!$ possible permutations of 10 kids
 $|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together

$$P(E) =$$

CLASS PICTURE

- At school, 4 girls and 6 boys are lining up for a class picture in a random order. What is the probability that ...

- **... all girls are standing next to each other?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 7!$ possible permutations where girls are together

$$P(E) = \frac{4! \cdot 7!}{10!} \approx 0.033$$

- **... all the girls are next to each other and all the boys as well?**

$|S| = 10!$ possible permutations of 10 kids

$|E| = 4! \cdot 6! \cdot 2!$ permutations where girls are together and boys are together

$$P(E) = \frac{4! \cdot 6! \cdot 2!}{10!} \approx 0.0096$$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- ... **it doesn't contain digit 7?**
- ... **it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- ... **it doesn't contain digit 7?**
 $|S| =$
- ... **it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| =$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| =$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| =$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| =$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...
- **... it doesn't contain digit 7?**
 $|S| = 101$ different numbers
 $|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7
 $|P| = 82/101 \approx 0.81$
- **... it contains unique digits (e.g., 123 but 111)?**
 $|S| = 101$ different numbers
 $|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...

- **... it doesn't contain digit 7?**

$|S| = 101$ different numbers

$|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7

$$|P| = 82/101 \approx 0.81$$

- **... it contains unique digits (e.g., 123 but 111)?**

$|S| = 101$ different numbers

$|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits

$$|P| =$$

RANDOM NUMBER

- A computer is generating a random number between 0 and 100.
What is the probability that ...

- **... it doesn't contain digit 7?**

$|S| = 101$ different numbers

$|E| = 9 + 8 \cdot 9 + 1 = 82$ numbers don't contain 7

$$|P| = 82/101 \approx 0.81$$

- **... it contains unique digits (e.g., 123 but 111)?**

$|S| = 101$ different numbers

$|E| = 10 + 9 \cdot 9 = 91$ numbers with unique digits

$$|P| = 91/101 \approx 0.9$$

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

ways to distribute candies

ways to distribute candies so that each kid gets at least one.

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

ways to distribute candies so that each kid gets at least one.

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

$$P(E) =$$

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

$$P(E) = \frac{C(14, 6)}{C(21, 6)} =$$

CANDIES

- You are distributing 15 candies among 7 kids. What is the probability that each kid gets at least one candy?

Combinations with repetitions:

candies = 'balls', kids = 'colors', 'categories'

$C(15 + 7 - 1, 7 - 1) = 54264$ ways to distribute candies

$C(8 + 7 - 1, 7 - 1) = 3003$ ways to distribute candies so that each kid gets at least one.

$$P(E) = \frac{C(14, 6)}{C(21, 6)} = \frac{3003}{54264} = 0.055$$

Monty Hall

<https://youtu.be/4Lb-6rxZxx0>

MONTY HALL

