

# PROBABILITY & STATISTICS

Lecture 11 – Confidence Intervals

# MLE FOR NORMAL DISTRIBUTIONS

estimating  $\mu$  and  $\sigma$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.
- $\hat{\mu} = ?$        $\hat{\sigma} = ?$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.
- $\hat{\mu} = ?$        $\hat{\sigma} = ?$
- Maximum likelihood.

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- $\hat{\mu} = ? \quad \hat{\sigma} = ?$

$$L(\mu, \sigma) =$$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- $\hat{\mu} = ? \quad \hat{\sigma} = ?$

$$L(\mu, \sigma) = \prod_{i=1}^n \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{X_i - \mu}{\sigma}\right)^2}$$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- $\hat{\mu} = ? \quad \hat{\sigma} = ?$

$$L(\mu, \sigma) = \prod_{i=1}^n \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{X_i - \mu}{\sigma}\right)^2}$$

$$\log L(\mu, \sigma) =$$



# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- $\hat{\mu} = ? \quad \hat{\sigma} = ?$

$$L(\mu, \sigma) = \prod_{i=1}^n \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{X_i - \mu}{\sigma}\right)^2}$$

$$\log L(\mu, \sigma) = - \sum_{i=1}^n \log(\sigma\sqrt{2\pi}) - \frac{1}{2} \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2 =$$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- $\hat{\mu} = ? \quad \hat{\sigma} = ?$

$$L(\mu, \sigma) = \prod_{i=1}^n \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{X_i - \mu}{\sigma}\right)^2}$$

$$\begin{aligned} \log L(\mu, \sigma) &= -\sum_{i=1}^n \log(\sigma\sqrt{2\pi}) - \frac{1}{2} \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2 = \\ &= -n \log \sigma - n \log \sqrt{2\pi} - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - \mu)^2 \end{aligned}$$

# MLE FOR $\mu$ AND $\sigma$

$$\log L(\mu, \sigma) = -n \log \sigma - n \log \sqrt{2\pi} - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - \mu)^2$$

$$\frac{d}{d\mu} \log L(\mu, \sigma) =$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) =$$

# MLE FOR $\mu$ AND $\sigma$

$$\log L(\mu, \sigma) = -n \log \sigma - n \log \sqrt{2\pi} - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - \mu)^2$$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) =$$

# MLE FOR $\mu$ AND $\sigma$

$$\log L(\mu, \sigma) = -n \log \sigma - n \log \sqrt{2\pi} - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - \mu)^2$$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \quad \frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \qquad \frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \quad \frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$



# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \quad \frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0 \quad \frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \mu)^2 = 0$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \bar{X})^2 = 0$$

# MLE FOR $\mu$ AND $\sigma$

$$\frac{d}{d\mu} \log L(\mu, \sigma) = \frac{1}{\sigma^2} \sum_{i=1}^n (X_i - \mu) = 0$$

$$\frac{1}{\sigma^2} \neq 0 \Rightarrow$$

$$\sum_{i=1}^n (X_i - \mu) = \sum_{i=1}^n X_i - n\mu = 0$$

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

$$\frac{d}{d\sigma} \log L(\mu, \sigma) = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \mu)^2 = 0$$

$$-n\sigma^2 + \sum_{i=1}^n (X_i - \bar{X})^2 = 0$$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = S^2$$

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.
- $\hat{\mu} = ?$        $\hat{\sigma} = ?$
- Maximum likelihood:

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.
- $\hat{\mu} = ?$        $\hat{\sigma} = ?$
- Maximum likelihood:
  - $\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$  - **MLE for  $\mu$  is sample mean**

# MLE FOR $\mu$ AND $\sigma$

- $X_1, X_2, \dots, X_n$  – samples from the normal distribution.
- Parameters  $\mu$  and  $\sigma$  are unknown.
- $\hat{\mu} = ?$        $\hat{\sigma} = ?$
- Maximum likelihood:
  - $\hat{\mu} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$  - **MLE for  $\mu$  is sample mean**
  - $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = S^2$  - **MLE for  $\sigma^2$  is sample variance**



# PROPERTIES OF ESTIMATORS

*Bias, variance and consistency*

# PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples  $X_1, X_2, \dots, X_n$

Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .

# PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples  $X_1, X_2, \dots, X_n$

Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .

- $\hat{\theta} = T(X_1, \dots, X_n),$                        $T(X)$  – estimator.

# PROPERTIES OF ESTIMATORS

- Parameter estimation:  
Given samples  $X_1, X_2, \dots, X_n$   
Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .
- $\hat{\theta} = T(X_1, \dots, X_n),$                        $T(X)$  – estimator.
- So far, we only constructed ML estimators.

# PROPERTIES OF ESTIMATORS

- Parameter estimation:  
Given samples  $X_1, X_2, \dots, X_n$   
Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .
- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$
- So far, we only constructed ML estimators.  
But *anything* can be an estimator:

# PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples  $X_1, X_2, \dots, X_n$

Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$

- So far, we only constructed ML estimators.  
But *anything* can be an estimator:

$$\hat{\theta}_1 = \frac{1}{n} \sum_{i=1}^n X_i, \quad \hat{\theta}_2 = \frac{X_1 + X_n}{2}, \quad \hat{\theta}_3 = \max(X_1, \dots, X_n), \quad \dots$$

# PROPERTIES OF ESTIMATORS

- Parameter estimation:

Given samples  $X_1, X_2, \dots, X_n$

Obtain estimate  $\hat{\theta}$  of an unknown parameter  $\theta$ .

- $\hat{\theta} = T(X_1, \dots, X_n), \quad T(X) - \text{estimator.}$

- So far, we only constructed ML estimators.  
But *anything* can be an estimator:

$$\hat{\theta}_1 = \frac{1}{n} \sum_{i=1}^n X_i, \quad \hat{\theta}_2 = \frac{X_1 + X_n}{2}, \quad \hat{\theta}_3 = \max(X_1, \dots, X_n), \quad \dots$$

- We need to compare different estimators.


# PROPERTIES OF ESTIMATORS

- Bias
- Variance
- Consistency




# BIAS VS VARIANCE


1



2



3




4




# BIAS VS VARIANCE


1



2



3




4




**LOW BIAS  
LOW VARIANCE**

# BIAS VS VARIANCE


1



2



3




4




# BIAS VS VARIANCE


1



2



3




4




**LOW BIAS  
HIGH VARIANCE**

# BIAS VS VARIANCE


1



2



3




4




# BIAS VS VARIANCE


1



2



3




4




**HIGH BIAS  
LOW VARIANCE**

# BIAS VS VARIANCE


1



2



3




4






# BIAS VS VARIANCE


1



2



3



4




**HIGH BIAS  
HIGH VARIANCE**




# BIAS VS VARIANCE

1




**LOW BIAS  
LOW VARIANCE**

2



**HIGH BIAS  
LOW VARIANCE**

3



**LOW BIAS  
HIGH VARIANCE**


4



**HIGH BIAS  
HIGH VARIANCE**

# BIAS VS VARIANCE


1



**LOW BIAS  
LOW VARIANCE**




2



**HIGH BIAS  
LOW VARIANCE**

3



**LOW BIAS  
HIGH VARIANCE**


4



**HIGH BIAS  
HIGH VARIANCE**

# BIAS VS VARIANCE


1



**LOW BIAS  
LOW VARIANCE**




2



**HIGH BIAS  
LOW VARIANCE**

3



**LOW BIAS  
HIGH VARIANCE**

4



**HIGH BIAS  
HIGH VARIANCE**



# BIAS

An estimator  $T(X)$  is unbiased of  $\theta$  if

$$E(T(X)) = \theta$$

# BIAS

An estimator  $T(X)$  is unbiased of  $\theta$  if

$$E(T(X)) = \theta$$

Bias is defined as

$$bias(T(X)) = \theta - E(T(X))$$

# BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

# BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) =$$

# BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i \text{ — (un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$



# BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) =$$

# BIAS: EXAMPLE 1

$$X_1, \dots, X_n \sim \text{Bernoulli}(p)$$

$$\hat{p}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{p}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot np = p$$

# BIAS: EXAMPLE 2

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

## BIAS: EXAMPLE 2

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i - \text{(un)biased?}$$

$$E(\hat{\mu}_{ML}) = E\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot n\mu = \mu$$

# BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

# BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 =$$

# BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \left( \frac{1}{n} \sum_{i=1}^n X_i^2 \right)^2 =$$

# BIAS: EXAMPLE 3

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\sigma}_{ML}^2 = s^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 - \text{(un)biased?}$$

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \left( \frac{1}{n} \sum_{i=1}^n X_i^2 \right)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$



# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) =$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

$$E X_i^2 = ?$$

$$Var(X_i) = \sigma^2 =$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n E X_i^2 - E \bar{X}^2$$

$$E X_i^2 = ?$$

$$Var(X_i) = \sigma^2 = E X_i^2 - (E X_i)^2 =$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

$$Var(X_i) = \sigma^2 = EX_i^2 - (EX_i)^2 = EX_i^2 - \mu^2$$

# BIAS: EXAMPLE 3

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}^2$$

$$E(\hat{\sigma}_{ML}^2) = \frac{1}{n} \sum_{i=1}^n EX_i^2 - E\bar{X}^2$$

$$EX_i^2 = ?$$

$$Var(X_i) = \sigma^2 = EX_i^2 - (EX_i)^2 = EX_i^2 - \mu^2$$

$$E(X_i^2) = \sigma^2 + \mu^2$$



# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$



# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - E\bar{X}^2$$

$$E\bar{X}^2 = ?$$

$$Var(\bar{X}) = E\bar{X}^2 - (E\bar{X})^2 = E\bar{X}^2 - \mu^2$$

$$Var(\bar{X}) = Var\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

$$E\bar{X}^2 = \frac{\sigma^2}{n} + \mu^2$$

## BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 =$$

## BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

## BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance  $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is a **biased** estimator for the parameter  $\sigma^2$  of the Normal distribution.

## BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance  $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is a **biased** estimator for the parameter  $\sigma^2$  of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance  $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is a **biased** estimator for the parameter  $\sigma^2$  of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 = \frac{n}{(n-1)} \cdot S^2 =$$

# BIAS: EXAMPLE 3

$$E(\hat{\sigma}_{ML}^2) = \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 = \frac{(n-1)\sigma^2}{n}$$

Sample variance  $S^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  is a **biased** estimator for the parameter  $\sigma^2$  of the Normal distribution.

- How to make it **unbiased**?

$$S_{unbiased}^2 = \frac{n}{(n-1)} \cdot S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

# VARIANCE

Variance of an estimator  $T(X)$  is defined as

$$Var(T(X)) = E\{T(X) - E(T(X))\}^2$$



# VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

# VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

# VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) =$$

# VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) =$$

# VARIANCE: EXAMPLE

$$X_1, \dots, X_n \sim N(\mu, \sigma^2)$$

$$\hat{\mu}_{ML} = \frac{1}{n} \sum_{i=1}^n X_i, = \bar{X}, \quad \text{Var}(\hat{\mu}_{ML}) = ?$$

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{n\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

# BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?

# BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

# BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$Var(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$



# BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$Var(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$

$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

*but*

$$Var(\hat{\mu}_2) =$$

# BIAS AND VARIANCE EXAMPLE

- Can a biased and an unbiased estimator have the same variance?
- Example:

$$\hat{\mu}_{ML} = \bar{X} - \text{unbiased}$$
$$\text{Var}(\hat{\mu}_{ML}) = \frac{\sigma^2}{n}$$


$$\hat{\mu}_2 = \hat{\mu}_{ML} + 1 - \text{biased}$$

*but*

$$\text{Var}(\hat{\mu}_2) = \frac{\sigma^2}{n}$$

# BIAS-VARIANCE TRADE-OFF

- Impossible to simultaneously optimize bias and variance.
- Related to *under-* and *overfitting* in Machine Learning.



**LOW BIAS  
LOW VARIANCE**

# CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

# CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

Let  $T_1, T_2, \dots, T_n$  be a sequence of estimators for  $\theta$ ,  $T_k = T(X_1, \dots, X_k)$ .

# CONSISTENCY

For a good estimator, as the sample size increases, the values of the estimator should get closer to the parameter being estimated.

Let  $T_1, T_2, \dots, T_n$  be a sequence of estimators for  $\theta$ ,  $T_k = T(X_1, \dots, X_k)$ .

Then  $\{T_n\}$  is consistent if  $\forall \epsilon > 0$

$$\lim_{n \rightarrow \infty} P(|T_n - \theta| < \epsilon) = 1$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$



# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq$$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} =$$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} = \frac{\sigma^2}{n\epsilon^2}$$

# CONSISTENCY: EXAMPLE

$$\hat{\mu} = \bar{X}, \quad E\hat{\mu} = \mu, \quad \text{Var}(\hat{\mu}) = \sigma^2/n$$

Let's show that  $\forall \epsilon > 0 \quad \lim_{n \rightarrow \infty} P(|T_n - \mu| \geq \epsilon) = 0$

Let's chose  $\epsilon = \frac{c\sigma}{\sqrt{n}}$

Chebyshev's inequality:

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

$$P\left(|T_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2} = \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \quad n \rightarrow \infty$$

# PROPERTIES OF MAX. LIKELIHOOD EST.



# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If  $\hat{\theta}$  is MLE for  $\theta$  than  $g(\hat{\theta})$  is MLE for  $g(\theta)$

# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If  $\hat{\theta}$  is MLE for  $\theta$  than  $g(\hat{\theta})$  is MLE for  $g(\theta)$

3. MLE is **asymptotically efficient**:

# PROPERTIES OF MAX. LIKELIHOOD EST.

1. The MLE is **consistent**:

$$\hat{\theta} \xrightarrow{P} \theta$$

2. The MLE is **equivariant**:

If  $\hat{\theta}$  is MLE for  $\theta$  than  $g(\hat{\theta})$  is MLE for  $g(\theta)$

3. MLE is **asymptotically efficient**:  
roughly speaking, among well-behaved estimators, it has the smallest variance, at least for large samples.