

# PROBABILITY & STATISTICS

Lecture 9 – Intro to Statistics, Parameter estimation

# STATISTICS

**LET'S STRAT!**



# WHAT IS STATISTICS?

- How do *you* imagine Statistics?

# WHAT IS STATISTICS?

- How do *you* imagine Statistics?
  - Can you think of any example of Statistics?

# WE'RE ALL FOLLOWING THESE STATISTICS...

## Coronavirus Statistics

Global

59229918  
Infected

1397583 2.4%  
Deaths

40976963 69.2%  
Recovered

Last update: 23.11.20 17:03 GMT

Show 10 countries

Search:

Country	Infected	Deaths	Recovered	Deaths Percent	Recovered Percent	Infected per million	Deaths per million
USA	12603019	262781	7453682	2.1%	59.1%	37987	792
India	9170825	134088	8592303	1.5%	93.7%	6620	97
Brazil	6073058	169213	5432505	2.8%	89.5%	28491	794
France	2140208	48732	149521	2.3%	7.0%	32760	746
Russia	2114502	36540	1611445	1.7%	76.2%	14487	250
Spain	1589219	42619	N/A	2.7%	0.0%	33985	911
UK	1527495	55230	N/A	3.6%	0.0%	22454	812
Italy	1431795	50453	584493	3.5%	40.8%	23695	835
Argentina	1370366	37002	1195492	2.7%	87.2%	30212	816
Colombia	1248417	35287	1150932	2.8%	92.2%	24433	691

Showing 1 to 10 of 209 countries

Previous 1 2 3 4 5 ... 21 Next

### ACTIVE CASES

16,862,071


Currently Infected Patients

16,758,720 (99.4%) in Mild Condition  
103,351 (0.6%) Serious or Critical

[Show Graph](#)

Percent of infected, deaths and recovered (Global)


Infected Deaths Recovered



linear logarithmic

### Daily New Cases

Cases per Day  
Data as of 0:00 GMT+0




7-day moving average

daily linear logarithmic

### Daily Deaths

Deaths per Day  
Data as of 0:00 GMT+0



7-day moving average

# WHAT IS STATISTICS?

- How do *you* imagine Statistics?
  - Can you think of any example of Statistics?
- What is it?

# WHAT IS STATISTICS?

- How do *you* imagine Statistics?
  - Can you think of any example of Statistics?
- What is it?
- Why do we need it?



# WHAT IS STATISTICS?

- Statistics is a collection of methods which help us to describe, summarize, interpret and analyse data.


# WHAT IS STATISTICS?

- Statistics is a collection of methods which help us to describe, summarize, interpret and analyse data.
- Vital in research, politics, management, business...

# WHAT IS STATISTICS?


- Statistics is a collection of methods which help us to describe, summarize, interpret and analyse data.
- Vital in research, politics, management, business...
- There are different kinds of Statistics...

# STATISTICS: EXAMPLE 1



Source: <https://www.statista.com/chart/4467/female-employees-at-tech-companies/>


# STATISTICS: EXAMPLE 2



Source: <https://www.statista.com/chart/20018/canada-most-comfortable-with-female-leaders/>


# WHAT'S THE DIFFERENCE BETWEEN THE TWO?

## EXAMPLE 1



Source: <https://www.statista.com/chart/4467/female-employees-at-tech-companies/>

## EXAMPLE 2




Source: <https://www.statista.com/chart/20018/canada-most-comfortable-with-female-leaders/>

**WATCH THE VIDEO:**

<https://bit.ly/3fy8nzd>


# WHAT'S THE DIFFERENCE BETWEEN THE TWO?

## EXAMPLE 1



Source: <https://www.statista.com/chart/4467/female-employees-at-tech-companies/>

## EXAMPLE 2




Source: <https://www.statista.com/chart/20018/canada-most-comfortable-with-female-leaders/>




# WHAT'S THE DIFFERENCE BETWEEN THE TWO?

## DESCRIPTIVE STATISTICS



Source: <https://www.statista.com/chart/4467/female-employees-at-tech-companies/>


## INFERENTIAL STATISTICS



Source: <https://www.statista.com/chart/20018/canada-most-comfortable-with-female-leaders/>


# DESCRIPTIVE VS INFERENTIAL STATISTICS

## DESCRIPTIVE STATISTICS



- *Describe the data at hand.*

## INFERENTIAL STATISTICS



- *From the data at hand, make conclusions about a larger group.*

# SO...

- DESCRIPTIVE STATISTICS

- Data about **the whole population** is available.
- Summarize it with
  - *summary statistics*;
  - *tables*;
  - *plots*.

- INFERENCE STATISTICS

- Data about **a sample of the whole population** is available.
- Reason about the whole population.

# SO...

*Explorative Data Analysis (EDA) in ML, Business Analytics, ...*

- DESCRIPTIVE STATISTICS

- Data about **the whole population** is available.
- Summarize it with
  - *summary statistics;*
  - *tables;*
  - *plots.*

- INFERENCE STATISTICS

- Data about **a sample of the whole population** is available.
- Reason about the whole population.

# OUR FOCUS

- DESCRIPTIVE STATISTICS

- Data about **the whole population** is available.
- Summarize it with
  - *summary statistics*;
  - *tables*;
  - *plots*.

- INFERENCE STATISTICS

- Data about **a sample of the whole population** is available.
- Reason about the whole population.



# Parameter estimation

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>



# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*
- **How many men and women are there in the world?**

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>

# MEN AND WOMEN

- *If we take a random person  $X$ , what's the probability that it's a man?*
- **How many men and women are there in the world?**

<b>X</b>	1	0
<b>P(X)</b>	<b>p</b>	<b>1-p</b>

- There is precise data! 😊

World Male Population  
3,720,696  
thousands

World Female Population  
3,659,101  
thousands


<https://knoema.com/gndibag/sex-ratio-around-the-world-men-per-100-women-1950-2100>

# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- How many men and women are there in the world?
- There is precise data! 😊



↓

**p = 0.5**


# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?

- There is precise data! 😊



↓

**p = 0.5**

# MEN AND WOMEN


- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?

$$N_{\text{men}} = X_1 + \dots + X_{100} \sim \text{Bi}(100, \mathbf{p})$$

- There is precise data! 😊



↓

$$\mathbf{p} = 0.5$$

# MEN AND WOMEN

- If we take a random person  $X$ , what's the probability that it's a man?
- How many men and women are there in the world?

<b>X</b>	1	0
<b>P(X)</b>	<b>p = 0.5</b>	<b>1-p = 0.5</b>

- If we take 100 random people  $X_1 \dots X_{100}$ , how many men do we expect to see?

$$N_{\text{men}} = X_1 + \dots + X_{100} \sim \text{Bi}(100, \mathbf{p})$$

$$E(N_{\text{men}}) = 100 * \mathbf{p} = 50$$

- There is precise data! 😊



↓

$$\mathbf{p} = 0.5$$

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$

$$P(X = 1) = \mathbf{p} = ?$$



# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*
- **How many vegetarians are there in the world?**

$$X \sim \text{Bernoulli}(\mathbf{p})$$

$$P(X = 1) = \mathbf{p} = ?$$

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*
- **How many vegetarians are there in the world?**
- **No precise data!** 😞

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

*What would you do?*

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*
- **How many vegetarians are there in the world?**
- **No precise data!** 😞

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

- *What is a good guess for the value of  $\mathbf{p}$ ?*

- **How many vegetarians are there in the world?**

- **No precise data!** 😞

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

- *If we take a random person  $X$ , what's the probability that it's a vegetarian?*

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

- *What is a good guess for the value of  $\mathbf{p}$ ?*

$$\hat{\mathbf{p}} = 13 / 100 = 0.13$$

- **How many vegetarians are there in the world?**

- **No precise data!** 😞

*What would you do?*

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

# (NON-)VEGETARIANS

## INFERENCE STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?
- How many vegetarians are there in the world?
- No precise data! 😞

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

What would you do?

- What is a good guess for the value of  $\mathbf{p}$ ?

$$\hat{\mathbf{p}} = 13 / 100 = 0.13$$

...get the parameter estimate

- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?

## (NON-)VEGETARIANS

### INFERENCE STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?
- **How many vegetarians are there in the world?**

$$X \sim \text{Bernoulli}(p)$$
$$P(X = 1) = p = ?$$

- No precise data! ☹️

What would you do?

- What is a good guess for the value of  $p$ ?

$$\hat{p} = 13 / 100 = 0.13$$

- Survey:

- 100 people:  $X_1 \dots X_{100}$
- 13 are vegetarians

...get the parameter estimate

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?

## (NON-)VEGETARIANS

### INFERENCE STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?
- **How many vegetarians are there in the world?**
- **No precise data!** ☹️

$$X \sim \text{Bernoulli}(\mathbf{p})$$
$$P(X = 1) = \mathbf{p} = ?$$

What would you do?

- What is a good guess for the value of  $p$ ?

$$\hat{p} = 13 / 100 = 0.13$$

- Survey:

- 100 people:  $X_1 \dots X_{100}$
- 13 are vegetarians

...get the parameter estimate

From the data...



# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?
- Would it be better if there were more people in the survey?

**(NON-)VEGETARIANS**

INFERENTIAL STATISTICS

- If we take a random person  $X$ , what's the probability that it's a vegetarian?  
 $X \sim \text{Bernoulli}(p)$   
 $P(X = 1) = p = ?$
- How many vegetarians are there in the world?
  - No precise data! ☹️
  - What would you do?
- What is a good guess for the value of  $p$ ?
  - Survey:
    - 100 people:  $X_1 \dots X_{100}$
    - 13 are vegetarians

$\hat{p} = 13 / 100 = 0.13$

...get the parameter estimate

From the data...

# SOME QUESTIONS TO ASK OURSELVES

- Was it a good way to obtain an estimate?
- How precise is our estimate?
- Would it be better if there were more people in the survey?

## (NON-)VEGETARIANS

**INFERENCE STATISTICS**

- If we take a random person  $X$ , what's the probability that it's a vegetarian?  
 $X \sim \text{Bernoulli}(p)$   
 $P(X = 1) = p = ?$
- How many vegetarians are there in the world?
  - No precise data! ☹️
  - What would you do?
- What is a good guess for the value of  $p$ ?  
 $\hat{p} = 13 / 100 = 0.13$
- Survey:
  - 100 people:  $X_1 \dots X_{100}$
  - 13 are vegetarians

...get the parameter estimate

From the data...

# THE METHOD OF MAXIMUM LIKELIHOOD

A general framework for parameter estimation

# THE METHOD OF MAXIMUM LIKELIHOOD

A general framework for parameter estimation

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\boldsymbol{p})$  – whether a person is a vegetarian (1) or not (0).
- $\boldsymbol{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\boldsymbol{p}$ ?

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\mathbf{p})$  – whether a person is a vegetarian (1) or not (0).
- $\mathbf{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\mathbf{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$



# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\mathbf{p})$  – whether a person is a vegetarian (1) or not (0).
- $\mathbf{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\mathbf{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid \mathbf{p}) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(\mathbf{p})$  – whether a person is a vegetarian (1) or not (0).
- $\mathbf{p}$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $\mathbf{p}$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid \mathbf{p}) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid \mathbf{p}) * \dots * P(X_{13} = 1 \mid \mathbf{p}) * P(X_{14} = 0 \mid \mathbf{p}) * \dots * P(X_{100} = 0 \mid \mathbf{p}) =$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(p)$  – whether a person is a vegetarian (1) or not (0).
- $p$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $p$ ?

$$P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid p) * \dots * P(X_{13} = 1 \mid p) * P(X_{14} = 0 \mid p) * \dots * P(X_{100} = 0 \mid p) =$$

$$= p^{13}(1-p)^{87}$$

# LIKELIHOOD FUNCTION

- $X \sim \text{Bernoulli}(p)$  – whether a person is a vegetarian (1) or not (0).
- $p$  - the true proportion of the vegetarians.
- Survey results: 13 out of 100 participants are vegetarians:

$$X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0$$

- What is the probability to obtain such results given  $p$ ? **Likelihood:**

$$L(p) = P(X_1 = 1, \dots, X_{13} = 1, X_{14} = 0, \dots, X_{100} = 0 \mid p) =$$

*$\{X_1, \dots, X_{100} \text{ are independent}\}$*

$$= P(X_1 = 1 \mid p) * \dots * P(X_{13} = 1 \mid p) * P(X_{14} = 0 \mid p) * \dots * P(X_{100} = 0 \mid p) =$$

$$= p^{13}(1-p)^{87}$$

**Likelihood function** is the joint probability of realized sample given the parameters.

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?

# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?
  - We want the data that we obtained to be the most likely one.



# MAXIMUM LIKELIHOOD

- The probability to observe such data given  $p$  is:

$$L(p) = p^{13}(1-p)^{87}$$

- How to choose the best  $p$ ?
  - We want the data that we obtained to be the most likely one.
  - **Maximize  $L(p)$  w.r.t.  $p$ !**

# CALCULUS 101

- **How to optimize a function?**

# CALCULUS 101

- **How to optimize a function?**
  1. Compute its derivative.
  2. Set it to zero.
  3. Get the critical point(s).
  4. Chose the point of maximum.

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1 - x} = 0$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1 - x} = 0$$

Critical points:

$$x = 0, \quad x = 1, \quad x = \frac{1}{2}$$

# CALCULUS 101

- **How to optimize a function?**

1. Compute its derivative.
2. Set it to zero.
3. Get the critical point(s).
4. Chose the point of maximum.

$$f(x) = \log x + \log(1 - x)$$

$$\frac{d}{dx} f(x) = \frac{1}{x} - \frac{1}{1 - x} = 0$$

Critical points:

$$x = 0, \quad x = 1, \quad x = \frac{1}{2}$$



# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$




# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$f(x) = x*(1-x)$$



**log f(x)**

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) =$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) = \frac{13}{p} - \frac{87}{1-p} = 0$$

# MAXIMUM LIKELIHOOD

$$\text{maximize } L(p) = p^{13}(1-p)^{87} \quad \text{w.r.t. } p$$

**USEFUL TRICK: MAXIMIZE LOG-LIKELIHOOD INSTEAD**

$$\text{maximize } \log L(p) = 13 \log(p) + 87 \log(1-p) \quad \text{w.r.t. } p$$

$$\frac{d}{dp} \log L(p) = \frac{13}{p} - \frac{87}{1-p} = 0$$

$$\hat{p} = \frac{13}{100}$$

# MAXIMUM LIKELIHOOD ESTIMATE

1. Write down the likelihood function:

$$L(\theta) = P(X_1, \dots, X_n \mid \theta) = \prod_{i=1}^n P(X_i \mid \theta)$$

2. Find its maximum w.r.t. the unknown parameter  $\theta$ :

$$\hat{\theta} = \operatorname{argmax}_{\theta} L(\theta) \text{ w.r.t. } \theta$$

(!) In many cases, it's easier to maximize **log-likelihood**:

$$\log L(\theta) = \log \prod_{i=1}^n P(X_i \mid \theta) = \sum_{i=1}^n \log P(X_i \mid \theta)$$

$$\hat{\theta} = \operatorname{argmax}_{\theta} \log L(\theta)$$

**Maximum Likelihood Estimate (MLE)** is the value which

...



**Maximum Likelihood Estimate (MLE)** is the value which maximizes the probability of observing the realized sample.

# **MLE FOR SOME DISCRETE DISTRIBUTIONS**

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

- *(General case of the previous example)*
- You want to estimate the proportion of the vegetarians.

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.
- (General case of the previous example)
- You want to estimate the proportion of the vegetarians.
- Survey: you ask  $N$  people whether they are vegetarians and get responses

$$X_1, X_2, \dots, X_N$$

# BERNOULLI DISTRIBUTION

$$X \sim \text{Bernoulli}(p)$$

$$E(X) = p$$

$$\text{Var}(X) = p(p-1)$$

- Models the probability of success in an experiment with two outcomes.

- *(General case of the previous example)*
- You want to estimate the proportion of the vegetarians.
- Survey: you ask  $N$  people whether they are vegetarians and get responses

$$X_1, X_2, \dots, X_N$$

- What's the MLE of  $p$ ?

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$



# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1 - p)^{1 - X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} =$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1-p} = 0$$

$$\hat{p} = \frac{1}{N} \sum_{i=1}^N X_i$$

# MLE FOR BERNOULLI DISTRIBUTION

$$L(p) = \prod_{i=1}^N p^{X_i} (1-p)^{1-X_i}$$

$$\log L(p) = \sum_{i=1}^N X_i \log p + (1 - X_i) \log(1 - p)$$

$$\frac{d}{dp} \log L(p) = \frac{\sum_{i=1}^N X_i}{p} - \frac{N - \sum_{i=1}^N X_i}{1-p} = 0$$

$$\hat{p} = \frac{1}{N} \sum_{i=1}^N X_i$$


# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.



# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Suppose you want to model the number of phone calls a call center receives in an hour.

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.



# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Suppose you want to model the number of phone calls a call center receives in an hour.
- You've been recording it for the past  $n$  hours:

$X_1, X_2, \dots, X_n$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.

# POISSON DISTRIBUTION

$$X \sim \text{Po}(\lambda), \quad \lambda > 0$$

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$E(X) = \lambda, \quad \text{Var}(X) = \lambda$$

- Models number of events occurring in a fixed interval of time.
- Assumptions: events occur
  - with a known constant mean rate;
  - independently of the time since the last event.

- Suppose you want to model the number of phone calls a call center receives in an hour.
- You've been recording it for the past  $n$  hours:

$$X_1, X_2, \dots, X_n$$

- What's the MLE of  $\lambda$ ?

# MLE FOR POISSON DISTRIBUTION

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) =$$

# MLE FOR POISSON DISTRIBUTION

$$P(X = k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) =$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) =$$



# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0,$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0, \quad \hat{\lambda} = \frac{1}{n} \sum_{i=1}^n X_i$$

# MLE FOR POISSON DISTRIBUTION

$$L(\lambda) = P(X_1, \dots, X_n | \lambda) = \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{X_i}}{X_i!}$$

$$\log L(\lambda) = \sum_{i=1}^n [-\lambda \log e + X_i \log \lambda - \log X_i!]$$

$$= -n\lambda + \log \lambda \sum_{i=1}^n X_i - \sum_{i=1}^n \log X_i!$$

$$\frac{d}{d\lambda} \log L(\lambda) = -n + \frac{\sum_{i=1}^n X_i}{\lambda} = 0,$$

$$\hat{\lambda} = \frac{1}{n} \sum_{i=1}^n X_i$$

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	?	?	?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	?	?	?

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:

<b><math>X</math></b>	1	2	3
<b><math>P(X)</math></b>	$p$	$q$	$1-p-q$

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:
- We observed it  $N$  times:

$X$	1	2	3
$P(X)$	$p$	$q$	$1-p-q$

$X$	1	2	3
#	$n_1$	$n_2$	$n_3$

- How many parameters does such a distribution have?

# MLE: YET ANOTHER EXAMPLE

- Random variable  $X$  takes three values with unknown probabilities:
- We observed it  $N$  times:

<b>X</b>	1	2	3
<b>P(X)</b>	$p$	$q$	$1-p-q$

<b>X</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- How many parameters does such a distribution have?
- What is the MLE of the parameters?



# CALCULUS 101

- **How to optimize a function of several variables?**

# CALCULUS 101

- **How to optimize a function of several variables?**

1. Compute partial derivatives.
2. Set them to zero.
3. Solve the equations and get the critical points.

# CALCULUS 101

- **How to optimize a function of several variables?**

1. Compute partial derivatives.
2. Set them to zero.
3. Solve the equations and get the critical points.

$$f(x, y) = x^2 + 2xy - 2x - 4y$$

$$\frac{d}{dx}f(x, y) = 2x + 2y - 2 = 0$$

$$\frac{d}{dy}f(x, y) = 2x - 4 = 0$$

$$x^* = 2, \quad y^* = 1$$

# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3
<b>P(X)</b>	$p$	$q$	$1-p-q$

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) =$$



# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$



# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$



$$\log L(p, q) =$$

# MLE: YET ANOTHER EXAMPLE

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

- What's the probability of observing such data? Likelihood:

$$L(p, q) = p^{n_1} \cdot q^{n_2} \cdot (1 - p - q)^{n_3}$$

maximize  $L(p, q)$  w.r.t.  $p, q$



$$\log L(p, q) = n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$$

maximize  $\log L(p, q)$  w.r.t.  $p, q$

# MLE: EXAMPLE 2

<b>X</b>	1	2	3	<b>Value</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>	<b>#</b>	$n_1$	$n_2$	$n_3$

maximize  $n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$  w.r.t.  $p, q$

$$\frac{d}{dp} \log L(p, q) = \frac{n_1}{p} - \frac{n_3}{1 - p - q} = 0$$

$$\frac{d}{dq} \log L(p, q) = \frac{n_2}{q} - \frac{n_3}{1 - p - q} = 0$$



## MLE: EXAMPLE 2


<b>X</b>	1	2	3
<b>P(X)</b>	<b>p</b>	<b>q</b>	<b>1-p-q</b>

<b>Value</b>	1	2	3
<b>#</b>	$n_1$	$n_2$	$n_3$

maximize  $n_1 \log p + n_2 \log q + n_3 \log(1 - p - q)$  w.r.t.  $p, q$

$$\frac{d}{dp} \log L(p, q) = \frac{n_1}{p} - \frac{n_3}{1 - p - q} = 0$$

$$\frac{d}{dq} \log L(p, q) = \frac{n_2}{q} - \frac{n_3}{1 - p - q} = 0$$



$$\hat{p} = \frac{n_1}{N}$$
$$\hat{q} = \frac{n_2}{N}$$

# TO SUM UP

- **Likelihood function** is the joint probability of realized sample given the parameters.
- **Maximum Likelihood Estimate (MLE)** is the value which maximizes the probability of observing the realized sample.

# RANDOMIZED RESPONSE

*Asking embarrassing questions*

# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.

# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.
- BUT WHAT IF the question is very sensitive?
  - People won't tell the truth.

# MOTIVATION

- BEFORE:
  - How many vegetarians are in the world?
  - Survey: ask “*are you a vegetarian?*”, estimate the true proportion.
- BUT WHAT IF the question is very sensitive?
  - People won't tell the truth.
- EXAMPLE: *Do you find this course boring?*
  - How do I find out what my students *actually* think?

# LET'S TRY THIS OUT!

- Toss a coin...

# LET'S TRY THIS OUT!

- Toss a coin...
- Now, answer one of the following questions:



# LET'S TRY THIS OUT!

- Toss a coin...
- Now, answer one of the following questions:

<b>If you got HEADS:</b>	DO YOU FIND THIS CLASS BORING?
<b>If you got TAILS:</b>	ARE YOU AT THE STATISTICS CLASS RIGHT NOW?


# BINOMIAL DISTRIBUTION

$$X \sim \text{Bi}(n, p), \quad n = 1, 2, \dots, \quad 0 < p < 1$$

$$P(X = k) = C_n^k p^k (1 - p)^{n-k}$$

$$E(X) = np, \quad \text{Var}(X) = np(1-p)$$

- Models the number of successes in a series of  $n$  independent Bernoulli trials, each of which has a success probability  $p$ .



# BINOMIAL DISTRIBUTION

$$X \sim \text{Bi}(n, p), \quad n = 1, 2, \dots, \quad 0 < p < 1$$

$$P(X = k) = C_n^k p^k (1 - p)^{n-k}$$

$$E(X) = np, \quad \text{Var}(X) = np(1-p)$$

- Models the number of successes in a series of  $n$  independent Bernoulli trials, each of which has a success probability  $p$ .

- You sell two types of sandwiches: chicken and vegetarian. Which one people like more?

- For the past  $N$  days, you were selling  $n=100$  sandwiches every day and recorded the number of the chicken ones:

$$X_1, X_2, \dots, X_N$$

- What is the MLE of the  $p$  parameter?

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) =$$



# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = 0$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} =$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{\sum_{i=1}^N X_i}{100N}$$

# MLE FOR BINOMIAL DISTRIBUTION

$$L(p) = \prod_{i=1}^N C_n^{X_i} p^{X_i} (1 - p)^{100 - X_i}$$

$$\log L(p) = \sum_{i=1}^N [\log C_n^{X_i} + X_i \log p + (100 - X_i) \log(1 - p)]$$

$$\frac{d}{dp} L(p) = \sum_{i=1}^N \frac{X_i}{p} - \frac{100 - X_i}{1 - p} = \frac{\sum_{i=1}^N X_i}{p} - \frac{100N - \sum_{i=1}^N X_i}{1 - p} = 0$$

$$\hat{p} = \frac{\sum_{i=1}^N X_i}{100N}$$

# TO SUM UP

- **Likelihood function** is the joint probability of realized sample given the parameters.
- **Maximum Likelihood Estimate (MLE)** is the value which maximizes the probability of observing the realized sample.

# MAXIMUM LIKELIHOOD ESTIMATE

## FOR PARAMETERS OF CONTINUOUS DISTRIBUTIONS

# LIKELIHOOD FUNCTION

- Given samples from a distribution:  $X_1, X_2, \dots, X_N$   
Obtain estimate of the parameter(s)  $\theta$



# LIKELIHOOD FUNCTION

- Given samples from a distribution:  $X_1, X_2, \dots, X_N$   
Obtain estimate of the parameter(s)  $\theta$
- **Likelihood function:**

# LIKELIHOOD FUNCTION

- Given samples from a distribution:  $X_1, X_2, \dots, X_N$   
Obtain estimate of the parameter(s)  $\theta$
- **Likelihood function:** joint *density function* of the obtained data given the model parameters

$$L(\theta) = p(X_1, X_2, \dots, X_n | \theta)$$

# LIKELIHOOD FUNCTION

- Given samples from a distribution:  $X_1, X_2, \dots, X_N$   
Obtain estimate of the parameter(s)  $\theta$
- **Likelihood function:** joint *density function* of the obtained data given the model parameters

$$L(\theta) = p(X_1, X_2, \dots, X_n | \theta)$$

- Assuming the independence of the samples:

# LIKELIHOOD FUNCTION

- Given samples from a distribution:  $X_1, X_2, \dots, X_N$   
Obtain estimate of the parameter(s)  $\theta$
- **Likelihood function:** joint *density function* of the obtained data given the model parameters

$$L(\theta) = p(X_1, X_2, \dots, X_n | \theta)$$

- Assuming the independence of the samples:

$$L(\theta) = \prod_{i=1}^n p(X_i | \theta)$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) =$$

$$p(x) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} =$$

$$p(x) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$



# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

maximize  $L(\lambda)$  w.r.t.  $\lambda$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

maximize  $L(\lambda)$  w.r.t.  $\lambda$

$$\log L(\lambda) =$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

$$\text{maximize } L(\lambda) \quad \text{w.r.t. } \lambda$$

$$\log L(\lambda) = n \cdot \log \lambda - \lambda \sum_{i=1}^N X_i$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

maximize  $L(\lambda)$  w.r.t.  $\lambda$

$$\log L(\lambda) = n \cdot \log \lambda - \lambda \sum_{i=1}^N X_i$$

$$\frac{d}{d\lambda} \log L(\lambda) =$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

maximize  $L(\lambda)$  w.r.t.  $\lambda$

$$\log L(\lambda) = n \cdot \log \lambda - \lambda \sum_{i=1}^N X_i$$

$$\frac{d}{d\lambda} \log L(\lambda) = \frac{n}{\lambda} - \sum_{i=1}^N X_i = 0$$

# ML EXAMPLE: EXPONENTIAL DISTRIBUTION

- Observed data:  $X_1, X_2, \dots, X_N$
- Likelihood:

$$L(\lambda) = p(X_1, \dots, X_N | \lambda) = \prod_{i=1}^N \lambda \cdot e^{-\lambda X_i} = \lambda^n \cdot \prod_{i=1}^N e^{-\lambda X_i}$$

maximize  $L(\lambda)$  w.r.t.  $\lambda$

$$\log L(\lambda) = n \cdot \log \lambda - \lambda \sum_{i=1}^N X_i$$

$$\frac{d}{d\lambda} \log L(\lambda) = \frac{n}{\lambda} - \sum_{i=1}^N X_i = 0 \quad \Rightarrow \quad \hat{\lambda} = \frac{1}{\frac{1}{N} \sum_{i=1}^N X_i}$$